

One Particle (improper) states and Scattering States in Nelson's Massless Model

Alessandro Pizzo*

S.I.S.S.A. -I.S.A.S., via Beirut 2-4 34014 Trieste, Italia

October 25, 2000

Abstract

In the one nucleon sector of Nelson's massless model with an ultraviolet cut-off and no infrared regularization, one particle improper states are constructed and a scattering theory is developed both for the massless scalar field and for the non-relativistic particle. One particle improper states are obtained constructively by iterating an analytic perturbation of isolated eigenvalues. Scattering states are constructed by exploiting some properties of (non-relativistic) locality of the model and starting from "minimal asymptotic nucleon states". They represent asymptotic configurations where a cloud of (soft) mesons always appears even if their energy may be arbitrarily small.

*Pizzo@sissa.it

Introduction

The model I will examine describes the covariant interaction (under spatial translation) between mesons and a non-relativistic nucleon. The rigorous study of the model was started by Nelson [1], in order to remove the ultraviolet cut-off in the interaction, in the case of massive mesons.

Then, it was used as a toy model to give a consistent explanation of interaction between radiation and matter, in particular of a single charged particle with radiation field.

The most important results on the subject are contained in [2] . They regard both the case of massive mesons and the massless case. In particular they show the existence of a subspace of one particle states for the hamiltonian (it means that the hamiltonian acts on them as a function of the total impulse) in the massive case and the absence of such a subspace in the other case (if there is no infrared regularization in the interaction term). Recently [3] the model was studied in order to prove asymptotic completeness, adding some more conditions (confinement of the non-relativistic particle).

In this paper I study the interaction of a non-relativistic nucleon with massless mesons, without infrared regularization. The aim is to prove that a description of the asymptotic dynamics exists once the ultraviolet cut-off is fixed in the interaction and for sufficiently small values of the coupling constant; the result is a scattering theory in a physical relevant subspace with an explicit interpretation with respect to the asymptotic dynamical variables.

The interaction without infrared regularization involves additional difficulties with respect to previous treated versions:

- the absence of a ground state for the hamiltonians at fixed total momentum;
- the massless dispersion of smearing test functions in L.S.Z. meson field operators. These operators have to define a free meson field in the asymptotic limit.

My approach to scattering is different from the traditional one in two aspects:

- the first one consists in using some mechanism of the scattering in quantum field theory (Haag-Ruelle formulation), by exploiting the locality properties at fixed time of the model;

- the second one is technical and regards the determination of the limit, for $\sigma \rightarrow 0$, of the ground states of the hamiltonians with an infrared cut-off σ , at fixed total momentum and properly transformed. I will use an iterative procedure (different from the operatorial renormalization group [4] by Bach, Froehlich and Segal) which gives a strong convergence with an error estimable in terms of the infrared cut-off that we have to remove.

Finally, I observe that some of these ideas and techniques can be used in the study of scattering in the one-electron sector of non relativistic quantum electrodynamics.

Definition of the model.

The system consists of a non-relativistic spinless quantum particle of mass m , which is coupled to the massless boson field. The non-relativistic particle is described by position and momentum variables with usual canonic commutation rules (c.c.r.)

$$[x_i, p_j] = i\delta_{i,j} \ (\hbar = 1) ;$$

the meson field is described by $\mathbf{A}(0, \mathbf{y}) = \frac{1}{\sqrt{2\pi^3}} \cdot \int \left(a^\dagger(\mathbf{k}) e^{-i\mathbf{k}\cdot\mathbf{y}} + a(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{y}} \right) \frac{d^3k}{\sqrt{2|\mathbf{k}|}}$ ($c = 1$), where $a^\dagger(\mathbf{k})$, $a(\mathbf{k})$ are creation and destruction operator valued distributions which satisfy the c.c.r.

$$\begin{aligned} [a(\mathbf{k}), a(\mathbf{q})] &= [a^\dagger(\mathbf{k}), a^\dagger(\mathbf{q})] = 0 \\ [a(\mathbf{k}), a^\dagger(\mathbf{q})] &= \delta^3(\mathbf{k} - \mathbf{q}) \end{aligned}$$

The spatial translations are implemented by the total momentum

$$\mathbf{P} = \mathbf{p} + \mathbf{P}^{e.m.} = \mathbf{p} + \int \mathbf{k} a^\dagger(\mathbf{k}) a(\mathbf{k}) d^3k ;$$

the time evolution is given by the covariant hamiltonian ($[H, \mathbf{P}] = 0$)

$$H = \frac{\mathbf{p}^2}{2m} + g \int_0^\kappa \left(a(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{x}} + a^\dagger(\mathbf{k}) e^{-i\mathbf{k}\cdot\mathbf{x}} \right) \frac{d^3\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{1}{2}}} + H^{mes}$$

where κ is the ultraviolet *cut-off* and g is the coupling constant.

The Hilbert space of the system is $\mathcal{H} = L^2(R^3) \otimes F$, where F is the Fock space

with respect to the operator valued distributions $\{a^\dagger(\mathbf{k}), a(\mathbf{k})\}$. An element of H is a sequence $\{\psi^n\}$ of functions on R^{3n+1} with $\|\psi\| < \infty$, where $\|\psi\|^2 = \sum_{n=0}^{\infty} \int \overline{\psi^n(\mathbf{x}, \mathbf{k}_1, \dots, \mathbf{k}_n)} \psi^n(\mathbf{x}, \mathbf{k}_1, \dots, \mathbf{k}_n) d^3k_1 \dots d^3k_n d^3x$ and each $\psi^n(\mathbf{x}, \mathbf{k}_1, \dots, \mathbf{k}_n)$ is symmetric in $\mathbf{k}_1, \dots, \mathbf{k}_n$. The $n = 0$ component corresponds to the vacuum subspace tensorized with the non-relativistic particle space $L^2(R^3)$.

Standard results about H and P :

i) The operators $\mathbf{P} = \mathbf{p} \otimes 1 + 1 \otimes \int \mathbf{k} a^\dagger(\mathbf{k}) a(\mathbf{k}) d^3k$ are essentially self-adjoint (e.s.a.) in $D \equiv \bigvee_{n \in \mathbb{N}} h \otimes \psi^n$, which is the set of finite linear combinations of vectors of wave function $h(\mathbf{x}) \psi^n(\mathbf{k}_1, \dots, \mathbf{k}_n)$, where $h(\mathbf{x}) \in S(R^3)$ and where $\psi^n(\mathbf{k}_1, \dots, \mathbf{k}_n) \in S(R^{3n})$ *simultaneously*, $n \in \mathbb{N}$. Since $\mathbf{p}, \int \mathbf{k} a^\dagger(\mathbf{k}) a(\mathbf{k}) d^3k$ are e.s.a. respectively in $S(R^3)$ and in $\bigvee_{n \in \mathbb{N}} \psi^n$, the result follows for $\mathbf{P}$ operators.

ii) The interaction term in the hamiltonian is a Kato's small perturbation with respect to $H_0 \equiv \frac{\mathbf{p}^2}{2m} + H^{mes}$; therefore the hamiltonian H is e.s.a. in $D \equiv \bigvee_{n \in \mathbb{N}} h \otimes \psi^n$ and $D(H) \equiv D(H_0)$.

iii) The groups $e^{i\mathbf{a} \cdot \mathbf{P}} e^{i\tau H}$ ($\tau, a^i \in R$) commute.

iv) The joined spectral decomposition of the space H , with respect to $\mathbf{P}$ operators, is written $H = \bigoplus \int H_{\mathbf{P}} d^3P$, where $H_{\mathbf{P}}$ is isomorphic to F .

In fact, to the improper eigenvectors (of the $\mathbf{P}$ operators) $\psi_{\mathbf{P}}^n$ of wave function

$$\psi_{\mathbf{P}}^n(\mathbf{x}, \mathbf{k}_1, \dots, \mathbf{k}_n) = (2\pi)^{\frac{-3}{2}} e^{i(\mathbf{P} - \mathbf{k}_1 - \dots - \mathbf{k}_n) \cdot \mathbf{x}} \psi_{\mathbf{P}}^n(\mathbf{k}_1, \dots, \mathbf{k}_n)$$

we can relate a natural scalar product:

$$(\phi_{\mathbf{P}}^n, \psi_{\mathbf{P}}^m) = \delta_{n,m} \int \overline{\phi_{\mathbf{P}}^n(\mathbf{k}_1, \dots, \mathbf{k}_n)} \psi_{\mathbf{P}}^m(\mathbf{k}_1, \dots, \mathbf{k}_n) d^3k_1 \dots d^3k_n \quad (0.1)$$

The vectorial space $\{\overline{\bigvee_{n \in \mathbb{N}} \psi_{\mathbf{P}}^n}\}$ is obtained as closure of the finite linear combinations of the $\psi_{\mathbf{P}}^n$, in the norm which derives from the scalar product (0.1). Starting from this space we uniquely define the linear application

$$I_{\mathbf{P}} : \bigvee_{n \in \mathbb{N}} \psi_{\mathbf{P}}^n \rightarrow F^b$$

by the prescription:

$$I_{\mathbf{P}}(\psi_{\mathbf{P}}^n(\mathbf{x}, \mathbf{k}_1, \dots, \mathbf{k}_n)) = \frac{1}{\sqrt{n!}} \int b^\dagger(\mathbf{k}_1) \dots b^\dagger(\mathbf{k}_n) \psi_{\mathbf{P}}^n(\mathbf{k}_1, \dots, \mathbf{k}_n) d^3k_1 \dots d^3k_n \psi_0$$

where $b(\mathbf{k}), b^\dagger(\mathbf{k})$ formally correspond to $a(\mathbf{k})e^{i\mathbf{k}\cdot\mathbf{x}}, a^\dagger(\mathbf{k})e^{-i\mathbf{k}\cdot\mathbf{x}}$. They are destruction and creation operator valued distributions in the Fock space $F^b \cong F$. The norm given by (0.1) for $\psi_{\mathbf{P}}^n$ is equal to $\|I_{\mathbf{P}}(\psi_{\mathbf{P}}^n)\|_F$ ($\|\cdot\|_F$ is the Fock norm).

Main results.

The results will be presented in two parts.

The first one regards the determination of the ground states $\psi_{\mathbf{P}}^{\sigma_j}$ of the hamiltonians

$$H_{\mathbf{P}, \sigma_j} = \frac{(\mathbf{P}^{mes})^2}{2m} - \frac{\mathbf{P} \cdot \mathbf{P}^{mes}}{m} + \frac{\mathbf{P}^2}{2m} + g \int_{\sigma_j}^\kappa (b(\mathbf{k}) + b^\dagger(\mathbf{k})) \frac{d^3\mathbf{k}}{\sqrt{2|\mathbf{k}|}} + H^{mes}$$

at fixed total momentum $\mathbf{P}$, restricted to the set $\Sigma \equiv \{\mathbf{P} : |\mathbf{P}| \leq \sqrt{m}\}$, where $\sigma_j = \epsilon^{\frac{j+1}{2}}$, $0 < \epsilon < \left(\frac{1}{4}\right)^{16}$ and $j \in \mathbb{N}$, is the infrared cut-off.

The main results are:

1) Theorem 1.5

$H_{\mathbf{P}, \epsilon\sqrt{\epsilon}}|_{F_{\epsilon\sqrt{\epsilon}}^+}$ has unique ground state $\psi_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}}$ with gap bigger than $\frac{\epsilon\sqrt{\epsilon}}{2}$; $\psi_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}}$ (unnormalized) is

$$\psi_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}} = -\frac{1}{2\pi i} \oint \frac{1}{H_{\mathbf{P}, \epsilon\sqrt{\epsilon}} - E} dE \psi_{\mathbf{P}}^\epsilon$$

where the integral is calculated around the point $E_{\mathbf{P}}^\epsilon$, $E \in \mathcal{C}$ and s.t. $|E - E_{\mathbf{P}}^\epsilon| = \frac{11\epsilon\sqrt{\epsilon}}{20}$

In its generalization to generic j , this result and **Corollary 1.6** allow to construct the sequence of ground states $\left\{\psi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}\right\}$ for sufficiently small g .

2) In **theorem 2.3** and **2.3bis**. the strong convergence of the sequence $\left\{\phi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}\right\}$ is proved for small coupling constant g , with an error $\left\|\phi_{\mathbf{P}} - \phi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}\right\| \leq \epsilon^{\frac{j+1}{16}}$, where $\phi_{\mathbf{P}} = s - \lim_{j \rightarrow \infty} \phi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}$. The vectors $\left\{\phi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}\right\}$ are the ground states of the transformed hamiltonians

$$H_{\mathbf{P},\epsilon}^w = W_{\epsilon}^{\frac{j+1}{2}} \left(\nabla E^{\epsilon \frac{j+1}{2}} (\mathbf{P}) \right) H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} W_{\epsilon}^{\frac{j+1}{2}} \left(\nabla E^{\epsilon \frac{j+1}{2}} (\mathbf{P}) \right)$$

$$\left(\text{where } W_{\epsilon}^{\frac{j+1}{2}} \left(\nabla E^{\epsilon \frac{j+1}{2}} (\mathbf{P}) \right) = e^{-g \int_{\epsilon}^{\kappa} \frac{j+1}{2} \frac{b(\mathbf{k}) - b^{\dagger}(\mathbf{k})}{|\mathbf{k}| \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}} (\mathbf{P}) \right)} \frac{d^3 k}{\sqrt{2} |\mathbf{k}|}} \right)$$

$\Leftrightarrow$

In the second part, I will treat the scattering. I will assume a technical hypothesis that is not proved in the spectral analysis:¹ there exists a positive constant m_r such that

$$\text{hypothesis B1} \quad \frac{\partial E^{\sigma}(\mathbf{P})}{\partial |\mathbf{P}|} \geq \frac{|\mathbf{P}|}{m_r} \quad \mathbf{e} \quad \frac{\partial^2 E^{\sigma}(\mathbf{P})}{\partial^2 |\mathbf{P}|} \geq \frac{1}{m_r} \quad \forall \sigma$$

The main result is the construction of the generic *minimal asymptotic nucleon state* $\psi_G^{out(in)}$ (G is the wave function, in variables $\mathbf{P}$, of the one-particle state from which the construction of the minimal asymptotic nucleon state starts). It is defined by the strong convergence of the approximating vector $\psi_G(t)$. This subject is discussed in the paragraph 4.2. The result is proved in **theorem 4.1**.

$$\exists C, \rho > 0 \quad \text{s.t.} \quad \|\psi_G(t) - \psi_G^{out}\| \leq \frac{C}{t^{\rho}}$$

From this result, the construction of scattering subspaces $H^{out(in)}$ and the definition of asymptotic dynamical variables easily follow:

theorem 5.2

The functions of nucleon mean velocity, continuous and of compact support, have asymptotically strong limits in H^{out} ; in particular:

$$s - \lim_{t \rightarrow +\infty} e^{iHt} f\left(\frac{\mathbf{x}}{t}\right) e^{-iHt} \psi_{G,\varphi}^{out} = \psi_{G,\widehat{f},\varphi}^{out} \quad \text{where} \quad \widehat{f}(\mathbf{P}) \equiv \lim_{\sigma \rightarrow 0} f(\nabla E^{\sigma}(\mathbf{P}))$$

corollary 5.3

In the spaces H^{out} , the asymptotic meson algebra $\mathcal{A}^{out}$ is defined as the norm closure of

¹ In this respect, I cite the result T.Chen communicated me: the function $E(\mathbf{P})$ has continuous 2^{nd} derivatives.

the $*$ algebra generated by the set of Weyl operators $\{W^{out}(\mu) : \tilde{\mu}(\mathbf{k}) \in C_0^\infty(R^3 \setminus 0)\}$:

$$W^{out}(\mu) = s - \lim_{t \rightarrow +\infty} e^{iHt} e^{-iH^{mes}t} e^{i(a(\mu) + a^\dagger(\mu))} e^{iH^{mes}t} e^{-iHt}$$

Part I

Spectral Analysis.

The iterative procedure aims at constructing the sequence $\{\psi_{\mathbf{P}}^{\sigma_j}\}$ of ground states of the hamiltonians $H_{\mathbf{P},\sigma_j}$

$$H_{\mathbf{P},\sigma_j} = \frac{(\mathbf{P}^{mes})^2}{2m} - \frac{\mathbf{P} \cdot \mathbf{P}^{mes}}{m} + \frac{\mathbf{P}^2}{2m} + g \int_{\sigma_j}^\kappa \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) \frac{d^3\mathbf{k}}{\sqrt{2|\mathbf{k}|}} + H^{mes}$$

where $\sigma_j = \epsilon^{\frac{j+1}{2}}$ $0 < \epsilon < \left(\frac{1}{4}\right)^{16}$ $j \in \mathbb{N}$.

The procedure starts from the comparisons between the resolvents of the hamiltonians $H_{\mathbf{P},\sigma_j}$ and $H_{\mathbf{P},\sigma_{j+1}}$. It recursively uses the Kato's theorem on analytic perturbation of isolated eigenvalues (of self-adjoint operators) to relate the ground states $\psi_{\mathbf{P}}^{\sigma_j}$ and $\psi_{\mathbf{P}}^{\sigma_{j+1}}$; at each step two pieces of information are required:

1) a lower bound for the gap (with respect to the ground eigenvalue) of the hamiltonian $H_{\mathbf{P},\sigma_j}$ restricted to the subspace

$$F_{\sigma_{j+1}}^+ \equiv \left\{ \overline{\bigvee_{n \in \mathbb{N}} \psi^n(\mathbf{k}_1, \dots, \mathbf{k}_n)} : n \in \mathbb{N}, \psi^n(\mathbf{k}_1, \dots, \mathbf{k}_n) \in L^2(R^{3n}) \text{ simm.}, |\mathbf{k}_i| \geq \sigma_{j+1} \ 1 \leq i \leq n \right\};$$

2) an estimate of the difference $\Delta H_{\mathbf{P}}|_{\sigma_{j+1}}^{\sigma_j} \equiv H_{\mathbf{P},\sigma_{j+1}}|_{F_{\sigma_{j+1}}^+} - H_{\mathbf{P},\sigma_j}|_{F_{\sigma_{j+1}}^+}$ between two subsequent infrared cut-off hamiltonians; this is small with respect to $H_{\mathbf{P},\sigma_j}|_{F_{\sigma_{j+1}}^+}$ in a generalized sense, which means that it is possible to expand the spectral projection of $H_{\mathbf{P},\sigma_{j+1}}|_{F_{\sigma_{j+1}}^+}$ on the ground eigenvalue in a perturbative series in terms of the resolvent of $H_{\mathbf{P},\sigma_j}|_{F_{\sigma_{j+1}}^+}$ and of the difference $\Delta H_{\mathbf{P}}|_{\sigma_{j+1}}^{\sigma_j}$.

The convergence of the sequence $\{\psi_{\mathbf{P}}^{\sigma_j}\}$ is a problem of perturbation of an eigenvalue in the continuum, precisely of the ground eigenvalue of the hamiltonian $H_{\mathbf{P}}^0 = \frac{(\mathbf{P}^{mes})^2}{2m} - \frac{\mathbf{P} \cdot \mathbf{P}^{mes}}{m} + \frac{\mathbf{P}^2}{2m} + H^{mes}$. If the exponent of $|\mathbf{k}|$ in the interaction term of

the hamiltonian were larger than $-\frac{1}{2}$, the norm estimates about resolvents would be sufficient not only to construct the sequence $\{\psi_{\mathbf{P}}^{\sigma_j}\}$ but also to gain the convergence. The physical case $-\frac{1}{2}$ is a limit case which requires inequivalent representations of the variables $\{b(\mathbf{k}), b^\dagger(\mathbf{k})\}$ at different $\mathbf{P}$, and strong estimates of the series expansion of the difference between two subsequent ground eigenvectors in order to have convergence.

In the case $\mathbf{P} = 0$ the right representation is explicitly known by symmetry. The only problem is to improve some estimates of some terms in the iterative procedure, in order to gain the convergence; in the case $\mathbf{P} \neq 0$, the representation problem is solved by transforming, step by step, the hamiltonian in a canonic form. Such canonic form is analogous, as regards the convergence problem, to the case $\mathbf{P} = 0$. In the limit we obtain the representation of $\{b(\mathbf{k}), b^\dagger(\mathbf{k})\}$ given by the non-Fock coherent transformation:

$$\mathcal{W}(\mathbf{P}) = e^{-g \int_0^\kappa \frac{b(\mathbf{k}) - b^\dagger(\mathbf{k})}{|\mathbf{k}|(1 - \mathbf{k} \cdot \nabla E(\mathbf{P}))} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}}$$

Since such representations of $\{b(\mathbf{k}), b^\dagger(\mathbf{k})\}$ are inequivalent for different $\mathbf{P}$, we have to face the problem of the existence of the vector $\int \phi_{\mathbf{P}} d^3 P$ where $\phi_{\mathbf{P}}$ is the limit in Fock space of the ground states of the transformed hamiltonians $\mathcal{W}_\sigma(\mathbf{P}) H_{\mathbf{P},\sigma} \mathcal{W}_\sigma^\dagger(\mathbf{P})$. The sufficient conditions to define $\int \phi_{\mathbf{P}} d^3 P$ are explored in chapter 3.

1 Construction of the sequence $\{\psi_{\mathbf{P}}^{\sigma_j}\}$.

In the present chapter I only construct the sequence. In order to do it, I introduce some preliminary lemmas (1.1, 1.2, 1.3, 1.4). They are necessary to perform the first step of the iterative procedure contained in theorem 1.5. In the results of theorem 1.5 there are the hypotheses to repeat the same procedure with a smaller infrared cut-off. Finally in corollary 1.6 the sequence of ground states $\{\psi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, j \in \mathbb{N}\}$ is constructed. The lemma 1.4 is crucial to prove theorem 1.5. Starting from the perturbative series of the resolvents of the hamiltonians, it allows to state that the norm difference between the ground states of $H_{\mathbf{P},\epsilon \frac{j+2}{2}}$ and of $H_{\mathbf{P},\epsilon \frac{j+1}{2}}$ is of order 1.

The initial hypotheses are:

- at the first step, the infrared cut-off is $\epsilon < \kappa$, $0 < \epsilon < \left(\frac{1}{4}\right)^{16}$;
- the mass m satisfies $m > 25 \cdot 4^{20}$;
- the coupling constant $g > 0$ and the ultraviolet cut-off κ satisfy the relation $2\pi g^2 \kappa \leq \frac{1}{4}$;
- the momentum $\mathbf{P}$ are restricted to the set $\Sigma \equiv \{\mathbf{P} : |\mathbf{P}| \leq \sqrt{m}\}$.

We synthesize the content of the lemmas:

- lemma 1.1 is a simple application of Kato's theorem to the hamiltonian with infrared cut-off ϵ in order to fix a coupling constant $\overline{g}_\epsilon$ such that a unique ground state $\psi_\mathbf{P}^\epsilon$ exists of energy $E_\mathbf{P}^\epsilon$ where the gap is bigger than $\frac{\epsilon}{2}$, for $\mathbf{P}$ in the set $\Sigma \equiv \{\mathbf{P} : |\mathbf{P}| \leq \sqrt{m}\}$;
- in the lemma 1.2 I study the operator $H_{\mathbf{P},\epsilon}$ restricted to the subspace $F_{\epsilon\sqrt{\epsilon}}^+$. Under the initial assumptions, $\psi_\mathbf{P}^\epsilon \otimes \psi_0$ is the unique ground state of $H_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+}$ of energy $E_\mathbf{P}^\epsilon$ and its gap is bigger than $\frac{3}{5}\epsilon\sqrt{\epsilon}$;
- lemma 1.3 proves that the ground energy is increasing in the infrared cut-off: $E_\mathbf{P}^\epsilon \geq E_\mathbf{P}^{\epsilon\sqrt{\epsilon}}$.

Lemma 1.1

Given ϵ and for $\mathbf{P} \in \Sigma \equiv \{\mathbf{P} : |\mathbf{P}| \leq \sqrt{m}\}$, there is a value for the coupling constant $\overline{g}_\epsilon \equiv \overline{g}_\epsilon(m, \kappa)$ such that the ground state $\psi_\mathbf{P}^\epsilon$ of $H_{\mathbf{P},\epsilon} |_{F_\epsilon^+}$ exists unique and the corresponding eigenvalue $E_\mathbf{P}^\epsilon$ is isolated. Its gap is bigger than $\frac{\epsilon}{2}$.

Proof.

If $\mathbf{P} \in \Sigma \equiv \{\mathbf{P} : |\mathbf{P}| \leq \sqrt{m}\}$ and $\varphi \in D^b \cap F_\epsilon^+$ ($D^b = \bigvee_{n \in \mathbb{N}} \psi^n$ the dense set in F^+ which is generated by the finite linear combinations of vectors of a finite number of mesons ($b, b^\dagger$) and of (symmetric) wave function $\psi^n(\mathbf{k}_1, \dots, \mathbf{k}_n) \in S(R^{3n})$, $n \in \mathbb{N}$) we have:

$$\left(\varphi, \left(H_\mathbf{P}^0 - \frac{\mathbf{P}^2}{2m}\right) \varphi\right) = \left(\varphi, \left(\frac{(\mathbf{P}^{mes})^2}{2m} - \frac{\mathbf{P} \cdot \mathbf{P}^{mes}}{m} + H^{mes}\right) \varphi\right) \geq$$

$$\geq \left(\varphi, \left(-\frac{\mathbf{P} \cdot \mathbf{P}^{mes}}{m} + H^{mes} \right) \varphi \right) = \left(\varphi, \int_{\epsilon}^{+\infty} \left(|\mathbf{k}| - \mathbf{k} \cdot \frac{\mathbf{P}}{m} \right) b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3 k \varphi \right);$$

therefore, if $\mathbf{P} \in \Sigma$, the vacuum vector ψ_0 is the ground eigenvector of $H_{\mathbf{P}}^0$, of energy $\frac{\mathbf{P}^2}{2m}$.

Let us consider the perturbation

$$H_{\mathbf{P},\epsilon}^I \equiv g \int_{\epsilon}^{\kappa} \frac{1}{\sqrt{2} |\mathbf{k}|^{\frac{1}{2}}} \left(b(\mathbf{k}) + b^{\dagger}(\mathbf{k}) \right) d^3 k$$

and an integration circle γ in the complex plane of radius $\frac{3}{4}\epsilon$ and centered in $E_{\mathbf{P}}^0 = \frac{\mathbf{P}^2}{2m}$. $E_{\mathbf{P}}^0 = \frac{\mathbf{P}^2}{2m}$ is the eigenvalue of the ground state ψ_0 of $H_{\mathbf{P}}^0 |_{F_{\epsilon}^+}$ when $\mathbf{P} \in \Sigma$. By Kato's theorem, for g sufficiently small and uniform in $\mathbf{P} \in \Sigma$, $g \leq \bar{g}_{\epsilon}$, there is a unique ground state $\psi_{\mathbf{P}}^{\epsilon}$ of $H_{\mathbf{P},\epsilon} |_{F_{\epsilon}^+}$, of energy $E_{\mathbf{P}}^{\epsilon} < \frac{1}{4}\epsilon + \frac{\mathbf{P}^2}{2m}$; moreover no other point of the spectrum of $H_{\mathbf{P},\epsilon} |_{F_{\epsilon}^+ \ominus \psi_{\mathbf{P}}^{\epsilon}}$ is inside γ .

Therefore the $\inf spec \left(H_{\mathbf{P},\epsilon} |_{F_{\epsilon}^+} \right)$ is $E_{\mathbf{P}}^{\epsilon}$ and the related gap is bigger than $\frac{1}{2}\epsilon$.

Remark

The ultraviolet *cut-off* κ and the mass m , with the initial constraints, are fixed. The value of g will be constrained several times during the procedure; at each time I will call $\bar{g}$ the maximum value such that the constraint under examination is satisfied and the previous constraints too.

Lemma 1.2

If $\psi_{\mathbf{P}}^{\epsilon}$ is the ground state of $H_{\mathbf{P},\epsilon} |_{F_{\epsilon}^+}$ with gap bigger than $\frac{1}{2}\epsilon$, then $\psi_{\mathbf{P}}^{\epsilon}$ is the ground state of $H_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+}$ (with the same eigenvalue) and its gap is bigger than $\frac{3}{5}\epsilon\sqrt{\epsilon}$.

Proof.

The proof is in two steps:

a) at first, I analyze the hamiltonian $H_{\mathbf{P},\epsilon} |_{F_{\epsilon}^+}$ plus terms of the difference $H_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} - H_{\mathbf{P},\epsilon} |_{F_{\epsilon}^+}$ in which the meson field modes of frequency between $\epsilon\sqrt{\epsilon}$ and ϵ do not interact with frequencies bigger than ϵ ;

b) then I will consider the interaction with frequencies bigger than ϵ too.

$\longleftrightarrow$

a)

We decompose $F_{\epsilon\sqrt{\epsilon}}^+$ as $F_{\epsilon}^+ \otimes F_{\epsilon\sqrt{\epsilon}}^{\epsilon}$, where $F_{\epsilon\sqrt{\epsilon}}^{\epsilon}$ is the tensorial sub-product defined as follows

$$F_{\epsilon\sqrt{\epsilon}}^{\epsilon} \equiv \left\{ \overline{\bigvee \psi^n(\mathbf{k}_1, \dots, \mathbf{k}_n)} : n \in \mathbb{N}, \psi^n(\mathbf{k}_1, \dots, \mathbf{k}_n) \in L^2(R^{3n}) \text{ simm.}, \epsilon \geq |\mathbf{k}_i| \geq \epsilon\sqrt{\epsilon}, 1 \leq i \leq n \right\}$$

I introduce the intermediate hamiltonian $\widehat{H}_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+}$:

$$\begin{aligned} \widehat{H}_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} &\equiv \frac{1}{2m} (\mathbf{P}^{mes} |_{\epsilon}^{+\infty})^2 \otimes 1 - \frac{\mathbf{P} \cdot \mathbf{P}^{mes} |_{\epsilon}^{+\infty}}{m} \otimes 1 + \frac{\mathbf{P}^2}{2m} \otimes 1 + \int_{\epsilon}^{+\infty} |\mathbf{k}| b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3k \otimes 1 + \\ &+ g \int_{\epsilon}^{\epsilon} (b(\mathbf{k}) + b^{\dagger}(\mathbf{k})) \frac{d^3\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{1}{2}}} \otimes 1 + 1 \otimes \left\{ \frac{1}{2m} (\mathbf{P}^{mes} |_{\epsilon\sqrt{\epsilon}}^{\epsilon})^2 + \frac{3}{4} \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3k \right\} \end{aligned}$$

that I denote as $H_1 \otimes 1 + 1 \otimes H_2$.

I observe that $\forall \varphi_1 \in F_{\epsilon}^+$ and $\forall \varphi_2 \in F_{\epsilon\sqrt{\epsilon}}^{\epsilon}$, φ_1 and φ_2 normalized vectors, we have

$$\left(\varphi_1 \otimes \varphi_2, \widehat{H}_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} \varphi_1 \otimes \varphi_2 \right) = (\varphi_1, H_1 \varphi_1) + (\varphi_2, H_2 \varphi_2) \geq (\varphi_1, H_1 \varphi_1) \geq E_{\mathbf{P}}^{\epsilon} \quad (1)$$

since

- $H_2 = \frac{1}{2m} (\mathbf{P}^{mes})^2 |_{\epsilon\sqrt{\epsilon}}^{\epsilon} + \frac{3}{4} \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3k$ is a positive operator
- $(\varphi_1, H_1 \varphi_1) = (\varphi_1, H_{\mathbf{P},\epsilon} |_{F_{\epsilon}^+} \varphi_1)$ if $\varphi_1 \in F_{\epsilon}^+$

Starting from the joined spectral decomposition of $H_1 \otimes 1$ and $1 \otimes H_2$ in $F_{\epsilon}^+ \otimes F_{\epsilon\sqrt{\epsilon}}^{\epsilon}$ we conclude that:

1) $E_{\mathbf{P}}^{\epsilon}$ is the ground energy of $\widehat{H}_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+}$, otherwise the condition (1) is not valid;

2) the gap of $E_{\mathbf{P}}^\epsilon$ (corresponding to the eigenvector $\psi_{\mathbf{P}}^\epsilon \otimes \psi_0$, ψ_0 is the Fock vacuum) is bigger than $\min \left\{ \frac{3}{4}\epsilon\sqrt{\epsilon}, \frac{\epsilon}{2} \right\}$; by the hypothesis $\epsilon < \left(\frac{1}{4} \right)^{16}$, then the gap is bigger than $\frac{3}{4}\epsilon\sqrt{\epsilon}$.

b)

$$H_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} = \widehat{H}_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} + \frac{\mathbf{P}^{mes}|_{\epsilon\sqrt{\epsilon}}^\epsilon}{m} (\mathbf{P}^{mes} |_{\epsilon}^{+\infty} - \mathbf{P}) + \frac{1}{4} \int_{\epsilon\sqrt{\epsilon}}^\epsilon |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k$$

I define:

$$\begin{aligned} \Delta \widehat{H}_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} &\equiv \frac{\mathbf{P}^{mes}|_{\epsilon\sqrt{\epsilon}}^\epsilon}{m} (\mathbf{P}^{mes} |_{\epsilon}^{+\infty} - \mathbf{P}) + \frac{1}{4} \int_{\epsilon\sqrt{\epsilon}}^\epsilon |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k = \\ &= \frac{\mathbf{P}^{mes}|_{\epsilon\sqrt{\epsilon}}^\epsilon}{m} (\mathbf{P}^{mes} |_{\epsilon}^{+\infty} - \mathbf{P}) + \frac{1}{5} \int_{\epsilon\sqrt{\epsilon}}^\epsilon |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k + \frac{1}{20} \int_{\epsilon\sqrt{\epsilon}}^\epsilon |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k \end{aligned}$$

Let us observe that from the decomposition of $F_{\epsilon\sqrt{\epsilon}}^+ = F_\epsilon^+ \otimes F_{\epsilon\sqrt{\epsilon}}^\epsilon$ in the two orthogonal subspaces $F_1 \oplus F_2$, where $F_1 \equiv \{\varphi \otimes \psi_0 : \varphi \in F_\epsilon^+\}$ and $F_2 \equiv \{\varphi \otimes \psi : \varphi \in F_\epsilon^+, \psi \in F_{\epsilon\sqrt{\epsilon}}^\epsilon, \psi \perp \psi_0\}$, we have:

$$\begin{aligned} H_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} &: F_1 \rightarrow F_1 \\ H_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} &: F_2 \rightarrow F_2. \end{aligned}$$

Spectrum of $H_{\mathbf{P},\epsilon} |_{F_1}$.

Since $H_{\mathbf{P},\epsilon} |_{F_1} \equiv \widehat{H}_{\mathbf{P},\epsilon} |_{F_1}$ the final results **1)** and **2)** of **a)** are valid.

Spectrum of $H_{\mathbf{P},\epsilon} |_{F_2}$.

In order to verify that $\inf spec(H_{\mathbf{P},\epsilon} |_{F_2}) \geq E_{\mathbf{P}}^\epsilon + \frac{3}{5}\epsilon\sqrt{\epsilon}$, I will prove the inequality

$$\inf spec \left(H_{\mathbf{P},\epsilon} |_{F_2} - \frac{1}{20} \int_{\epsilon\sqrt{\epsilon}}^\epsilon |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k |_{F_2} \right) \geq E_{\mathbf{P}}^\epsilon + \frac{3}{5}\epsilon\sqrt{\epsilon}$$

which implies the previous one and that will be useful in the lemma 1.4.

Given the joined spectral decomposition of the operators:

$$\begin{aligned} & - H_2 = \frac{1}{2m} (\mathbf{P}^{mes})^2 \big|_{\epsilon\sqrt{\epsilon}}^\epsilon + \frac{3}{4} H^{mes} \big|_{\epsilon\sqrt{\epsilon}}^\epsilon, \\ & - P^{mes^l} \big|_{\epsilon\sqrt{\epsilon}}^\epsilon \quad l = 1, 2, 3 \\ & - \frac{1}{5} H^{mes} \big|_{\epsilon\sqrt{\epsilon}}^\epsilon \end{aligned}$$

it is easy to verify that if two vectors φ_j and $\varphi_{j'}$ in $F_{\epsilon\sqrt{\epsilon}}^\epsilon$ have disjoint spectral supports, we have

$$(\varphi \otimes \varphi_{j'}, H_{\mathbf{P},\epsilon} \big|_{F_2} \varphi \otimes \varphi_j) = (\varphi \otimes \varphi_{j'}, (\widehat{H}_{\mathbf{P},\epsilon} \big|_{F_2} + \Delta \widehat{H}_{\mathbf{P},\epsilon} \big|_{F_2}) \varphi \otimes \varphi_j) = 0$$

Therefore, it is possible to restrict the analysis to the mean value of $H_{\mathbf{P},\epsilon} \big|_{F_2} - \frac{1}{20} \int_{\epsilon\sqrt{\epsilon}}^\epsilon |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k \big|_{F_2}$ applied to normalized vectors like $\varphi \otimes \varphi_j$, where $\varphi \in F_\epsilon^+$ is in the domain of $\widehat{H}_{\mathbf{P},\epsilon}$, $\varphi_j \in F_{\epsilon\sqrt{\epsilon}}^\epsilon$ is in the domain of $H^{mes} \big|_{\epsilon\sqrt{\epsilon}}^\epsilon$:

$$\begin{aligned} & (\varphi \otimes \varphi_j, H_{\mathbf{P},\epsilon} \big|_{F_2} - \frac{1}{20} \int_{\epsilon\sqrt{\epsilon}}^\epsilon |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k \big|_{F_2} \varphi \otimes \varphi_j) = \\ & = \left\{ (\varphi \otimes \varphi_j, \widehat{H}_{\mathbf{P},\epsilon} \big|_{F_2} \varphi \otimes \varphi_j) + 2 \left(\varphi \otimes \varphi_j, \frac{\mathbf{P}^{mes} \big|_{\epsilon\sqrt{\epsilon}}^\epsilon}{2m} (\mathbf{P}^{mes} \big|_\epsilon^{+\infty} - \mathbf{P}) \varphi \otimes \varphi_j \right) + \right. \\ & \left. + (\varphi \otimes \varphi_j, \frac{1}{5} H^{mes} \big|_{\epsilon\sqrt{\epsilon}}^\epsilon \varphi \otimes \varphi_j) \right\} \end{aligned}$$

I study the quantity written above as a function of two independent variables:

- $c_j \equiv (\varphi_j, \int_{\epsilon\sqrt{\epsilon}}^\epsilon |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k \varphi_j)$
- $E_{\mathbf{P},\varphi \otimes \varphi_j} \equiv (\varphi \otimes \varphi_j, \widehat{H}_{\mathbf{P},\epsilon} \big|_{F_{\epsilon\sqrt{\epsilon}}^+} \varphi \otimes \varphi_j)$

with the constraints $c_j > 0$ and $E_{\mathbf{P},\varphi \otimes \varphi_j} \geq E_{\mathbf{P}}^\epsilon + \frac{3}{4} \epsilon \sqrt{\epsilon}$.

Let us note that

$$\begin{cases} \widehat{H}_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} - \frac{1}{2m} (\mathbf{P}^{mes} |_{\epsilon}^{+\infty} - \mathbf{P})^2 - \frac{1}{2m} (\mathbf{P}^{mes} |_{\epsilon\sqrt{\epsilon}}^{\epsilon})^2 + 2\pi g^2 \kappa \geq 0 \\ [\mathbf{P}^{mes} |_{\epsilon\sqrt{\epsilon}}^{\epsilon}, \mathbf{P}^{mes} |_{\epsilon}^{+\infty}] = 0 \end{cases}$$

from which it follows that

$$(\varphi \otimes \varphi_j, \widehat{H}_{\mathbf{P},\epsilon} |_{F_2} \varphi \otimes \varphi_j) + \left(\varphi \otimes \varphi_j, \frac{\mathbf{P}^{mes} |_{\epsilon\sqrt{\epsilon}}^{\epsilon}}{m} (\mathbf{P}^{mes} |_{\epsilon}^{+\infty} - \mathbf{P}) \varphi \otimes \varphi_j \right) + 2\pi g^2 \kappa \geq 0$$

Since

$$H_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} = \widehat{H}_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} + \frac{\mathbf{P}^{mes} |_{\epsilon\sqrt{\epsilon}}^{\epsilon}}{m} (\mathbf{P}^{mes} |_{\epsilon}^{+\infty} - \mathbf{P}) + \frac{1}{4} \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k$$

for $c_j \geq 5$

$$\begin{aligned} & (\varphi \otimes \varphi_j, \{ H_{\mathbf{P},\epsilon} |_{F_2} - \frac{1}{20} \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k |_{F_2} \} \varphi \otimes \varphi_j) = \\ & = \left(\varphi \otimes \varphi_j, \{ \widehat{H}_{\mathbf{P},\epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} + \frac{\mathbf{P}^{mes} |_{\epsilon\sqrt{\epsilon}}^{\epsilon}}{m} (\mathbf{P}^{mes} |_{\epsilon}^{+\infty} - \mathbf{P}) + \frac{1}{5} \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k \} \varphi \otimes \varphi_j \right) \geq \\ & \geq \frac{1}{5} c_j - 2\pi g^2 \kappa \geq 1 - 2\pi g^2 \kappa \end{aligned}$$

In order to study the case $0 < c_j < 5$, since $\widehat{H}_{\mathbf{P},\epsilon} + 2\pi g^2 \kappa - \frac{(\mathbf{P}^{mes^l} |_{\epsilon}^{+\infty} - P^l)^2}{2m}$ is a positive operator, I observe that

$$\begin{aligned} & \left| \left(\varphi \otimes \varphi_j, \frac{\mathbf{P}^{mes} |_{\epsilon\sqrt{\epsilon}}^{\epsilon}}{m} (\mathbf{P}^{mes} |_{\epsilon}^{+\infty} - \mathbf{P}) \varphi \otimes \varphi_j \right) \right| = \left| (\varphi, (\mathbf{P}^{mes} |_{\epsilon}^{+\infty} - \mathbf{P}) \varphi) \cdot \left(\varphi_j, \frac{\mathbf{P}^{mes} |_{\epsilon\sqrt{\epsilon}}^{\epsilon}}{m} \varphi_j \right) \right| \leq \\ & \leq \sum_{l=1}^3 \frac{1}{m} \left| (\varphi, (P^{mes^l} |_{\epsilon}^{+\infty} - P^l) \varphi) \right| \left| (\varphi_j, \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k \varphi_j) \right| \leq \\ & \leq (E_{\mathbf{P},\varphi \otimes \varphi_j} + 2\pi g^2 \kappa)^{\frac{1}{2}} \cdot 3\sqrt{\frac{2}{m}} (\varphi_j, \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k \varphi_j) \leq \\ & \leq (E_{\mathbf{P},\varphi \otimes \varphi_j} + 2\pi g^2 \kappa)^{\frac{1}{2}} \cdot 3c_j \sqrt{\frac{2}{m}} \end{aligned}$$

from which

$$(\varphi \otimes \varphi_j, (H_{\mathbf{P},\epsilon} |_{F_2} - \frac{1}{20} \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k |_{F_2}) \varphi \otimes \varphi_j) =$$

$$\begin{aligned}
&= \left\{ \left(\varphi \otimes \varphi_j, \widehat{H}_{\mathbf{P},\epsilon} |_{F_2} \varphi \otimes \varphi_j \right) + \left(\varphi \otimes \varphi_j, \frac{\mathbf{P}^{mes}|_{\epsilon\sqrt{\epsilon}}}{m} (\mathbf{P}^{mes}|_{\epsilon}^{+\infty} - \mathbf{P}) \varphi \otimes \varphi_j \right) + \right. \\
&\quad \left. + \left(\varphi \otimes \varphi_j, \frac{1}{5} H^{mes}|_{\epsilon\sqrt{\epsilon}} \varphi \otimes \varphi_j \right) \right\} \geq E_{\mathbf{P},\varphi \otimes \varphi_j} - 3c_j \sqrt{\frac{2}{m}} \left(E_{\mathbf{P},\varphi \otimes \varphi_j} + 2\pi g^2 \kappa \right)^{\frac{1}{2}} + \frac{1}{5} c_j
\end{aligned}$$

I define the function:

$$\begin{aligned}
f(E_{\mathbf{P},\varphi \otimes \varphi_j}, c_j) &\equiv E_{\mathbf{P},\varphi \otimes \varphi_j} - 3c_j \sqrt{\frac{2}{m}} \left(E_{\mathbf{P},\varphi \otimes \varphi_j} + 2\pi g^2 \kappa \right)^{\frac{1}{2}} + \frac{1}{5} c_j - \left\{ \frac{4}{5} (E_{\mathbf{P},\varphi \otimes \varphi_j} - E_{\mathbf{P}}^{\epsilon}) + E_{\mathbf{P}}^{\epsilon} \right\} = \\
&= \frac{1}{5} E_{\mathbf{P},\varphi \otimes \varphi_j} - 3c_j \sqrt{\frac{2}{m}} \left(E_{\mathbf{P},\varphi \otimes \varphi_j} + 2\pi g^2 \kappa \right)^{\frac{1}{2}} + \frac{1}{5} c_j - \frac{1}{5} E_{\mathbf{P}}^{\epsilon}
\end{aligned}$$

Analysis of $f(E_{\mathbf{P},\varphi \otimes \varphi_j}, c_j)$.

First of all we observe that the positivity of $f(E_{\mathbf{P},\varphi \otimes \varphi_j}, c_j)$ implies:

$$\begin{aligned}
&\left(\varphi \otimes \varphi_j, \left(H_{\mathbf{P},\epsilon} |_{F_2} - \frac{1}{20} \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3 k \right) |_{F_2} \varphi \otimes \varphi_j \right) \geq \\
&\geq \frac{4}{5} (E_{\mathbf{P},\varphi \otimes \varphi_j} - E_{\mathbf{P}}^{\epsilon}) + E_{\mathbf{P}}^{\epsilon} \geq \frac{3}{5} \epsilon \sqrt{\epsilon} + E_{\mathbf{P}}^{\epsilon}
\end{aligned}$$

Discussion

For the initial assumptions $m > \left(\frac{75}{\sqrt{2}}\right)^2$ e $2\pi g^2 \kappa < \frac{1}{4}$; the values of the variables of f to be considered are: $0 < c_j < 5$ and $E_{\mathbf{P},\varphi \otimes \varphi_j} \geq E_{\mathbf{P}}^{\epsilon} + \frac{3}{4} \epsilon \sqrt{\epsilon}$.

I will consider separately two cases:

i) if $E_{\mathbf{P},\varphi \otimes \varphi_j} \leq 1 - 2\pi g^2 \kappa$,

then $\frac{1}{5} c_j - 3c_j \sqrt{\frac{2}{m}} \left(E_{\mathbf{P},\varphi \otimes \varphi_j} + 2\pi g^2 \kappa \right)^{\frac{1}{2}} \geq 0$

from which $f(E_{\mathbf{P},\varphi \otimes \varphi_j}, c_j) \geq \frac{1}{5} (E_{\mathbf{P},\varphi \otimes \varphi_j} - E_{\mathbf{P}}^{\epsilon}) > 0$;

ii) if $E_{\mathbf{P}, \varphi \otimes \varphi_j} > 1 - 2\pi g^2 \kappa$,

let us observe that $\frac{\partial}{\partial E_{\mathbf{P}, \varphi \otimes \varphi_j}} f(E_{\mathbf{P}, \varphi \otimes \varphi_j}, c_j) \equiv \frac{1}{5} - 3c_j \frac{1}{\sqrt{2m}} (E_{\mathbf{P}, \varphi \otimes \varphi_j} + 2\pi g^2 \kappa)^{-\frac{1}{2}}$ from which $\frac{\partial}{\partial E_{\mathbf{P}, \varphi \otimes \varphi_j}} f(E_{\mathbf{P}, \varphi \otimes \varphi_j}, c_j) \geq 0$ for $E_{\mathbf{P}, \varphi \otimes \varphi_j} \geq \left(15c_j \frac{1}{\sqrt{2m}}\right)^2 - 2\pi g^2 \kappa$, that is a minimum value at fixed c_j ; since $\left(15c_j \frac{1}{\sqrt{2m}}\right)^2 < 1$, by exploiting the result i), we arrive at $f(E_{\mathbf{P}, \varphi \otimes \varphi_j}, c_j) \geq f\left(\left(15c_j \frac{1}{\sqrt{2m}}\right)^2 - 2\pi g^2 \kappa, c_j\right) > 0$.

If $E_{\mathbf{P}}^\epsilon + \frac{3}{5}\epsilon\sqrt{\epsilon} \leq 1 - 2\pi g^2 \kappa$, the previous results tell us that

$$\inf_{\varphi \otimes \varphi_j} \left(\varphi \otimes \varphi_j, \left\{ H_{\mathbf{P}, \epsilon} \big|_{F_2} - \frac{1}{20} \int_{\epsilon\sqrt{\epsilon}}^\epsilon |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k \big|_{F_2} \right\} \varphi \otimes \varphi_j \right) \geq E_{\mathbf{P}}^\epsilon + \frac{3}{5}\epsilon\sqrt{\epsilon}$$

Conclusion

The results in **a)** and **b)** imply that, **if $E_{\mathbf{P}}^\epsilon + \frac{3}{5}\epsilon\sqrt{\epsilon} \leq 1 - 2\pi g^2 \kappa$ then the minimum value of the spectrum of $H_{\mathbf{P}, \epsilon} \big|_{F_{\epsilon\sqrt{\epsilon}}^+}$ is $E_{\mathbf{P}}^\epsilon$ and the gap is bigger or equal to $\frac{3}{5}\epsilon\sqrt{\epsilon}$.** For the lemma 1.1 the value of the coupling constant is such that $E_{\mathbf{P}}^\epsilon < \frac{1}{4}\epsilon + \frac{\mathbf{P}^2}{2m}$. Since $\epsilon < \left(\frac{1}{4}\right)^{16}$, $2\pi g^2 \kappa < \frac{1}{4}$ and $\mathbf{P} \in \Sigma$, the condition $E_{\mathbf{P}}^\epsilon + \frac{3}{5}\epsilon\sqrt{\epsilon} \leq 1 - 2\pi g^2 \kappa$ is satisfied.

Lemma 1.3

The following relation between $E_{\mathbf{P}}^\epsilon$, $E_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}}$ (ground energy of $H_{\mathbf{P}, \epsilon\sqrt{\epsilon}} \big|_{F_{\epsilon\sqrt{\epsilon}}^+}$) holds:

$$E_{\mathbf{P}}^\epsilon \geq E_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}} \geq E_{\mathbf{P}}^\epsilon - 40\pi g^2 \epsilon$$

Proof.

$$H_{\mathbf{P}, \epsilon\sqrt{\epsilon}} \big|_{F_{\epsilon\sqrt{\epsilon}}^+} = \frac{1}{2m} (\mathbf{P}^{mes} \big|_{\epsilon}^{+\infty})^2 \otimes 1 + \int_{\epsilon}^{+\infty} |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k \otimes 1 - \frac{\mathbf{P} \cdot \mathbf{P}^{mes} \big|_{\epsilon}^{+\infty}}{m} \otimes 1 + \frac{\mathbf{P}^2}{2m} \otimes 1 +$$

$$\begin{aligned}
& +g \int_{\epsilon}^{\kappa} \left(b(\mathbf{k}) + b^{\dagger}(\mathbf{k}) \right) \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{1}{2}}} \otimes 1 + 1 \otimes \frac{1}{2m} \left(\mathbf{P}^{mes} |_{\epsilon\sqrt{\epsilon}}^{\epsilon} \right)^2 + \mathbf{P}^{mes} |_{\epsilon}^{+\infty} \otimes \frac{\mathbf{P}^{mes} |_{\epsilon\sqrt{\epsilon}}^{\epsilon}}{m} + \\
& -1 \otimes \frac{\mathbf{P} \cdot \mathbf{P}^{mes} |_{\epsilon\sqrt{\epsilon}}^{\epsilon}}{m} + 1 \otimes \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3 k + 1 \otimes g \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} \left(b(\mathbf{k}) + b^{\dagger}(\mathbf{k}) \right) \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{1}{2}}}
\end{aligned}$$

The mean value of $H_{\mathbf{P}, \epsilon\sqrt{\epsilon}} |_{F_{\epsilon\sqrt{\epsilon}}^+}$ on $\psi_{\mathbf{P}}^{\epsilon} \otimes \psi_0$ (normalized) is $E_{\mathbf{P}}^{\epsilon}$.

By definition, $E_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}}$ is the inf of the mean value of $H_{\mathbf{P}, \epsilon\sqrt{\epsilon}} |_{F_{\epsilon\sqrt{\epsilon}}^+}$ on the normalized vectors in $F_{\epsilon\sqrt{\epsilon}}^+$ belonging to the operator domain. Therefore $E_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}} \leq E_{\mathbf{P}}^{\epsilon}$ and in general for $\sigma_1 > \sigma_2$ $E_{\mathbf{P}}^{\sigma_2} \leq E_{\mathbf{P}}^{\sigma_1}$.

Moreover, as proved in the previous lemma

$$\inf spec \left(H_{\mathbf{P}, \epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} - \frac{1}{20} \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3 k \right) > E_{\mathbf{P}}^{\epsilon}$$

while completing the square

$$\frac{1}{20} \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3 k + g \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} \left(b(\mathbf{k}) + b^{\dagger}(\mathbf{k}) \right) \frac{d^3 k}{\sqrt{2}|\mathbf{k}|^{\frac{1}{2}}} + 40\pi g^2 \epsilon \geq 0$$

It follows that

$$\begin{aligned}
E_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}} &= \inf spec \left(H_{\mathbf{P}, \epsilon\sqrt{\epsilon}} |_{F_{\epsilon\sqrt{\epsilon}}^+} \right) = \inf spec \left(H_{\mathbf{P}, \epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} + g \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} \left(b(\mathbf{k}) + b^{\dagger}(\mathbf{k}) \right) \frac{d^3 k}{\sqrt{2}|\mathbf{k}|^{\frac{1}{2}}} \right) \geq \\
&\geq \inf spec \left(H_{\mathbf{P}, \epsilon} |_{F_{\epsilon\sqrt{\epsilon}}^+} - \frac{1}{20} \int_{\epsilon\sqrt{\epsilon}}^{\epsilon} |\mathbf{k}| b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3 k - 40\pi g^2 \epsilon \right) \geq E_{\mathbf{P}}^{\epsilon} - 40\pi g^2 \epsilon
\end{aligned}$$

Coupling constant constraints.

In the following lemma 1.4 a value $\bar{g}$ will be fixed. It allows to estimate the difference between the ground states at different and arbitrarily small cut-off σ_j . The validity of lemma 1.2 at each step for fixed $\bar{g}$ is necessary for the consistency of the iteration. The relation in lemma 1.3 assures the validity of lemma 1.2 since:

$$E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} + \frac{3}{5} \epsilon^{\frac{j+2}{2}} \leq E_{\mathbf{P}}^{\epsilon} + \frac{3}{5} \epsilon \sqrt{\epsilon} < 1 - 2\pi \bar{g}^2 \kappa \quad \forall j$$

Definitions

$$H_{\mathbf{P},\epsilon\sqrt{\epsilon}}|_{F_{\epsilon\sqrt{\epsilon}}^+} = H_{\mathbf{P},\epsilon}|_{F_{\epsilon\sqrt{\epsilon}}^+} + (\Delta H_{\mathbf{P}})_{\epsilon\sqrt{\epsilon}}^\epsilon$$

$$(\Delta H_{\mathbf{P}})_{\epsilon\sqrt{\epsilon}}^\epsilon \equiv g \int_{\epsilon\sqrt{\epsilon}}^\epsilon \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) \frac{d^3\mathbf{k}}{\sqrt{2}|\mathbf{k}|}$$

Lemma 1.4

For a properly small $\overline{g}$, $(\Delta H_{\mathbf{P}})_{\epsilon\sqrt{\epsilon}}^\epsilon$ is small of order 1 with respect to $H_{\mathbf{P},\epsilon}|_{F_{\epsilon\sqrt{\epsilon}}^+}$, in the following sense:

given E such that $\left| E - \inf H_{\mathbf{P},\epsilon}|_{F_{\epsilon\sqrt{\epsilon}}^+} \right| = |E - E_{\mathbf{P}}^\epsilon| = \frac{20\epsilon\sqrt{\epsilon}}{11}$

$$\frac{1}{H_{\mathbf{P},\epsilon} + (\Delta H_{\mathbf{P}})_{\epsilon\sqrt{\epsilon}}^\epsilon - E} \Big|_{F_{\epsilon\sqrt{\epsilon}}^+} = \frac{1}{H_{\mathbf{P},\epsilon} - E} \Big|_{F_{\epsilon\sqrt{\epsilon}}^+} + \frac{1}{H_{\mathbf{P},\epsilon} - E} \sum_{n=1}^{+\infty} \left(- (\Delta H_{\mathbf{P}})_{\epsilon\sqrt{\epsilon}}^\epsilon \frac{1}{H_{\mathbf{P},\epsilon}|_{F_{\epsilon\sqrt{\epsilon}}^+} - E} \right)^n \Big|_{F_{\epsilon\sqrt{\epsilon}}^+}$$

and

$$\left\| \frac{1}{H_{\mathbf{P},\epsilon}|_{F_{\epsilon\sqrt{\epsilon}}^+} - E} \left(- (\Delta H_{\mathbf{P}})_{\epsilon\sqrt{\epsilon}}^\epsilon \frac{1}{H_{\mathbf{P},\epsilon}|_{F_{\epsilon\sqrt{\epsilon}}^+} - E} \right)^n \right\|_{F_{\epsilon\sqrt{\epsilon}}^+} \leq \frac{20(C(\overline{g}, \overline{m}))^n}{\epsilon\sqrt{\epsilon}}$$

where $0 < C(\overline{g}, m) < \frac{1}{12}$.

Theorem 1.5

$H_{\mathbf{P},\epsilon\sqrt{\epsilon}}|_{F_{\epsilon\sqrt{\epsilon}}^+}$ has a unique ground eigenvector $\psi_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}}$ of energy $E_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}}$ and gap $\frac{\epsilon\sqrt{\epsilon}}{2}$; the un-normalized vector $\psi_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}}$ corresponds to

$$\psi_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}} = -\frac{1}{2\pi i} \oint \frac{1}{H_{\mathbf{P},\epsilon\sqrt{\epsilon}} - E} dE \psi_{\mathbf{P}}^\epsilon \quad (2)$$


where $E \in \mathcal{C}$ and $|E - E_{\mathbf{P}}^\epsilon| = \frac{11}{20}\epsilon\sqrt{\epsilon}$.

Proof.

Continuity argument.

I distinguish the coupling constant g in $H_{\mathbf{P},\epsilon}|_{F_{\epsilon\sqrt{\epsilon}}^+}$ from that one in $(\Delta H_{\mathbf{P}})_{\epsilon\sqrt{\epsilon}}^\epsilon$, and I call the last one $g^\sharp$. Kato's theorem ensures that (2) is verified for sufficiently small $g^\sharp$, since the gap of $H_{\mathbf{P},\epsilon}|_{F_{\epsilon\sqrt{\epsilon}}^+}$ is bigger or equal to $\frac{3}{5}\epsilon\sqrt{\epsilon}$ and $(\Delta H_{\mathbf{P}}(g^\sharp))_{\epsilon\sqrt{\epsilon}}^\epsilon$ is a small Kato perturbation with respect to $H_{\mathbf{P},\epsilon}|_{F_{\epsilon\sqrt{\epsilon}}^+}$.

Now look at the figure



if $g^\sharp$ increases, the equation (2) is valid till the eigenvalue $E_{\mathbf{P},g^\sharp}^{\epsilon\sqrt{\epsilon}}$ remains inside the circle of integration and the remaining spectrum of $H_{\mathbf{P},\epsilon\sqrt{\epsilon}}|_{F_{\epsilon\sqrt{\epsilon}}^+}$ remains outside of the circle of integration; a limit value $\overline{g^\sharp}$ exists for which the expression $-\frac{1}{2\pi i} \oint \left\| \frac{1}{H_{\mathbf{P},\epsilon\sqrt{\epsilon}}|_{F_{\epsilon\sqrt{\epsilon}}^+} - E} \right\| dE$ diverges because the spectrum intersects the circle of integration.

According to the estimates in lemma 1.4 we can conclude that:

- the integral $-\frac{1}{2\pi i} \oint \left\| \frac{1}{H_{\mathbf{P},\epsilon\sqrt{\epsilon}}|_{F_{\epsilon\sqrt{\epsilon}}^+} - E} \right\| dE$ exists for $0 \leq g^\sharp \leq g = \overline{g}$ and then $\overline{g} < \overline{g^\sharp}$
- the ground state of $H_{\mathbf{P},\epsilon\sqrt{\epsilon}}|_{F_{\epsilon\sqrt{\epsilon}}^+}$ is unique and it is not zero since

$$\psi_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}} = \psi_{\mathbf{P}}^\epsilon - \frac{1}{2\pi i} \sum_{n=1}^{+\infty} \oint \frac{1}{H_{\mathbf{P},\epsilon\sqrt{\epsilon}}|_{F_{\epsilon\sqrt{\epsilon}}^+} - E} \left(-(\Delta H_{\mathbf{P}})_{\epsilon\sqrt{\epsilon}}^\epsilon \frac{1}{H_{\mathbf{P},\epsilon\sqrt{\epsilon}}|_{F_{\epsilon\sqrt{\epsilon}}^+} - E} \right)^n dE \psi_{\mathbf{P}}^\epsilon \quad (3)$$

where the norm of the remainder $-\frac{1}{2\pi i} \sum_{n=1}^{+\infty} \oint \frac{1}{H_{\mathbf{P},\epsilon}|_{F_{\epsilon\sqrt{\epsilon}}^+} - E} \left(-(\Delta H_{\mathbf{P}})_{\epsilon\sqrt{\epsilon}} \frac{1}{H_{\mathbf{P},\epsilon}|_{F_{\epsilon\sqrt{\epsilon}}^+} - E} \right)^n dE \psi_{\mathbf{P}}^{\epsilon}$ is less than $\frac{11C(\bar{g},m)}{1-C(\bar{g},m)} \|\psi_{\mathbf{P}}^{\epsilon}\|$. Therefore $\psi_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}}$ is not zero since:

$$\|\psi_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}}\| \geq \|\psi_{\mathbf{P}}^{\epsilon}\| - \frac{11C(\bar{g},m)}{1-C(\bar{g},m)} \|\psi_{\mathbf{P}}^{\epsilon}\| \geq \frac{1-12C(\bar{g},m)}{1-C(\bar{g},m)} \|\psi_{\mathbf{P}}^{\epsilon}\| > 0$$

- since for lemma 1.3 $E_{\mathbf{P}}^{\epsilon\sqrt{\epsilon}} \leq E_{\mathbf{P}}^{\epsilon}$, the gap is bigger than $\frac{\epsilon\sqrt{\epsilon}}{2}$.

Corollary 1.6

Thanks to the results of theorem 1.5 about the existence of the ground state of $H_{\mathbf{P},\epsilon\sqrt{\epsilon}}|_{F_{\epsilon\sqrt{\epsilon}}^+}$ and about the *gap* (bigger than $\frac{\epsilon\sqrt{\epsilon}}{2}$), it is possible to iterate the procedure at fixed $\bar{g}$, by applying lemmas 1.2, 1.3 to $H_{\mathbf{P},\epsilon\sqrt{\epsilon}}|_{F_{\epsilon\sqrt{\epsilon}}^+}$ and $H_{\mathbf{P},\epsilon^2}|_{F_{\epsilon^2}^+}$ and by using properly adapted versions of lemma 1.4 and of theorem 1.5 to compare $H_{\mathbf{P},\epsilon\sqrt{\epsilon}}|_{F_{\epsilon\sqrt{\epsilon}}^+}$ and $H_{\mathbf{P},\epsilon^2}|_{F_{\epsilon^2}^+}$.

Therefore the iteration is consistent and it does not end since the vector obtained at the step $j+1$ has norm bigger than a fixed fraction of the norm of the vector at the j step. At each step the cut-off is reduced by a factor $\epsilon^{\frac{1}{2}}$, so that at the j step we obtain:

$$\psi_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} = \psi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} - \frac{1}{2\pi i} \sum_{n=1}^{+\infty} \oint \frac{1}{H_{\mathbf{P},\epsilon^{\frac{j+1}{2}}}|_{F_{\epsilon^{\frac{j+2}{2}}}^+} - E} \left(-(\Delta H_{\mathbf{P}})_{\epsilon^{\frac{j+1}{2}}} \frac{1}{H_{\mathbf{P},\epsilon^{\frac{j+1}{2}}}|_{F_{\epsilon^{\frac{j+2}{2}}}^+} - E} \right)^n dE \psi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}$$

Lemma 1.4

For fixed and properly small $\bar{g}$, $(\Delta H_{\mathbf{P}})_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j+2}{2}}}$ is small of order 1 with respect to $H_{\mathbf{P},\epsilon^{\frac{j+1}{2}}}|_{F_{\epsilon^{\frac{j+2}{2}}}^+}$, in the following sense:

given E such that $\left| E - \inf H_{\mathbf{P},\epsilon^{\frac{j+1}{2}}}|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \right| = \left| E - E_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} \right| = \frac{11\epsilon^{\frac{j+2}{2}}}{20}$

$$\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} + (\Delta H_{\mathbf{P}})^{\epsilon}_{\frac{j+2}{2}} - E} \Big|_{F_{\epsilon}^{+\frac{j+2}{2}}} = \frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \sum_{n=0}^{+\infty} \left(-(\Delta H_{\mathbf{P}})^{\epsilon}_{\frac{j+2}{2}} \frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^n \Big|_{F_{\epsilon}^{+\frac{j+2}{2}}}$$

and $\left\| \frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \left(-(\Delta H_{\mathbf{P}})^{\epsilon}_{\frac{j+2}{2}} \frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^n \Big|_{F_{\epsilon}^{+\frac{j+2}{2}}} \right\|_{F_{\epsilon}^{+\frac{j+2}{2}}} \leq \frac{20(C(\bar{g}, \bar{m}))^n}{\epsilon^{\frac{j+2}{2}}}$

where $0 < C(\bar{g}, m) < \frac{1}{12}$ is a constant independent of $\epsilon^{\frac{j+1}{2}}$.

Proof

Let us analyze the n^{th} term of the following sum

$$\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \sum_{n=1}^{+\infty} \left(-(\Delta H_{\mathbf{P}})^{\epsilon}_{\frac{j+2}{2}} \frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)$$

$$(-1)^n \frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} (\Delta H_{\mathbf{P}})^{\epsilon}_{\frac{j+2}{2}} \frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \dots (\Delta H_{\mathbf{P}})^{\epsilon}_{\frac{j+2}{2}} \frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} =$$

$$= (-1)^n \left(\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \dots \left(\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} (\Delta H_{\mathbf{P}})^{\epsilon}_{\frac{j+2}{2}} \left(\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \dots \left(\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}}$$

where $\left(\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}}$ is defined starting from the spectral representation of $H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}}$ by using the convention to take the branch of the square root with smaller argument in $(-\pi, \pi]$.

Study of the norm of $\left(\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} (\Delta H_{\mathbf{P}})^{\epsilon}_{\frac{j+2}{2}} \left(\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \Big|_{F_{\epsilon}^{+\frac{j+2}{2}}}$.

$$\left\| \left(\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} (\Delta H_{\mathbf{P}})^{\epsilon}_{\frac{j+2}{2}} \left(\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon}^{+\frac{j+2}{2}}} =$$

$$= \left\| \left(\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \left(g \int_{\epsilon^{\frac{j+2}{2}}}^{\frac{j+1}{2}} \frac{1}{\sqrt{2}|\mathbf{k}|^{\frac{1}{2}}} (b(\mathbf{k}) + b^{\dagger}(\mathbf{k})) d^3k \right) \left(\frac{1}{H_{\mathbf{P},\epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon}^{+\frac{j+2}{2}}} =$$

$$\leq 2g \left\| \left(\frac{1}{H_{\mathbf{P},\epsilon} \frac{j+1}{2} - E} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon \frac{j+2}{2}}^+} \cdot \left\| \left(\int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{\sqrt{2}|\mathbf{k}|^{\frac{1}{2}}} b(\mathbf{k}) d^3 k \right) \left(\frac{1}{H_{\mathbf{P},\epsilon} \frac{j+1}{2} - E} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon \frac{j+2}{2}}^+}$$

if the above quantities exist.

The following estimate holds:

$$\left\| \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} b(\mathbf{k}) \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|} \left(\frac{1}{H_{\mathbf{P},\epsilon} \frac{j+1}{2} - E} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon \frac{j+2}{2}}^+} \leq \sqrt{40\pi} \cdot \epsilon^{\frac{j+1}{4}} \quad (4)$$

Proof

Let us consider vectors $\varphi \in D^b \cap F_{\epsilon \frac{j+2}{2}}^+$; then

$$\begin{aligned} & \left\| \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} b(\mathbf{k}) \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|} \left(\frac{1}{H_{\mathbf{P},\epsilon} \frac{j+1}{2} - E} \right)^{\frac{1}{2}} \varphi \right\|_{F_{\epsilon \frac{j+2}{2}}^+}^2 = \\ & = \left(\left(\frac{1}{H_{\mathbf{P},\epsilon} \frac{j+1}{2} - E} \right)^{\frac{1}{2}} \varphi, \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} b^\dagger(\mathbf{k}) \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|} \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} b(\mathbf{k}) \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|} \left(\frac{1}{H_{\mathbf{P},\epsilon} \frac{j+1}{2} - E} \right)^{\frac{1}{2}} \varphi \right) \end{aligned}$$

In general

$$\begin{aligned} & \left(\varphi, \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{\sqrt{2}|\mathbf{k}|} \frac{1}{\sqrt{2}|\mathbf{q}|} b^\dagger(\mathbf{k}) b(\mathbf{q}) d^3 k d^3 q \varphi \right) = \\ & = \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{\sqrt{2}|\mathbf{k}|} \frac{1}{\sqrt{2}|\mathbf{q}|} \left(\varphi, b^\dagger(\mathbf{k}) b(\mathbf{q}) \varphi \right) d^3 k d^3 q \leq \\ & \leq \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{\sqrt{2}|\mathbf{k}|} \frac{1}{\sqrt{2}|\mathbf{q}|} \left(\varphi, b^\dagger(\mathbf{k}) b(\mathbf{k}) \varphi \right)^{\frac{1}{2}} \left(\varphi, b^\dagger(\mathbf{q}) b(\mathbf{q}) \varphi \right)^{\frac{1}{2}} d^3 k d^3 q \leq \\ & \leq \left(\varphi, H^{mes} \Big|_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \varphi \right) \cdot \left(\int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{2|\mathbf{k}|^2} d^3 k \right)^{\frac{1}{2}} \cdot \left(\int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{2|\mathbf{q}|^2} d^3 q \right)^{\frac{1}{2}} \leq \\ & \leq 2\pi\epsilon^{\frac{j+1}{2}} \left(\varphi, H^{mes} \Big|_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \varphi \right) \end{aligned} \quad (5)$$

and the inequality holds in the case of vectors in $D^b(H^{mes})$ too.

Therefore we arrive at

$$\begin{aligned} & \left(\left(\frac{1}{H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \varphi, H^{mes} \Big|_{\epsilon^{\frac{j+2}{2}}} \left(\frac{1}{H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \varphi \right) \leq \\ & \leq \|\varphi\| \cdot \left\| H^{mes} \Big|_{\epsilon^{\frac{j+2}{2}}} \left[\left(\frac{1}{H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \right]^{\dagger} \left(\frac{1}{H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \varphi \right\|_{F_{\epsilon}^{+ \frac{j+2}{2}}} \\ & \left(\left[H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}}, \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} |\mathbf{k}| b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3 k \right] = 0 \right) \end{aligned}$$

The operator norm of $H^{mes} \Big|_{\epsilon^{\frac{j+2}{2}}} \cdot \left[\left(\frac{1}{H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \right]^{\dagger} \cdot \left(\frac{1}{H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}}$ has to be studied separately on $\psi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}$ and on $F_{\epsilon}^{+ \frac{j+2}{2}} \ominus \psi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}$. The operator vanishes on $\psi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}$ (put $H^{mes} \Big|_{\epsilon^{\frac{j+2}{2}}}$ on the right). The discussion is restricted to the subspace $F_{\epsilon}^{+ \frac{j+2}{2}} \ominus \psi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}$. Moreover, as already seen in lemma 1.2. adapted to the hamiltonian $H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} \Big|_{F_{\epsilon}^{+ \frac{j+2}{2}}}$, we have

- $\inf spec \left(H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} \Big|_{F_{\epsilon}^{+ \frac{j+2}{2}}} - \frac{1}{20} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} |\mathbf{k}| b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3 k \Big|_{F_{\epsilon}^{+ \frac{j+2}{2}}} \right) \geq E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} + \frac{3}{5} \epsilon^{\frac{j+2}{2}}$
- $\inf spec \left(H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} \Big|_{F_{\epsilon}^{+ \frac{j+2}{2}}} - \frac{1}{20} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} |\mathbf{k}| b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3 k \Big|_{F_{\epsilon}^{+ \frac{j+2}{2}}} - Re E \right) \geq$
 $\geq \frac{3}{5} \epsilon^{\frac{j+2}{2}} - \frac{11}{20} \epsilon^{\frac{j+2}{2}} > 0$

Going to the **joined spectral representation** of $H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}}$ and $H^{mes} \Big|_{\epsilon^{\frac{j+2}{2}}}$, we obtain

$$\left\| H^{mes} \Big|_{\epsilon^{\frac{j+2}{2}}} \left[\left(\frac{1}{H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \right]^{\dagger} \left(\frac{1}{H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon}^{+ \frac{j+2}{2}}} \leq 20$$

(6)

Putting together the inequalities (5) and (6) we arrive at the inequality (4).

Conclusion

If g is less than a limit value $\bar{g}$, the thesis is proved since $\left\| \left(\frac{1}{H_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} - E} \right)^{\frac{1}{2}} \right\|_{F^+_{\epsilon^{\frac{j+2}{2}}}}$ is of order $\frac{1}{\epsilon^{\frac{j+2}{4}}}$.

2 Convergence of the ground states of transformed $H_{\mathbf{P}, \sigma_j}$.

I conjecture that the hamiltonians $H_{\mathbf{P}}$ have a ground state for the representations of $\{b(\mathbf{k}), b^\dagger(\mathbf{k})\}$ which are coherent in the infrared region ($\mathbf{k} = 0$). Then an argument is developed which explicitly identifies the eventual coherent factor in the case $\mathbf{P} = 0$ and implicitly in the case $\mathbf{P} \neq 0$. Such a heuristic information will be used in a rigorous proof which is based on the iterative procedure of construction of the ground state. The two cases $\mathbf{P} = 0$ and $\mathbf{P} \neq 0$ are treated separately.

Derivation of the coherent factor.

Let us assume that $\psi_{\mathbf{P}}$ is an eigenvector of $H_{\mathbf{P}}$ and that it is “coherent in the infrared region”, which means $b(\mathbf{k}) \psi_{\mathbf{P}} \approx f_{\mathbf{P}}(\mathbf{k}) \psi_{\mathbf{P}}$ for $\mathbf{k} \rightarrow 0$, where the meaning of the limit is given only “a posteriori”. Then the coherent function $f_{\mathbf{P}}(\mathbf{k})$ has to satisfies the following relation, in the neighborhood of $\mathbf{k} = 0$:

$$(\psi_{\mathbf{P}}, [H_{\mathbf{P}}, b(\mathbf{k})] \psi_{\mathbf{P}}) = 0 \quad \text{for } \mathbf{k} \rightarrow 0$$

$$\begin{aligned} & (\psi_{\mathbf{P}}, [H_{\mathbf{P}}, b(\mathbf{k})] \psi_{\mathbf{P}}) = \\ & = \left(\psi_{\mathbf{P}}, \left[\frac{(\mathbf{P}^{mes})^2}{2m} - \frac{\mathbf{P} \cdot \mathbf{P}^{mes}}{m} + g \int_0^\kappa \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) \frac{d^3 \mathbf{k}}{\sqrt{2|\mathbf{k}|}} + H^{mes}, b(\mathbf{k}) \right] \psi_{\mathbf{P}} \right) = \\ & = \left(\psi_{\mathbf{P}}, \left[\frac{(\mathbf{P}^{mes})^2}{2m} - \frac{\mathbf{P} \cdot \mathbf{P}^{mes}}{m}, b(\mathbf{k}) \right] \psi_{\mathbf{P}} \right) + \left(\psi_{\mathbf{P}}, \left[g \int_0^\kappa \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) \frac{d^3 \mathbf{k}}{\sqrt{2|\mathbf{k}|}}, b(\mathbf{k}) \right] \psi_{\mathbf{P}} \right) + \\ & + (\psi_{\mathbf{P}}, [H^{mes}, b(\mathbf{k})] \psi_{\mathbf{P}}) = \end{aligned}$$

$$= -\frac{\mathbf{k} \cdot (\psi_{\mathbf{P}}, \mathbf{P}^{mes} b(\mathbf{k}) \psi_{\mathbf{P}})}{2m} - \frac{(\psi_{\mathbf{P}}, b(\mathbf{k}) \mathbf{P}^{mes} \psi_{\mathbf{P}}) \cdot \mathbf{k}}{2m} + \frac{\mathbf{P} \cdot \mathbf{k}}{m} (\psi_{\mathbf{P}}, b(\mathbf{k}) \psi_{\mathbf{P}}) - \frac{g \cdot \chi_0^\kappa(\mathbf{k})}{\sqrt{2|\mathbf{k}|}} \|\psi_{\mathbf{P}}\|^2 - |\mathbf{k}| (\psi_{\mathbf{P}}, b(\mathbf{k}) \psi_{\mathbf{P}})$$

($\chi_0^\kappa(\mathbf{k})$ characteristic function of $\{\mathbf{k} : |\mathbf{k}| \leq \kappa\}$)

then

$$\begin{aligned} & -\frac{\mathbf{k} \cdot (\psi_{\mathbf{P}}, \mathbf{P}^{mes} b(\mathbf{k}) \psi_{\mathbf{P}})}{2m} - \frac{(\psi_{\mathbf{P}}, b(\mathbf{k}) \mathbf{P}^{mes} \psi_{\mathbf{P}}) \cdot \mathbf{k}}{2m} + \frac{\mathbf{P} \cdot \mathbf{k}}{m} (\psi_{\mathbf{P}}, b(\mathbf{k}) \psi_{\mathbf{P}}) - \frac{g \cdot \chi_0^\kappa(\mathbf{k})}{\sqrt{2|\mathbf{k}|}} \|\psi_{\mathbf{P}}\|^2 - |\mathbf{k}| (\psi_{\mathbf{P}}, b(\mathbf{k}) \psi_{\mathbf{P}}) = \\ & = 0 \\ & -\frac{\mathbf{k} \cdot (\psi_{\mathbf{P}}, \mathbf{P}^{mes} b(\mathbf{k}) \psi_{\mathbf{P}})}{m} - \frac{(\psi_{\mathbf{P}}, b(\mathbf{k}) \psi_{\mathbf{P}}) |\mathbf{k}|^2}{2m} + \frac{\mathbf{P} \cdot \mathbf{k}}{m} (\psi_{\mathbf{P}}, b(\mathbf{k}) \psi_{\mathbf{P}}) - \frac{g \cdot \chi_0^\kappa(\mathbf{k})}{\sqrt{2|\mathbf{k}|}} \|\psi_{\mathbf{P}}\|^2 - |\mathbf{k}| (\psi_{\mathbf{P}}, b(\mathbf{k}) \psi_{\mathbf{P}}) = \\ & = 0 \end{aligned}$$

$$\begin{aligned} & -\frac{\mathbf{k} \cdot (\psi_{\mathbf{P}}, \mathbf{P}^{mes} \psi_{\mathbf{P}})}{m} f_{\mathbf{P}}(\mathbf{k}) - \frac{|\mathbf{k}|^2}{2m} \|\psi_{\mathbf{P}}\|^2 f_{\mathbf{P}}(\mathbf{k}) + \frac{\mathbf{P} \cdot \mathbf{k}}{m} \|\psi_{\mathbf{P}}\|^2 f_{\mathbf{P}}(\mathbf{k}) - \frac{g \cdot \chi_0^\kappa(\mathbf{k})}{\sqrt{2|\mathbf{k}|}} \|\psi_{\mathbf{P}}\|^2 - |\mathbf{k}| \|\psi_{\mathbf{P}}\|^2 f_{\mathbf{P}}(\mathbf{k}) = \\ & = 0 \end{aligned}$$

Therefore the expected behavior is

$$f_{\mathbf{P}}(\mathbf{k}) \approx_{\mathbf{k} \rightarrow 0} -\frac{g \cdot \chi_0^\kappa(\mathbf{k})}{\sqrt{2|\mathbf{k}|}} \cdot \frac{1}{\left(|\mathbf{k}| + \frac{|\mathbf{k}|^2}{2m} - \frac{\mathbf{P} \cdot \mathbf{k}}{m} + \frac{\mathbf{k} \cdot (\psi_{\mathbf{P}}, \mathbf{P}^{mes} \psi_{\mathbf{P}})}{m \cdot \|\psi_{\mathbf{P}}\|^2}\right)} = -\frac{g \cdot \chi_0^\kappa(\mathbf{k})}{\sqrt{2|\mathbf{k}|}} \cdot \frac{1}{\left(|\mathbf{k}| + \frac{|\mathbf{k}|^2}{2m} - \frac{\mathbf{k} \cdot (\mathbf{P} - \overline{\mathbf{P}^{mes}})}{m}\right)}$$

and the coherent factor is labelled by $\mathbf{P}_1 = \mathbf{P} - \overline{\mathbf{P}^{mes}}$.

The argument proves that if the ground state is “coherent in the infrared region”, a non Fock state is necessarily. Starting from this result I operate a proper coherent transformation on the variables $\{b(\mathbf{k}), b^\dagger(\mathbf{k})\}$ of the hamiltonian $H_{\mathbf{P}}$ and I look for a ground state of the transformed hamiltonian in the Fock space with respect to $\{b(\mathbf{k}), b^\dagger(\mathbf{k})\}$.

Coherent transformation.

$$b(\mathbf{k}) \longrightarrow b(\mathbf{k}) - \frac{g}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} \quad \text{for } \mathbf{k} : 0 \leq |\mathbf{k}| \leq \kappa$$

using the *inter-twiner*

$$\mathcal{W}(\mathbf{P}) = W\left(\frac{\mathbf{P}_1}{m}\right) = e^{-g \int_0^\kappa \frac{b(\mathbf{k}) - b^\dagger(\mathbf{k})}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} \quad (7)$$

Transformed hamiltonian.

I rewrite $H_{\mathbf{P}}$, $\mathbf{P} = \mathbf{P}_1 + \mathbf{P}_2$, as

$$\begin{aligned} H_{\mathbf{P}} &= \frac{(\mathbf{P}_1 + \mathbf{P}_2 - \mathbf{P}^{mes})^2}{2m} + g \int_0^\kappa \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|} + H^{mes} = \\ &= \frac{(\mathbf{P}_1 + \mathbf{P}_2)^2}{2m} - \frac{(\mathbf{P}_1 + \mathbf{P}_2) \cdot \mathbf{P}^{mes}}{m} + \frac{\mathbf{P}^{mes^2}}{2m} + g \int_0^\kappa \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|} + H^{mes} = \\ &= \frac{(\mathbf{P}_1 + \mathbf{P}_2)^2}{2m} + \frac{\mathbf{P}^{mes^2}}{2m} - \frac{\mathbf{P}_2 \cdot \mathbf{P}^{mes}}{m} + \int_\kappa^\infty \left(|\mathbf{k}| - \mathbf{k} \cdot \frac{\mathbf{P}_1}{m} \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k + \\ &+ \int_0^\kappa \left(|\mathbf{k}| - \mathbf{k} \cdot \frac{\mathbf{P}_1}{m} \right) \left(b^\dagger(\mathbf{k}) + \frac{g}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} \right) \left(b(\mathbf{k}) + \frac{g}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} \right) d^3 k + \\ &- g^2 \int_0^\kappa \frac{1}{2|\mathbf{k}|^2(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} d^3 k \end{aligned}$$

and I act on it with the coherent transformation:

$$\begin{aligned} W\left(\frac{\mathbf{P}_1}{m}\right) H_{\mathbf{P}} W^\dagger\left(\frac{\mathbf{P}_1}{m}\right) &= \quad (8) \\ &= \frac{(\mathbf{P}_1 + \mathbf{P}_2)^2}{2m} + \frac{1}{2m} \left(\mathbf{P}^{mes} - g \int_0^\kappa \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) d^3 k + g^2 \int_0^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3 k \right)^2 + \\ &- \frac{\mathbf{P}_2}{m} \left(\mathbf{P}^{mes} - g \int_0^\kappa \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) d^3 k + g^2 \int_0^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3 k \right) + \\ &+ \int_0^\infty \left(|\mathbf{k}| - \mathbf{k} \cdot \frac{\mathbf{P}_1}{m} \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k - g^2 \int_0^\kappa \frac{1}{2|\mathbf{k}|^2(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} d^3 k \end{aligned}$$

Remark

Given $\phi_{\mathbf{P}} = W\left(\frac{\mathbf{P}_1}{m}\right) \psi_{\mathbf{P}}$ we have formally that

$$\mathbf{P}_2 = \mathbf{P} - \mathbf{P}_1 = \overline{\mathbf{P}^{mes}} = \frac{(\psi_{\mathbf{P}}, \mathbf{P}^{mes} \psi_{\mathbf{P}})}{\|\psi_{\mathbf{P}}\|^2} = \frac{(\phi_{\mathbf{P}}, W \mathbf{P}^{mes} W^\dagger \phi_{\mathbf{P}})}{\|\phi_{\mathbf{P}}\|^2} =$$

$$= \frac{1}{\|\phi_{\mathbf{P}}\|^2} (\phi_{\mathbf{P}}, \mathbf{P}^{mes} \phi_{\mathbf{P}}) - \frac{1}{\|\phi_{\mathbf{P}}\|^2} \left(\phi_{\mathbf{P}}, g \int_0^\kappa \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k \phi_{\mathbf{P}} \right) + g^2 \int_0^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k$$

Therefore $\mathbf{P}_1$ has to satisfy the equation

$$\mathbf{P} - \mathbf{P}_1 =$$

$$= \frac{(\phi_{\mathbf{P}}, \mathbf{P}^{mes} \phi_{\mathbf{P}})}{\|\phi_{\mathbf{P}}\|^2} - \frac{1}{\|\phi_{\mathbf{P}}\|^2} \left(\phi_{\mathbf{P}}, g \int_0^\kappa \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k \phi_{\mathbf{P}} \right) + g^2 \int_0^{\kappa_1} \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \quad (9)$$

where $\phi_{\mathbf{P}}$ is a ground state of the transformed hamiltonian $W\left(\frac{\mathbf{P}_1}{m}\right) H_{\mathbf{P}} W^\dagger\left(\frac{\mathbf{P}_1}{m}\right)$.

I define $\Pi_{\mathbf{P}} = \mathbf{P}^{mes} - g \int_0^\kappa \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k$ and by a substitution in the expressions (9) and (8) one arrives at

$$\begin{aligned} \bullet \mathbf{P}_2 &= \frac{1}{\|\phi_{\mathbf{P}}\|^2} (\phi_{\mathbf{P}}, \Pi_{\mathbf{P}} \phi_{\mathbf{P}}) + g^2 \int_0^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} \\ \bullet W\left(\frac{\mathbf{P}_1}{m}\right) H_{\mathbf{P}} W^\dagger\left(\frac{\mathbf{P}_1}{m}\right) &= \\ &= \frac{1}{2m} \Pi_{\mathbf{P}}^2 + \frac{1}{2m} \left(g^2 \int_0^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right)^2 + \frac{1}{m} \Pi_{\mathbf{P}} \cdot \left(g^2 \int_0^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right) + \\ &- \frac{1}{m} \left(g^2 \int_0^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right) \cdot \Pi_{\mathbf{P}} - \frac{1}{m} \frac{(\phi_{\mathbf{P}}, \Pi_{\mathbf{P}} \phi_{\mathbf{P}})}{\|\phi_{\mathbf{P}}\|^2} \cdot \Pi_{\mathbf{P}} + \int_0^\infty \left(|\mathbf{k}| - \mathbf{k} \cdot \frac{\mathbf{P}_1}{m} \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k + \\ &+ \text{constants} \end{aligned}$$

The transformed hamiltonian corresponds to:

$$H_{\mathbf{P}}^w = \frac{1}{2m} \left(\Pi_{\mathbf{P}} - \frac{(\phi_{\mathbf{P}}, \Pi_{\mathbf{P}} \phi_{\mathbf{P}})}{\|\phi_{\mathbf{P}}\|^2} \right)^2 + \int_0^\infty \left(|\mathbf{k}| - \mathbf{k} \cdot \frac{\mathbf{P}_1}{m} \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k + \text{constants} \quad (10)$$

Now I distinguish the case $\mathbf{P} = 0$ from $\mathbf{P} \neq 0$ in order to rigorously apply the iterative procedure to the transformed hamiltonians with an infrared cut-off σ_j , written in a canonic form analogous to the expression (10). In the first case, the coherent representation is explicit, since $\overline{\mathbf{P}^{mes}} = 0$ by symmetry and then $\mathbf{P}_1 = 0$, while in the second case the procedure is more lengthy.

2.1 Case $\mathbf{P} = 0$.

I perform the coherent transformation (7) on the hamiltonian $H_{\mathbf{P}=0, \sigma_j}$ with infrared cut-off σ_j :

$$\begin{aligned} H_{\epsilon^{\frac{j+1}{2}}}^w &\equiv e^{-g \int_{\epsilon^{\frac{j+1}{2}}}^{\kappa} \frac{b(\mathbf{k}) - b^\dagger(\mathbf{k})}{|\mathbf{k}|} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} H_{\mathbf{P}=0, \epsilon^{\frac{j+1}{2}}} e^{g \int_{\epsilon^{\frac{j+1}{2}}}^{\kappa} \frac{b(\mathbf{k}) - b^\dagger(\mathbf{k})}{|\mathbf{k}|} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} = \\ &= \frac{1}{2m} \left(\mathbf{P}^{mes} - g \int_{\epsilon^{\frac{j+1}{2}}}^{\kappa} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) d^3 k \right)^2 + \int |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k + c(j+1) \end{aligned}$$

where $c(j+1) \equiv -g^2 \int_{\epsilon^{\frac{j+1}{2}}}^{\kappa} \frac{1}{2|\mathbf{k}|^2} d^3 k$.

The domain of selfadjointness (s.a.) of the transformed hamiltonian $H_{\epsilon^{\frac{j+1}{2}}}^w$ coincides with $D^b \left(H_{\mathbf{P}=0, \epsilon^{\frac{j+1}{2}}} \right)$ (see an analogous proof in [1]).

Preliminaries

From the results of the previous chapter and by unitarity I can conclude that the following properties hold $\forall j$ (these properties are assumed in lemma A1, Appendix A):

- i) $H_{\epsilon^{\frac{j+1}{2}}}^w \big|_{F_{\epsilon^{\frac{j+1}{2}}}^+}$ has ground eigenvalue $E_{\mathbf{P}=0}^{\epsilon^{\frac{j+1}{2}}}$ with the corresponding gap bigger than $\frac{\epsilon^{\frac{j+1}{2}}}{2}$;
- ii) $H_{\epsilon^{\frac{j+1}{2}}}^w \big|_{F_{\epsilon^{\frac{j+2}{2}}}^+}$ has ground eigenvalue $E_{\mathbf{P}=0}^{\epsilon^{\frac{j+1}{2}}}$ with the corresponding gap bigger than $\frac{3\epsilon^{\frac{j+2}{2}}}{5}$.

Now, for the values of g allowed by lemma A1 and on the basis of the results of the lemma, starting from $\phi^\epsilon = e^{-g \int_{\epsilon}^{\kappa} \frac{b(\mathbf{k}) - b^\dagger(\mathbf{k})}{|\mathbf{k}|} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} \psi_{\mathbf{P}=0}^\epsilon$ and using an iteration analogous to the one of chapter 1, I construct the vector $\phi^{\epsilon^{\frac{j+2}{2}}}$ applying the spectral projector to $\phi^{\epsilon^{\frac{j+1}{2}}}$. We can point out that in the difference

$$\phi^{\epsilon \frac{j+2}{2}} - \phi^{\epsilon \frac{j+1}{2}} = -\frac{1}{2\pi i} \oint \sum_n \frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \left(-(\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^n \phi^{\epsilon \frac{j+1}{2}} dE(j+1) \quad (11)$$

the terms that we cannot evaluate (in norm) with a power of the cut-off $\epsilon^{\frac{j+1}{2}}$ with positive exponent are vectors for which for all the n factors in the difference $(\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} = H_{\epsilon \frac{j+2}{2}}^w + c(j+1) - c(j+2) - H_{\epsilon \frac{j+1}{2}}^w$ only the “mixed” terms are present:

$$\frac{g}{2m} \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k \cdot \Pi_{\epsilon \frac{j+1}{2}} + \Pi_{\epsilon \frac{j+1}{2}} \cdot \frac{g}{2m} \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k.$$

If there are these terms, the estimate of the following norm, given in lemma A1,

$$\left\| \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \left(-(\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \right) \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|$$

is only of order 1.

We can have a more precise estimate of the norm of $\phi^{\epsilon \frac{j+2}{2}} - \phi^{\epsilon \frac{j+1}{2}}$ examining the first factor on the right of the product (11)

$$\left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \left(-(\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \right) \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \phi^{\epsilon \frac{j+1}{2}}$$

and noting that, if for the mixed terms

$$\frac{g}{2m} \sum_i \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \left(\int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k \cdot \Pi_{\epsilon \frac{j+1}{2}}^i \right) \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \phi^{\epsilon \frac{j+1}{2}}$$

the following inequality were true

$$\begin{aligned} & \frac{g}{2m} \left\| \sum_i \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \left(\int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k \cdot \Pi_{\epsilon \frac{j+1}{2}}^i \right) \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \phi^{\epsilon \frac{j+1}{2}} \right\| \leq \\ & \leq \frac{1}{4} \cdot \frac{\epsilon^{\frac{j+1}{8}}}{2}, \end{aligned}$$

then we would have an estimate of order $\epsilon^{\frac{j+1}{8}}$ for $\left\| \phi^{\epsilon \frac{j+2}{2}} - \phi^{\epsilon \frac{j+1}{2}} \right\|$. This is due to the fact that, using the lemma A1, the norm of the sum of the quadratic terms (for a proper g) can be bounded by $\frac{\epsilon^{\frac{j+1}{8}}}{40}$ while the norm of the other factors of the product

is of order 1, in particular less than $\frac{1}{12}$. Therefore we would have

$$\begin{aligned}
& \left\| \sum_{n=1}^{\infty} \frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \left(-(\Delta H^w)_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^n \phi^{\epsilon^{\frac{j+1}{2}}} \right\| \leq \\
& \leq \left| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right) \right|^{\frac{1}{2}} \cdot \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right) \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+}^{\frac{1}{2}} \left(\epsilon^{\frac{j+1}{2}} \right)^{\frac{1}{4}} \sum_{n=1}^{\infty} \left(\frac{11}{40} \right) \left(\frac{1}{12} \right)^{n-1} \leq \\
& \leq \left| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right) \right| \cdot \left(\epsilon^{\frac{j+1}{2}} \right)^{\frac{1}{4}}
\end{aligned}$$

As it will be shown in corollary 2.4 (in the general case $\mathbf{P} \in \Sigma$), an estimate like $\left\| \phi^{\epsilon^{\frac{j+2}{2}}} - \phi^{\epsilon^{\frac{j+1}{2}}} \right\| \leq \epsilon^{\frac{j+1}{8}}$ implies the convergence of the sequence $\left\{ \phi^{\epsilon^{\frac{j+1}{2}}} \right\}$. Therefore it is crucial to prove the following inequality:

$$\begin{aligned}
& \frac{g}{2m} \left\| \sum_i \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \left(\int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k \cdot \Pi_{\epsilon^{\frac{j+1}{2}}}^i \right) \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \phi^{\epsilon^{\frac{j+1}{2}}} \right\| = \\
& = \frac{g}{2m} \left\| \sum_i \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \left(\int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3k \cdot \Pi_{\epsilon^{\frac{j+1}{2}}}^i \right) \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \phi^{\epsilon^{\frac{j+1}{2}}} \right\| \leq \\
& \leq \frac{1}{4} \cdot \frac{\epsilon^{\frac{j+1}{8}}}{2}.
\end{aligned}$$

Lemma 2.1

The following inequalities hold:

$$\begin{aligned}
\text{I)} & \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3k \cdot \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right\|^2 = \\
& = \left(\int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3k \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right| \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3k \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \leq
\end{aligned}$$

$$\begin{aligned}
&\leq \sqrt{112} \left| \left(\int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|} b^\dagger(\mathbf{k}) d^3k \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|} b^\dagger(\mathbf{k}) d^3k \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| \\
\text{II)} &\left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right| \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \leq \\
&\leq \sqrt{1 + \left(\frac{11\sqrt{\epsilon}}{10-11\sqrt{\epsilon}} \right)^2} \cdot \left| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right|
\end{aligned}$$

Proof

I define the wave functions $\zeta_I(z), \zeta_{II}(z)$ of $\int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|} b^\dagger(\mathbf{k}) d^3k \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}$ and $\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}$, in the spectral variable of $H_{\epsilon^{\frac{j+1}{2}}}^w - \text{Re}E(j+1)$. Note that:

- the operator $H_{\epsilon^{\frac{j+1}{2}}}^w - \text{Re}E(j+1)$, applied to the vector $\int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|} b^\dagger(\mathbf{k}) d^3k \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}$, takes values bigger or equal to $\frac{1}{20}\epsilon^{\frac{j+2}{2}}$ $\left(= \frac{3}{5}\epsilon^{\frac{j+2}{2}} - \frac{11}{20}\epsilon^{\frac{j+2}{2}} \right)$ because of lemma 1.2;
- the operator $H_{\epsilon^{\frac{j+1}{2}}}^w - \text{Re}E(j+1)$ takes values bigger or equal to $\frac{10-11\sqrt{\epsilon}}{20} \cdot \epsilon^{\frac{j+1}{2}}$ $\left(= \frac{1}{2}\epsilon^{\frac{j+1}{2}} - \frac{11}{20}\epsilon^{\frac{j+1}{2}} \right)$ if applied to the vector $\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}$ because of theorem 1.5

I write the scalar products I) and II) using the spectral representation of the operator $H_{\epsilon^{\frac{j+1}{2}}}^w - \text{Re}E(j+1)$ and getting rid of the remaining degrees of freedom. In the chosen spectral representation, the following inequalities are evident:

$$\begin{aligned}
&\left| \int \frac{|\zeta_{I,II}(z)|^2}{z - i\text{Im}(E(j+1))} dz \right| = \left\{ \left| \int \frac{z|\zeta_{I,II}(z)|^2}{z^2 + [\text{Im}(E(j+1))]^2} dz \right|^2 + [\text{Im}(E(j+1))]^2 \left| \int \frac{|\zeta_{I,II}(z)|^2}{z^2 + [\text{Im}(E(j+1))]^2} dz \right|^2 \right\}^{\frac{1}{2}} \geq \\
&\geq \left| \int \frac{|\zeta_{I,II}(z)|^2}{\sqrt{z^2 + [\text{Im}(E(j+1))]^2}} \cdot \frac{z}{\sqrt{z^2 + [\text{Im}(E(j+1))]^2}} dz \right| \geq \left| \int \frac{|\zeta_{I,II}(z)|^2}{\sqrt{z^2 + [\text{Im}(E(j+1))]^2}} \cdot \frac{z}{\sqrt{z^2 + |E(j+1)|^2}} dz \right| \geq \\
&\geq \frac{z_{\min}}{\sqrt{z_{\min}^2 + |E(j+1)|^2}} \cdot \left| \int \frac{|\zeta_{I,II}(z)|^2}{\sqrt{z^2 + [\text{Im}(E(j+1))]^2}} dz \right| \geq \\
&\geq \frac{z_{\min}}{\sqrt{z_{\min}^2 + |E(j+1)|^2}} \cdot \left| \int \frac{|\zeta_{I,II}(z)|^2}{\sqrt{z^2 + |\mathbf{k}|^2 + 2|\mathbf{k}|z + [\text{Im}(E(j+1))]^2}} dz \right|
\end{aligned}$$

It follows that:

- in the case I), being $z_{min} \geq \frac{1}{20}\epsilon^{\frac{j+2}{2}}$, $\frac{z_{min}}{\sqrt{z_{min}^2 + |E(j+1)|^2}} \geq \frac{1}{\sqrt{112}}$

$$\left| \int \frac{|\zeta_I(z)|^2}{\sqrt{z^2 + [Im(E(j+1))]^2}} dz \right| \leq \sqrt{112} \cdot \left| \int \frac{|\zeta_I(z)|^2}{z - iIm(E(j+1))} dz \right|$$
- in the case II), being $z_{min} \geq \frac{10-11\sqrt{\epsilon}}{20} \cdot \epsilon^{\frac{j+1}{2}}$, $\frac{z_{min}}{\sqrt{z_{min}^2 + |E(j+1)|^2}} \geq \frac{1}{\sqrt{1 + \left(\frac{11\sqrt{\epsilon}}{10-11\sqrt{\epsilon}}\right)^2}}$

$$\left| \int \frac{|\zeta_{II}(z)|^2}{\sqrt{z^2 + |\mathbf{k}|^2 + 2|\mathbf{k}|z + [Im(E(j+1))]^2}} dz \right| \leq \sqrt{1 + \left(\frac{11\sqrt{\epsilon}}{10-11\sqrt{\epsilon}}\right)^2} \cdot \left| \int \frac{|\zeta_{II}(z)|^2}{z - iIm(E(j+1))} dz \right| =$$

$$= Q(\epsilon) \cdot \left| \int \frac{|\zeta_{II}(z)|^2}{z - iIm(E(j+1))} dz \right| \quad \text{where } Q(\epsilon) \equiv \sqrt{1 + \left(\frac{11\sqrt{\epsilon}}{10-11\sqrt{\epsilon}}\right)^2}.$$

Lemma 2.2

Taking into account lemma 2.1, we have that:

$$\left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3k \cdot \Pi_{\epsilon^{\frac{j+1}{2}}}^i \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \phi^{\epsilon^{\frac{j+1}{2}}} \right\|^2 \leq$$

$$\leq 2 \cdot Q(\epsilon) \cdot \sqrt{112} \cdot \left| \frac{1}{E_{\epsilon^{\frac{j+1}{2}}} - E(j+1)} \right| \cdot \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{2|\mathbf{k}|^3} \left| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right) \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| d^3k$$

Proof

$$\left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3k \cdot \Pi_{\epsilon^{\frac{j+1}{2}}}^i \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \phi^{\epsilon^{\frac{j+1}{2}}} \right\|^2 =$$

$$= \left| \frac{1}{E_{\epsilon^{\frac{j+1}{2}}} - E(j+1)} \right| \cdot \left(\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3k \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \right.$$

$$\left. \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \cdot \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3k \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) =$$

$$= \left| \frac{1}{E_{\epsilon^{\frac{j+1}{2}}} - E(j+1)} \right| \cdot \left(\int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3k \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right| \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3k \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \leq$$

$$\leq \sqrt{112} \left| \frac{1}{E^\epsilon \frac{j+1}{2} - E(j+1)} \right| \cdot \left| \left(\int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3 k \Pi_{\epsilon \frac{j+1}{2}}^i \phi^{\epsilon \frac{j+1}{2}} \right), \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right) \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3 k \Pi_{\epsilon \frac{j+1}{2}}^i \phi^{\epsilon \frac{j+1}{2}} \right) \right|$$

Starting from the expression $H_{\epsilon \frac{j+1}{2}}^w = \frac{1}{2m} \Pi_{\epsilon \frac{j+1}{2}}^2 + \int |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k + c(j+1)$, for $\mathbf{k} : |\mathbf{k}| \leq \epsilon \frac{j+1}{2}$, the following identity holds in distributional sense

$$\begin{aligned} \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right) b^\dagger(\mathbf{k}) &= b^\dagger(\mathbf{k}) \left(\frac{1}{\frac{1}{2m} \left(\Pi_{\epsilon \frac{j+1}{2}} + \mathbf{k} \right)^2 + \int |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) + |\mathbf{k}| + c(j+1) - E(j+1)} \right) = \\ &= b^\dagger(\mathbf{k}) \left(\frac{1}{\frac{1}{2m} \left(\Pi_{\epsilon \frac{j+1}{2}}^2 + 2\mathbf{k} \cdot \Pi_{\epsilon \frac{j+1}{2}} + \mathbf{k}^2 \right) + \int |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) + |\mathbf{k}| + c(j+1) - E(j+1)} \right) \end{aligned}$$

And also, for the assumptions made on m , ϵ and $\left| E^{\epsilon \frac{j+1}{2}} - E(j+1) \right|$, and since $\epsilon \frac{j+2}{2} \leq |\mathbf{k}| \leq \epsilon \frac{j+1}{2}$, the following bound holds in the subspace $F_{\epsilon \frac{j+2}{2}}^+$

$$\left\| \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} \left(\frac{1}{m} \mathbf{k} \cdot \Pi_{\epsilon \frac{j+1}{2}} + \frac{\mathbf{k}^2}{2m} \right) \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon \frac{j+2}{2}}^+} \leq \frac{1}{2}$$

which well defines the series expansion

$$\begin{aligned} b^\dagger(\mathbf{k}) \left(\frac{1}{\frac{1}{2m} \left(\Pi_{\epsilon \frac{j+1}{2}}^2 + 2\mathbf{k} \cdot \Pi_{\epsilon \frac{j+1}{2}} + \mathbf{k}^2 \right) + \int |\mathbf{k}| b^\dagger(\mathbf{k}) b(\mathbf{k}) + |\mathbf{k}| - E(j+1)} \right) &= \\ = b^\dagger(\mathbf{k}) \left(\left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w + |\mathbf{k}| - E(j+1)} \right) \sum_{n=0}^{+\infty} \left(- \left(\frac{1}{m} \mathbf{k} \cdot \Pi_{\epsilon \frac{j+1}{2}} + \frac{\mathbf{k}^2}{2m} \right) \frac{1}{H_{\epsilon \frac{j+1}{2}}^w + |\mathbf{k}| - E(j+1)} \right)^n \right) \end{aligned}$$

from which

$$\begin{aligned} \left(\int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3 k \Pi_{\epsilon \frac{j+1}{2}}^i \phi^{\epsilon \frac{j+1}{2}} \right), \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right) \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3 k \Pi_{\epsilon \frac{j+1}{2}}^i \phi^{\epsilon \frac{j+1}{2}} \right) &= \quad (12) \\ = \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{k^i}{2|\mathbf{k}|^3} \left(\Pi_{\epsilon \frac{j+1}{2}}^i \phi^{\epsilon \frac{j+1}{2}} \right), \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w + |\mathbf{k}| - E(j+1)} \right) \sum_{n=0}^{\infty} \left(- \left(\frac{1}{m} \mathbf{k} \cdot \Pi_{\epsilon \frac{j+1}{2}} + \frac{\mathbf{k}^2}{2m} \right) \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w + |\mathbf{k}| - E(j+1)} \right) \right)^n \Pi_{\epsilon \frac{j+1}{2}}^i \phi^{\epsilon \frac{j+1}{2}} d^3 k &= \end{aligned}$$

$$= \sum_{n=0}^{\infty} \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j+2}{2}}} \frac{k^{i^2}}{2|\mathbf{k}|^3} \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right) \left(- \left(\frac{1}{m} \mathbf{k} \cdot \Pi_{\epsilon^{\frac{j+1}{2}}} + \frac{\mathbf{k}^2}{2m} \right) \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right) \right) \right)^n \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} d^3 k$$

I prove that the module of the n^{th} term of the series can be reduced to the one of the first term, so that the whole sum is of the same order of the term at $n = 0$.

Exploiting the Schwartz inequality and the identity

$$\begin{aligned} & \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right) \left((-) \left(\frac{1}{m} \mathbf{k} \cdot \Pi_{\epsilon^{\frac{j+1}{2}}} + \frac{\mathbf{k}^2}{2m} \right) \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right) \right)^n = \\ & = \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} \cdot \\ & \cdot \left(\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} (-) \left(\frac{1}{m} \mathbf{k} \cdot \Pi_{\epsilon^{\frac{j+1}{2}}} + \frac{\mathbf{k}^2}{2m} \right) \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} \right)^n \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} \end{aligned}$$

we have that

$$\begin{aligned} & \left| \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j+2}{2}}} \frac{k^{i^2}}{2|\mathbf{k}|^3} \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right) \left((-) \left(\frac{1}{m} \mathbf{k} \cdot \Pi_{\epsilon^{\frac{j+1}{2}}} + \frac{\mathbf{k}^2}{2m} \right) \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right) \right) \right)^n \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} d^3 k \right| \leq \\ & \leq \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j+2}{2}}} \frac{k^{i^2}}{2|\mathbf{k}|^3} \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right\|^2 \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} \left(\frac{1}{m} \mathbf{k} \cdot \Pi_{\epsilon^{\frac{j+1}{2}}} + \frac{\mathbf{k}^2}{2m} \right) \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} \right\|^n d^3 k \\ & \quad \text{(remember that } \left\| \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} \right]^{\dagger} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right\| = \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right\|) \end{aligned}$$

Therefore, thanks to the series expansion and to the lemma 2.1, the module of the scalar product (12) is bounded by

$$\begin{aligned} & \sum_{n=0}^{\infty} \left\{ \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j+2}{2}}} \frac{k^{i^2}}{2|\mathbf{k}|^3} \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right\|^2 d^3 k \right\} \left(\frac{1}{2} \right)^n \leq \\ & \leq 2 \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j+2}{2}}} \frac{k^{i^2}}{2|\mathbf{k}|^3} \cdot \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right)^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right\|^2 d^3 k \leq \\ & \leq 2 \cdot Q(\epsilon) \cdot \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j+2}{2}}} \frac{k^{i^2}}{2|\mathbf{k}|^3} \cdot \left\| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w + |\mathbf{k}| - E(j+1)} \right) \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right\| d^3 k \end{aligned}$$

Theorem 2.3

$$\left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \frac{g}{2m} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} b^\dagger(\mathbf{k}) d^3k \cdot \Pi_{\epsilon^{\frac{j+1}{2}}} \phi^{\epsilon^{\frac{j+1}{2}}} \left(\frac{1}{E_{\epsilon^{\frac{j+1}{2}}} - E(j+1)} \right)^{\frac{1}{2}} \right\| \leq \frac{\epsilon^{\frac{j+1}{8}}}{8} \quad (13)$$

Proof

Due to the results of the lemma 2.2 the inequality (13) is true if the following estimate holds:

$$2\sqrt{112} \cdot Q(\epsilon) \left| \left(\frac{1}{E_{\epsilon^{\frac{j+1}{2}}} - E(j+1)} \right) \frac{g^2}{4m^2} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{k^2}{2|\mathbf{k}|^3} d^3k \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right) \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| \leq \left(\frac{\epsilon^{\frac{j+1}{8}}}{24} \right)^2$$

For this purpose I prove by induction that

$$\frac{g^2}{4m^2} \left| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right) \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| < \frac{11}{\sqrt{112} \cdot Q(\epsilon) 40\pi} \cdot \frac{1}{(4^4 \cdot 24)^2 \cdot \epsilon^{\frac{1}{4}(j+1)}} \quad (14)$$

I note that the inequality (14) implies the thesis of the theorem and therefore, as discussed in pag.29, also the bound $\left\| \phi^{\epsilon^{\frac{j+2}{2}}} - \phi^{\epsilon^{\frac{j+1}{2}}} \right\| \leq \epsilon^{\frac{j+1}{8}}$.

In order to prove the inequality (14) I start analyzing

$$\begin{aligned} & \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right) \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) = \\ & = \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - H_{\epsilon^{\frac{j}{2}}}^w + H_{\epsilon^{\frac{j}{2}}}^w - E(j) + E(j) - E(j+1)} \right) \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \end{aligned} \quad (15)$$

In lemma A1 I defined $(\Delta H_{\mathbf{P}}^w)_{\epsilon^{\frac{j+1}{2}}}^j \equiv \widehat{H}_{\mathbf{P}, \epsilon^{\frac{j+1}{2}}}^w + c_{\mathbf{P}}(j) - \widehat{c}_{\mathbf{P}}(j+1) - H_{\mathbf{P}, \epsilon^{\frac{j}{2}}}^w$, that for $\mathbf{P} = 0$ becomes $(\Delta H^w)_{\epsilon^{\frac{j+1}{2}}}^j \equiv H_{\epsilon^{\frac{j+1}{2}}}^w + c(j) - c(j+1) - H_{\epsilon^{\frac{j}{2}}}^w$.

Now I define $\tilde{\Delta}(H^w)^{\epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} \equiv \Delta H^{w, \epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} - c(j) + c(j+1) + E(j) - E(j+1)$ and I observe that choosing a proper value $\overline{g}_1$ for g , the following inequalities hold for all j :

$$\begin{aligned}
& \bullet \left\| \left(\frac{1}{H^{w, \epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} - E(j)} \right)^{\frac{1}{2}} (-) \Delta(H^w)^{\epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} \left(\frac{1}{H^{w, \epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} - E(j)} \right)^{\frac{1}{2}} \right\|_{F^{+}_{\epsilon \frac{j+1}{2}}} < \frac{1}{6} \\
& \bullet \left\| \left(\frac{1}{H^{w, \epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} - E(j)} \right)^{\frac{1}{2}} (-) (-c(j) + c(j+1)) \left(\frac{1}{H^{w, \epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} - E(j)} \right)^{\frac{1}{2}} \right\|_{F^{+}_{\epsilon \frac{j+1}{2}}} < \frac{1}{6} \\
& \bullet \left\| \left(\frac{1}{H^{w, \epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} - E(j)} \right)^{\frac{1}{2}} (-) (E(j) - E(j+1)) \left(\frac{1}{H^{w, \epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} - E(j)} \right)^{\frac{1}{2}} \right\|_{F^{+}_{\epsilon \frac{j+1}{2}}} < \frac{1}{6}
\end{aligned}$$

Observations

The first inequality is the content of lemma A1. The second is true as $c(j+1) - c(j) = -2\pi g^2 \epsilon^{\frac{j}{2}} (1 - \sqrt{\epsilon})$. Concerning the third inequality we arrive at the given estimate by re-conducting $|E(j) - E(j+1)|$ to $\left| E^{\epsilon \frac{j+1}{2}} - E^{\epsilon \frac{j}{2}} \right|$ and using the generalized lemma 1.3, for g sufficiently small and uniform in j .

In conclusion:

$$\left\| \left(\frac{1}{H^{w, \epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} - E(j)} \right)^{\frac{1}{2}} (-) \tilde{\Delta}(H^w)^{\epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} \left(\frac{1}{H^{w, \epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} - E(j)} \right)^{\frac{1}{2}} \right\|_{F^{+}_{\epsilon \frac{j+1}{2}}} < \frac{1}{2}$$

It is therefore possible to expand in series the expression (15):

$$\left(\Pi^i_{\epsilon \frac{j+1}{2}} \phi^{\epsilon \frac{j+1}{2}}, \left(\frac{1}{H^{w, \epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} - E(j)} \right) \sum_{n=0} \left((-) \tilde{\Delta}(H^w)^{\epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} \left(\frac{1}{H^{w, \epsilon \frac{j}{2}}_{\epsilon \frac{j+1}{2}} - E(j)} \right) \right)^n \Pi^i_{\epsilon \frac{j+1}{2}} \phi^{\epsilon \frac{j+1}{2}} \right)$$

$$\begin{aligned}
&= \sum_{n=0} \left(\left[\left(\frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right)^{\frac{1}{2}} \right]^{\dagger} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \right. \\
&\quad \left. \left[\left(\frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right)^{\frac{1}{2}} (-1) \tilde{\Delta} (H^w)_{\epsilon^{\frac{j+1}{2}}}^{\frac{j}{2}} \left(\frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right)^{\frac{1}{2}} \right]^n \left(\frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right)^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right)
\end{aligned}$$

so that we obtain the bound

$$\left| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right) \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| \leq 2 \left| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| \quad (16)$$

Proof by induction of the inequality (14)

For $j = 1, 2, 3, \dots, 7$ and g sufficiently small (that I can always assume as bigger or equal to $\bar{g}_1$, redefining $\bar{g}_1$), the bound (14) is valid. If one assumes that it holds for $j = 1, \dots, j-1, j \in g = \bar{g} \equiv \min(\bar{g}_1, \bar{g}_2, 1)$ where

$$\bar{g}_2 = \frac{2m^2}{W} \cdot \frac{1}{\sqrt{112} \cdot Q(\epsilon)} \frac{11}{40\pi S} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{1}{4}}} \quad \text{and}$$

$$\begin{aligned}
&\bullet W = 2 \left\{ \left[2m\sqrt{2m} \cdot (\epsilon^{-\frac{1}{4}}\sqrt{20}) + 2m \left((\epsilon^{-\frac{1}{4}}\sqrt{20}) + 2\sqrt{40\pi} \right) \right] \cdot \sqrt{\frac{11}{\sqrt{112} \cdot 40\pi} \cdot \frac{1}{(24 \cdot 4^4)^2}} \right\} + \\
&\quad + \left\{ 2m \left(\epsilon^{-\frac{1}{4}}\sqrt{20} \right)^2 + \left((\epsilon^{-\frac{1}{4}}\sqrt{20\pi}) + 2\sqrt{40\pi} \right)^2 + 2 \cdot \sqrt{2m} \cdot (\epsilon^{-\frac{1}{4}}\sqrt{20}) \left((\epsilon^{-\frac{1}{4}}\sqrt{20\pi}) + 2\sqrt{40\pi} \right) \right\} \\
&\bullet S = \sum_{n=0}^{\infty} \left(2Q(\epsilon) \cdot \epsilon^{\frac{1}{4}} \right)^n + \frac{20\epsilon^{\frac{1}{4}}}{11W} \cdot \left(2Q(\epsilon) \epsilon^{\frac{1}{12}} \right)^6
\end{aligned}$$

then the bound (14) holds for $j+1$, where $j \geq 7$.

The starting point of the procedure is the inequality (16). Adding and subtracting $\Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}}$ on the left and on the right of the scalar product, I bound the new terms that I get, using elementary properties of the scalar product:

$$\begin{aligned}
& 2 \left| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| \leq \\
& \leq 2 \left| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right) \right| + \tag{17}
\end{aligned}$$

$$+ 4 \left| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right| + \tag{18}$$

$$+ 2 \left| \left(\Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right| \tag{19}$$

Observation

The above expression displays the inductive procedure to arrive at the thesis of the theorem, as the quantity (19) has the same form of the (15), up to a factor $2 \cdot Q(\epsilon)$, where the cut-off is $\epsilon^{\frac{j}{2}}$ instead of $\epsilon^{\frac{j+1}{2}}$.

In the bound of the expressions (17) and (18) I will use:

$$1) \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} = \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - g \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} \frac{\mathbf{q}}{\sqrt{2}|\mathbf{q}|^{\frac{3}{2}}} \left(b(\mathbf{q}) + b^\dagger(\mathbf{q}) \right) d^3q \phi^{\epsilon^{\frac{j+1}{2}}}$$

$$2) \left\| \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} q^i b(\mathbf{k}) \frac{d^3q}{|\mathbf{q}| \sqrt{2}|\mathbf{q}|} \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \right\|_{F_{\epsilon^{\frac{j+1}{2}}}^+}^{\frac{1}{2}} \leq \sqrt{40\pi} \cdot \epsilon^{\frac{j}{4}}$$

(analogous to the (a1) of the lemma A1 (Appendix A));

3)

$$\left\| \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \right\|_{F_{\epsilon^{\frac{j+1}{2}}}^+}^{\frac{1}{2}} \leq \left(\epsilon^{-\frac{1}{4}} \sqrt{20} \right) \cdot \epsilon^{-\frac{j}{4}}$$

3bis)

$$\begin{aligned}
& \left\| \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 q}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right|^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+1}{2}}}^+} \leq \sqrt{\pi} \epsilon^{\frac{j}{2}} \left\| \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right|^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+1}{2}}}^+} + \\
& + \left\| \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} q^i b(\mathbf{q}) \frac{d^3 q}{|\mathbf{q}| \sqrt{2} |\mathbf{q}|} \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right|^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+1}{2}}}^+} \leq \\
& \leq \sqrt{\pi} \epsilon^{\frac{j}{2}} \left(\epsilon^{-\frac{1}{4}} \sqrt{20} \right) \cdot \epsilon^{-\frac{j}{4}} + \sqrt{40\pi} \cdot \epsilon^{\frac{j}{4}} \leq \left(\left(\epsilon^{-\frac{1}{4}} \sqrt{20\pi} \right) + \sqrt{40\pi} \right) \cdot \epsilon^{\frac{j}{4}}
\end{aligned}$$

from which

$$\left\| \left[\left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right|^{\frac{1}{2}} \right]^\dagger \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} k^i \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) \frac{d^3 q}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \right\|_{F_{\epsilon^{\frac{j+1}{2}}}^+} \leq \left(\left(\epsilon^{-\frac{1}{4}} \sqrt{20\pi} \right) + 2\sqrt{40\pi} \right) \cdot \epsilon^{\frac{j}{4}}$$

4) for the inductive hypothesis we have that:

$$\begin{aligned}
\text{i) } & \frac{g^2}{4m^2} \left\| \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right|^{\frac{1}{2}} d^3 k \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right\|^2 \leq \frac{g^2}{4m^2} Q(\epsilon) \left| \left(\Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right) \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right| \leq \\
& \leq \frac{11}{\sqrt{112 \cdot 40\pi}} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{j}{4}}} \\
\text{ii) } & \left\| \phi^{\epsilon^{\frac{j+1}{2}}} - \phi^{\epsilon^{\frac{j}{2}}} \right\| < \epsilon^{\frac{j}{8}} = \sigma_{j-1}^{\frac{1}{4}} \text{ from which } \left\| \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right\| < \sqrt{2m} \cdot \left(E^{\epsilon^{\frac{j}{2}}} - c(j) \right)^{\frac{1}{2}} \epsilon^{\frac{j}{8}} < \\
& < \sqrt{2m} \cdot \epsilon^{\frac{j}{8}}
\end{aligned}$$

(for the initial assumptions $\left(E^{\epsilon^{\frac{j}{2}}} - c(j) \right) < 1$)

Bound of (17)

$$\left| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right), \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right) \right| \leq$$

$$\begin{aligned}
&= \left| \left(\Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \left(\Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right) \right| + \\
&+ \left| \left(g \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k \phi^{\epsilon^{\frac{j+1}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| g \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| + \\
&+ 2 \left| \left(g \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k \phi^{\epsilon^{\frac{j+1}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \left(\Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right) \right|
\end{aligned}$$

The (17) is therefore bounded by

$$2 \left\{ 2m \cdot \left(\epsilon^{-\frac{1}{4}} \sqrt{20} \right)^2 + \left(\left(\epsilon^{-\frac{1}{4}} \sqrt{20\pi} \right) + 2\sqrt{40\pi} \right)^2 + 2 \cdot \left(\epsilon^{-\frac{1}{4}} \sqrt{20} \right) \cdot \left(\left(\epsilon^{-\frac{1}{4}} \sqrt{20\pi} \right) + 2\sqrt{40\pi} \right) \right\} \cdot \epsilon^{-\frac{j}{4}}$$

Bound of (18).

$$\begin{aligned}
&\left| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right| \leq \\
&\leq \left| \left(\Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right| + \tag{18.1}
\end{aligned}$$

$$+ \left| \left(g \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k \phi^{\epsilon^{\frac{j+1}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right| \tag{18.2}$$

I examine the two terms on the right side:

18.1)

$$\begin{aligned}
&\left| \left(\Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right| = \\
&= \left| \left(\Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right|^{\frac{1}{2}} \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right|^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right| \leq
\end{aligned}$$

$$\begin{aligned}
&\leq \left\| \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right\| \cdot \left\| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right\|_{F_{\epsilon^{\frac{j+1}{2}}}^+}^{\frac{1}{2}} \cdot \left\| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right\|^{\frac{1}{2}} d^3 k \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right\| \\
&\leq \epsilon^{\frac{j}{8}} \cdot \sqrt{2m} \cdot (\epsilon^{-\frac{1}{4}} \sqrt{20}) \cdot \epsilon^{-\frac{j}{4}} \cdot \frac{2m}{g} \cdot \sqrt{\frac{11}{\sqrt{112 \cdot 40\pi}}} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{j}{4}}} \\
&18.2)
\end{aligned}$$

$$\begin{aligned}
&\left| \left(g \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3 k \phi^{\epsilon^{\frac{j+1}{2}}}, \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right| \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right) \right| \leq \\
&\leq \left\| g \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} k^i (b(\mathbf{k}) + b^\dagger(\mathbf{k})) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2}|\mathbf{k}|} \left| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right|_{F_{\epsilon^{\frac{j+1}{2}}}^+}^{\frac{1}{2}} \cdot \left\| \frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right\|^{\frac{1}{2}} d^3 k \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j}{2}}} \right\| \leq \\
&\leq g \cdot \left((\epsilon^{-\frac{1}{4}} \sqrt{20}) + 2\sqrt{40\pi} \right) \cdot \epsilon^{\frac{j}{4}} \cdot \frac{2m}{g} \cdot \sqrt{\frac{11}{\sqrt{112 \cdot 40\pi}}} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{j}{4}}} \leq \\
&\leq 2m \cdot \left((\epsilon^{-\frac{1}{4}} \sqrt{20}) + 2\sqrt{40\pi} \right) \cdot \sqrt{\frac{11}{\sqrt{112 \cdot 40\pi}}} \cdot \frac{1}{(24 \cdot 4^4)^2} \cdot \epsilon^{\frac{j}{8}}
\end{aligned}$$

We conclude that (18) is bounded by

$$4 \left\{ \left[\frac{2m}{g} \sqrt{2m} \cdot (\epsilon^{-\frac{1}{4}} \sqrt{20}) + 2m \left((\epsilon^{-\frac{1}{4}} \sqrt{20}) + 2\sqrt{40\pi} \right) \right] \cdot \sqrt{\frac{11}{\sqrt{112 \cdot 40\pi}}} \cdot \frac{1}{(24 \cdot 4^4)^2} \right\} \cdot \epsilon^{-\frac{j}{4}}$$

From the previous estimates we have that for $g = \bar{g}$ ($\bar{g} \leq 1$), fixed from the beginning of the theorem, the sum of (17) and (18) times $\frac{\bar{g}^2}{4m^2}$ is less than

$$\begin{aligned}
&\frac{\bar{g}^2}{4m^2} \cdot \left[2 \left\{ 2m \left(\epsilon^{-\frac{1}{4}} \sqrt{20} \right)^2 + \left((\epsilon^{-\frac{1}{4}} \sqrt{20\pi}) + 2\sqrt{40\pi} \right)^2 + 2 \cdot \sqrt{2m} \cdot (\epsilon^{-\frac{1}{4}} \sqrt{20}) \cdot \left((\epsilon^{-\frac{1}{4}} \sqrt{20\pi}) + 2\sqrt{40\pi} \right) \right\} + \right. \\
&+ 4 \left\{ \left[\frac{2m}{g} \sqrt{2m} \cdot (\epsilon^{-\frac{1}{4}} \sqrt{20}) + 2m \left((\epsilon^{-\frac{1}{4}} \sqrt{20}) + 2\sqrt{40\pi} \right) \right] \cdot \sqrt{\frac{11}{\sqrt{112 \cdot 40\pi}}} \cdot \frac{1}{(24 \cdot 4^4)^2} \right\} \cdot \epsilon^{-\frac{j}{4}} \leq \\
&\leq \frac{\bar{g}}{2m^2} \left[\left\{ 2m \left(\epsilon^{-\frac{1}{4}} \sqrt{20} \right)^2 + \left((\epsilon^{-\frac{1}{4}} \sqrt{20\pi}) + 2\sqrt{40\pi} \right)^2 + 2 \cdot \sqrt{2m} \cdot (\epsilon^{-\frac{1}{4}} \sqrt{20}) \cdot \left((\epsilon^{-\frac{1}{4}} \sqrt{20\pi}) + 2\sqrt{40\pi} \right) \right\} + \right. \\
&+ 2 \left\{ \left[2m \sqrt{2m} \cdot (\epsilon^{-\frac{1}{4}} \sqrt{20}) + 2m \left((\epsilon^{-\frac{1}{4}} \sqrt{20}) + 2\sqrt{40\pi} \right) \right] \cdot \sqrt{\frac{11}{\sqrt{112 \cdot 40\pi}}} \cdot \frac{1}{(24 \cdot 4^4)^2} \right\} \cdot \epsilon^{-\frac{j}{4}} \leq \\
&\leq \frac{11}{40\pi S} \cdot \frac{1}{\sqrt{112 \cdot Q(\epsilon)}} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{j}{4}(j+1)}}.
\end{aligned}$$

At this point the procedure has to be repeated nearly up to the cut-off $\epsilon^{\frac{j+1}{8}}$, precisely the cut-off $\epsilon^{\frac{[\hat{j}+1]+1}{2}}$ where $\frac{\hat{j}+1}{2} = \frac{j+1}{8}$ e $[\hat{j}+1]$ =integer part of $\hat{j}+1$:

$$\begin{aligned}
& \frac{\bar{g}^2}{4m^2} \left| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right) \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| \leq \\
& \leq \frac{11}{40\pi S} \cdot \frac{1}{\sqrt{112} \cdot Q(\epsilon)} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{1}{4}(j+1)}} + \frac{\bar{g}^2}{4m^2} \cdot 2 \left| \left(\Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right) \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| \leq \\
& \leq \frac{11}{40\pi S} \cdot \frac{1}{\sqrt{112} \cdot Q(\epsilon)} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{1}{4}(j+1)}} + \frac{\bar{g}^2}{4m^2} \cdot 2Q(\epsilon) \left| \left(\Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j}{2}}}^w - E(j)} \right) \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| \leq \\
& \leq \frac{11}{40\pi S} \cdot \frac{1}{\sqrt{112} \cdot Q(\epsilon)} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{1}{4}(j+1)}} + \frac{\bar{g}^2}{4m^2} \cdot 2^2 Q(\epsilon) \left| \left(\Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j-1}{2}}}^w - E(j-1)} \right) \Pi_{\epsilon^{\frac{j}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| \leq \\
& \leq \frac{11}{40\pi S} \cdot \frac{1}{\sqrt{112} \cdot Q(\epsilon)} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{1}{4}(j+1)}} \left(1 + (2Q(\epsilon) \cdot \epsilon^{\frac{1}{4}}) + \dots + (2Q(\epsilon) \cdot \epsilon^{\frac{1}{4}})^{\frac{3j+3}{4}} \right) + \\
& + \frac{\bar{g}^2}{4m^2} \cdot \frac{20}{11} \cdot \frac{2\epsilon^{\frac{1}{2}}}{\epsilon^{\frac{1}{4}(j+1)}} (2Q(\epsilon))^{\frac{3j+3}{4}} \cdot \epsilon^{\frac{j-3}{8}}
\end{aligned}$$

the formula above can be explained with:

- $j - [\hat{j} + 1] \leq j + 1 - \hat{j} - 1 = \frac{3j+3}{4}$
- being $Q(\epsilon) > 1$ $(2Q(\epsilon))^{\frac{3j+3}{4}} \geq (2Q(\epsilon))^{j - [\hat{j} + 1]}$
- $\frac{1}{\epsilon^{\frac{[\hat{j}+1]+1}{2}}} \leq \frac{1}{\epsilon^{\frac{\frac{j+1}{4}+1}{2}}} = \frac{1}{\epsilon^{\frac{j+5}{8}}} = \frac{1}{\epsilon^{\frac{1}{4}(j+1)}} \cdot \epsilon^{\frac{j-3}{8}}$

then

$$\begin{aligned}
& \frac{\bar{g}^2}{4m^2} \left| \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}}, \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right) \Pi_{\epsilon^{\frac{j+1}{2}}}^i \phi^{\epsilon^{\frac{j+1}{2}}} \right) \right| \leq \\
& \leq \frac{11}{40\pi S} \cdot \frac{1}{\sqrt{112} \cdot Q(\epsilon)} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{1}{4}(j+1)}} \left(1 + (2Q(\epsilon) \cdot \epsilon^{\frac{1}{4}})^1 + \dots + (2Q(\epsilon) \cdot \epsilon^{\frac{1}{4}})^{j - [\hat{j} + 1]} \right) + \\
& + \frac{\bar{g}^2}{4m^2} \cdot \frac{40}{11} \cdot \frac{\epsilon^{\frac{1}{2}}}{\epsilon^{\frac{1}{4}(j+1)}} \left(2Q(\epsilon) \epsilon^{\frac{j-3}{8}} \right)^{\frac{3j+3}{4}} \leq \\
& \leq \frac{11}{40\pi S} \cdot \frac{1}{\sqrt{112} \cdot Q(\epsilon)} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{1}{4}(j+1)}} \left(1 + (2Q(\epsilon) \cdot \epsilon^{\frac{1}{4}})^1 + \dots + (2Q(\epsilon) \cdot \epsilon^{\frac{1}{4}})^{j - [\hat{j} + 1]} + \frac{20 \cdot \epsilon^{\frac{1}{4}}}{11 \cdot W} \left(2Q(\epsilon) \epsilon^{\frac{j-3}{8}} \right)^{\frac{3j+3}{4}} \right) \\
& \leq \frac{11}{40\pi} \cdot \frac{1}{\sqrt{112} \cdot Q(\epsilon)} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{1}{4}(j+1)}}
\end{aligned}$$

Notes

- by hypothesis, $0 < \bar{g} \leq \frac{2m^2}{W} \cdot \frac{1}{\sqrt{112} \cdot Q(\epsilon)} \cdot \frac{11}{40\pi S} \cdot \frac{1}{(24 \cdot 4^4)^2 \cdot \epsilon^{\frac{1}{4}}}$ and $\bar{g} \leq 1$, therefore we have that $\frac{\bar{g}^2}{4m^2} \cdot \frac{40\epsilon^{\frac{1}{2}}}{11} \leq \frac{20\epsilon^{\frac{1}{4}}}{11 \cdot W} \cdot \frac{1}{\sqrt{112} \cdot Q(\epsilon)} \cdot \frac{11}{40\pi S} \cdot \frac{1}{(24 \cdot 4^4)^2}$
- note that $\left(2Q(\epsilon) \epsilon^{\frac{j-3}{6j+6}}\right)^{\frac{3j+3}{4}} \leq \left(2Q(\epsilon) \epsilon^{\frac{1}{12}}\right)^6 \quad \forall j \geq 7$, as $0 < \epsilon < \left(\frac{1}{4}\right)^{16}$, $1 < Q(\epsilon) < 2 \Rightarrow 2Q(\epsilon) \cdot \epsilon^{\frac{1}{12}} < 1$
- by definition $S = \sum_{n=0}^{\infty} \left(2Q(\epsilon) \cdot \epsilon^{\frac{1}{4}}\right)^n + \frac{20\epsilon^{\frac{1}{4}}}{11 \cdot W} \cdot \left(2Q(\epsilon) \epsilon^{\frac{1}{12}}\right)^6$

2.2 Case $\mathbf{P} \neq 0$.

I repeat on the hamiltonian $H_{\mathbf{P},\sigma}$ the same operations made for the canonic form (10) of the transformed hamiltonian $H_{\mathbf{P}}$. Unlike the case without infrared cut-off, the transformation of $H_{\mathbf{P},\sigma}$ is implemented by the unitary operator

$$\mathcal{W}_{\sigma}(\mathbf{P}) = W_{\sigma}\left(\frac{\mathbf{P}_1}{m}\right) = e^{-g \int_{\sigma}^{\kappa} \frac{b(\mathbf{k}) - b^{\dagger}(\mathbf{k})}{|\mathbf{k}| \left(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m}\right)} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}}$$

where $\mathbf{P}_1 \equiv \mathbf{P} - \overline{\mathbf{P}^{mes}} = m \nabla E^{\sigma}(\mathbf{P})$. The last inequality follows from the perturbation of the isolated eigenvalue $E_{\mathbf{P}}^0$ of $H_{\mathbf{P}}^0|_{F_{\sigma}^+}$ (see [2]).

$\mathbf{P}_1$ verifies the equation:

$$\mathbf{P} - \mathbf{P}_1 = \frac{1}{\|\phi_{\mathbf{P}}^{\sigma}\|^2} (\phi_{\mathbf{P}}^{\sigma}, \mathbf{P}^{mes} \phi_{\mathbf{P}}^{\sigma}) - \frac{1}{\|\phi_{\mathbf{P}}^{\sigma}\|^2} \left(\phi_{\mathbf{P}}^{\sigma}, g \int_{\sigma}^{\kappa} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m}\right)} (b(\mathbf{k}) + b^{\dagger}(\mathbf{k})) d^3 k \phi_{\mathbf{P}}^{\sigma} \right) + g^2 \int_{\sigma}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m}\right)^2} d^3 k$$

as $\phi_{\mathbf{P}}^{\sigma}$ is the ground state of the transformed hamiltonian $W_{\sigma}\left(\frac{\mathbf{P}_1}{m}\right) H_{\mathbf{P},\sigma} W_{\sigma}^{\dagger}\left(\frac{\mathbf{P}_1}{m}\right)$

Transformed hamiltonian.

I will write $H_{\mathbf{P},\sigma}$, $\mathbf{P} = \mathbf{P}_1 + \mathbf{P}_2$, as

$$\begin{aligned}
H_{\mathbf{P},\sigma} &= \frac{(\mathbf{P}_1 + \mathbf{P}_2 - \mathbf{P}^{mes})^2}{2m} + g \int_{\sigma}^{\kappa} \left(b(\mathbf{k}) + b^{\dagger}(\mathbf{k}) \right) \frac{d^3 \mathbf{k}}{\sqrt{2|\mathbf{k}|}} + H^{mes} = \\
&= \frac{(\mathbf{P}_1 + \mathbf{P}_2)^2}{2m} - \frac{(\mathbf{P}_1 + \mathbf{P}_2) \cdot \mathbf{P}^{mes}}{m} + \frac{\mathbf{P}^{mes^2}}{2m} + g \int_{\sigma}^{\kappa} \left(b(\mathbf{k}) + b^{\dagger}(\mathbf{k}) \right) \frac{d^3 \mathbf{k}}{\sqrt{2|\mathbf{k}|}} + H^{mes} = \\
&= \frac{(\mathbf{P}_1 + \mathbf{P}_2)^2}{2m} + \frac{\mathbf{P}^{mes^2}}{2m} - \frac{\mathbf{P}_2 \cdot \mathbf{P}^{mes}}{m} + \int_{\kappa}^{\infty} \left(|\mathbf{k}| - \mathbf{k} \cdot \frac{\mathbf{P}_1}{m} \right) b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3 k + \\
&+ \int_{\sigma}^{\kappa} \left(|\mathbf{k}| - \mathbf{k} \cdot \frac{\mathbf{P}_1}{m} \right) \left(b^{\dagger}(\mathbf{k}) + \frac{g}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} \right) \left(b(\mathbf{k}) + \frac{g}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} \right) d^3 k \\
&+ \int_0^{\sigma} \left(|\mathbf{k}| - \mathbf{k} \cdot \frac{\mathbf{P}_1}{m} \right) b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3 k - g^2 \int_{\sigma}^{\kappa} \frac{1}{2|\mathbf{k}|^2(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} d^3 k
\end{aligned}$$

and I will perform the coherent transformation:

$$\begin{aligned}
W_{\sigma} \left(\frac{\mathbf{P}_1}{m} \right) H_{\mathbf{P},\sigma} W_{\sigma}^{\dagger} \left(\frac{\mathbf{P}_1}{m} \right) &= \\
&= \frac{(\mathbf{P}_1 + \mathbf{P}_2)^2}{2m} + \frac{1}{2m} \left(\mathbf{P}^{mes} - g \int_{\sigma}^{\kappa} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} \left(b(\mathbf{k}) + b^{\dagger}(\mathbf{k}) \right) d^3 k + g^2 \int_{\sigma}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3 k \right)^2 + \\
&- \frac{\mathbf{P}_2}{m} \left(\mathbf{P}^{mes} - g \int_{\sigma}^{\kappa} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} \left(b(\mathbf{k}) + b^{\dagger}(\mathbf{k}) \right) d^3 k + g^2 \int_{\sigma}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3 k \right) + \\
&+ \int_0^{\infty} \left(|\mathbf{k}| - \mathbf{k} \cdot \frac{\mathbf{P}_1}{m} \right) b^{\dagger}(\mathbf{k}) b(\mathbf{k}) d^3 k - g^2 \int_{\sigma}^{\kappa} \frac{1}{2|\mathbf{k}|^2(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} d^3 k
\end{aligned}$$

Considerations

Having defined $\Pi_{\mathbf{P},\sigma} = \mathbf{P}^{mes} - g \int_{\sigma}^{\kappa} \frac{\mathbf{k}(b(\mathbf{k}) + b^{\dagger}(\mathbf{k}))}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} d^3 k$ it follows that

$$\mathbf{P}_2 = \frac{1}{\|\phi_{\mathbf{P}}^{\sigma}\|^2} (\phi_{\mathbf{P}}^{\sigma}, \Pi_{\mathbf{P},\sigma} \phi_{\mathbf{P}}^{\sigma}) + g^2 \int_{\sigma}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3 k$$

By substitution we obtain

$$\begin{aligned}
H_{\mathbf{P},\sigma}^w &\equiv W_{\sigma} \left(\frac{\mathbf{P}_1}{m} \right) H_{\mathbf{P},\sigma} W_{\sigma}^{\dagger} \left(\frac{\mathbf{P}_1}{m} \right) = \\
&= \frac{\mathbf{P}_2^2}{2m} + \frac{1}{2m} \left(\Pi_{\mathbf{P},\sigma} + g^2 \int_{\sigma}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3 k \right)^2 - \frac{\mathbf{P}_2}{m} \left(\Pi_{\mathbf{P},\sigma} + g^2 \int_{\sigma}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3(1 - \widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3 k \right) +
\end{aligned}$$

$$\begin{aligned}
& + \int_0^\infty \left(|\mathbf{k}| - \mathbf{k} \cdot \frac{\mathbf{P}_1}{m} \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k - g^2 \int_\sigma^\kappa \frac{1}{2|\mathbf{k}|^2(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} d^3k = \\
& = \frac{1}{2m} (\Pi_{\mathbf{P},\sigma})^2 + \frac{1}{2m} \left(g^2 \int_\sigma^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right)^2 + \frac{1}{m} \Pi_{\mathbf{P},\sigma} \cdot \left(g^2 \int_\sigma^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right) + \\
& - \frac{1}{m} \left(\frac{1}{\|\phi_{\mathbf{P}}^\sigma\|^2} \cdot (\phi_{\mathbf{P}}^\sigma, \Pi_{\mathbf{P},\sigma} \phi_{\mathbf{P}}^\sigma) + g^2 \int_\sigma^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right) \cdot \left(\Pi_{\mathbf{P},\sigma} + g^2 \int_\sigma^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right) + \\
& + \int_0^\infty \left(|\mathbf{k}| - \mathbf{k} \cdot \frac{\mathbf{P}_1}{m} \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k - g^2 \int_\sigma^\kappa \frac{1}{2|\mathbf{k}|^2(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} d^3k + \frac{\mathbf{P}^2}{2m} = \\
& = \frac{1}{2m} \left(\Pi_{\mathbf{P},\sigma} - \frac{1}{\|\phi_{\mathbf{P}}^\sigma\|^2} \cdot (\phi_{\mathbf{P}}^\sigma, \Pi_{\mathbf{P},\sigma} \phi_{\mathbf{P}}^\sigma) \right)^2 + \int_0^\infty (|\mathbf{k}| - \mathbf{k} \cdot \nabla E^\sigma(\mathbf{P})) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k + \\
& + \frac{1}{2m} \left(g^2 \int_\sigma^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right)^2 - \frac{1}{m} \left(g^2 \int_\sigma^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right) \cdot \left(g^2 \int_\sigma^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right) + \\
& - g^2 \int_\sigma^\kappa \frac{1}{2|\mathbf{k}|^2(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} d^3k + \frac{\mathbf{P}^2}{2m} - \frac{1}{2m} \left(\frac{1}{\|\phi_{\mathbf{P}}^\sigma\|^2} \cdot (\phi_{\mathbf{P}}^\sigma, \Pi_{\mathbf{P},\sigma} \phi_{\mathbf{P}}^\sigma) \right)^2 + \\
& - \frac{1}{m} \left(\frac{1}{\|\phi_{\mathbf{P}}^\sigma\|^2} \cdot (\phi_{\mathbf{P}}^\sigma, \Pi_{\mathbf{P},\sigma} \phi_{\mathbf{P}}^\sigma) \right) \cdot \left(g^2 \int_\sigma^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right) = \\
& = \frac{1}{2m} \left(\Pi_{\mathbf{P},\sigma} - \frac{1}{\|\phi_{\mathbf{P}}^\sigma\|^2} \cdot (\phi_{\mathbf{P}}^\sigma, \Pi_{\mathbf{P},\sigma} \phi_{\mathbf{P}}^\sigma) \right)^2 + \int_0^\infty (|\mathbf{k}| - \mathbf{k} \cdot \nabla E^\sigma(\mathbf{P})) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k + c_{\mathbf{P}}(\sigma)
\end{aligned}$$

where

$$\begin{aligned}
c_{\mathbf{P}}(\sigma) & = -\frac{1}{2m} \left(g^2 \int_\sigma^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right)^2 - \frac{1}{2m} \left(\frac{1}{\|\phi_{\mathbf{P}}^\sigma\|^2} \cdot (\phi_{\mathbf{P}}^\sigma, \Pi_{\mathbf{P},\sigma} \phi_{\mathbf{P}}^\sigma) \right)^2 + \\
& - \frac{1}{m} \left(\frac{1}{\|\phi_{\mathbf{P}}^\sigma\|^2} \cdot (\phi_{\mathbf{P}}^\sigma, \Pi_{\mathbf{P},\sigma} \phi_{\mathbf{P}}^\sigma) \right) \cdot \left(g^2 \int_\sigma^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right) - g^2 \int_\sigma^\kappa \frac{1}{2|\mathbf{k}|^2(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} d^3k + \frac{\mathbf{P}^2}{2m} = \\
& = -\frac{1}{2m} \left[\left(g^2 \int_\sigma^\kappa \frac{\mathbf{k}}{2|\mathbf{k}|^3(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})^2} d^3k \right) + \left(\frac{1}{\|\phi_{\mathbf{P}}^\sigma\|^2} \cdot (\phi_{\mathbf{P}}^\sigma, \Pi_{\mathbf{P},\sigma} \phi_{\mathbf{P}}^\sigma) \right) \right]^2 - g^2 \int_\sigma^\kappa \frac{1}{2|\mathbf{k}|^2(1-\widehat{\mathbf{k}} \cdot \frac{\mathbf{P}_1}{m})} d^3k + \\
& + \frac{\mathbf{P}^2}{2m} = -\frac{1}{2m} [\mathbf{P} - m \nabla E^\sigma(\mathbf{P})]^2 - g^2 \int_\sigma^\kappa \frac{1}{2|\mathbf{k}|^2(1-\widehat{\mathbf{k}} \cdot \nabla E^\sigma)} d^3k + \frac{\mathbf{P}^2}{2m}
\end{aligned}$$

From lemma A2 (point 1), $|c_{\mathbf{P}}(\sigma)|$ will be bounded uniformly in σ and $|\nabla E^\sigma(\mathbf{P})| < 1$.

Now, given the infrared cut-off $\epsilon^{\frac{j+1}{2}}$ and $\epsilon^{\frac{j+2}{2}}$, I perform the unitary transformations

$$i) \ H_{\mathbf{P}, \epsilon^{\frac{j+1}{2}}} \rightarrow W_{\epsilon^{\frac{j+1}{2}}} \left(\nabla E^{\epsilon^{\frac{j+1}{2}}}(\mathbf{P}) \right) H_{\mathbf{P}, \epsilon^{\frac{j+1}{2}}} W_{\epsilon^{\frac{j+1}{2}}}^\dagger \left(\nabla E^{\epsilon^{\frac{j+1}{2}}}(\mathbf{P}) \right)$$

$$\text{ii) } H_{\mathbf{P},\epsilon}^{j+2} \rightarrow W_{\epsilon}^{j+2} \left(\nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) H_{\mathbf{P},\epsilon}^{j+2} W_{\epsilon}^{\dagger j+2} \left(\nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right)$$

Note that the two transformations are different in the cut-off but not in the coherent factor.

Note

In order to compress the formulas, from now on I will use the following

$$\Gamma_{\mathbf{P},\epsilon}^{j+1} \equiv \Pi_{\mathbf{P},\epsilon}^{j+1} - \frac{\left(\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \Pi_{\mathbf{P},\epsilon}^{j+1} \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)}{\left\| \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|^2}$$

We define:

$$\begin{aligned} H_{\mathbf{P},\epsilon}^{j+1} &\equiv W_{\epsilon}^{j+1} \left(\nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) H_{\mathbf{P},\epsilon}^{j+1} W_{\epsilon}^{\dagger j+1} \left(\nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) = \\ &= \frac{1}{2m} \left(\Gamma_{\mathbf{P},\epsilon}^{j+1} \right)^2 + \int_0^\infty \left(|\mathbf{k}| - \mathbf{k} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k + c_{\mathbf{P}}(j+1) \end{aligned}$$

$$\begin{aligned} \widehat{H}_{\mathbf{P},\epsilon}^{j+2} &\equiv W_{\epsilon}^{j+2} \left(\nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) H_{\mathbf{P},\epsilon}^{j+2} W_{\epsilon}^{\dagger j+2} \left(\nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) = \\ &= \frac{1}{2m} \left(\Gamma_{\mathbf{P},\epsilon}^{j+1} - g \int_{\epsilon}^{\frac{j+1}{2}} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right)} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3k + g^2 \int_{\epsilon}^{\frac{j+1}{2}} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right)^2} d^3k \right)^2 + \\ &+ \int_0^\infty \left(|\mathbf{k}| - \mathbf{k} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k + \widehat{c}_{\mathbf{P}}(j+2) \end{aligned}$$

where

$$\begin{aligned} \widehat{c}_{\mathbf{P}}(j+2) &= -\frac{1}{2m} \left(\frac{1}{\left\| \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|^2} \left(\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \Pi_{\mathbf{P},\epsilon}^{j+1} \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) + g^2 \int_{\epsilon}^{\frac{j+1}{2}} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right)^2} d^3k \right)^2 + \\ &- g^2 \int_{\epsilon}^{\frac{j+1}{2}} \frac{1}{2|\mathbf{k}|^2 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right)} d^3k + \frac{\mathbf{P}^2}{2m}. \end{aligned}$$

By construction, the following equality holds

$$\begin{aligned} H_{\mathbf{P},\epsilon}^w \frac{j+2}{2} &= W_{\epsilon} \frac{j+2}{2} \left(\nabla E^{\epsilon \frac{j+2}{2}}(\mathbf{P}) \right) H_{\mathbf{P},\epsilon} \frac{j+2}{2} W_{\epsilon}^{\dagger} \frac{j+2}{2} \left(\nabla E^{\epsilon \frac{j+2}{2}}(\mathbf{P}) \right) = \\ &= W_{\epsilon} \frac{j+2}{2} \left(\nabla E^{\epsilon \frac{j+2}{2}}(\mathbf{P}) \right) W_{\epsilon}^{\dagger} \frac{j+2}{2} \left(\nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) \widehat{H}_{\mathbf{P},\epsilon}^w \frac{j+2}{2} W_{\epsilon} \frac{j+2}{2} \left(\nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) W_{\epsilon}^{\dagger} \frac{j+2}{2} \left(\nabla E^{\epsilon \frac{j+2}{2}}(\mathbf{P}) \right) \end{aligned}$$

(the hamiltonians $H_{\mathbf{P},\epsilon} \frac{j+1}{2}, H_{\mathbf{P},\epsilon}^w \frac{j+1}{2}$ and $\widehat{H}_{\mathbf{P},\epsilon}^w \frac{j+1}{2}$ are s.a. on the same domain and the formal equalities are well defined from an operatorial point of view).

Definitions

$$\begin{aligned} \widehat{\Pi}_{\mathbf{P},\epsilon}^i \frac{j+1}{2} &\equiv W_{\epsilon} \frac{j+1}{2} \left(\nabla E^{\epsilon \frac{j}{2}}(\mathbf{P}) \right) W_{\epsilon}^{\dagger} \frac{j+1}{2} \left(\nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) \Pi_{\mathbf{P},\epsilon}^i \frac{j+1}{2} W_{\epsilon} \frac{j+1}{2} \left(\nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) W_{\epsilon}^{\dagger} \frac{j+1}{2} \left(\nabla E^{\epsilon \frac{j}{2}}(\mathbf{P}) \right) = \\ &= \mathbf{P}^{mes^i} - g \int_{\epsilon}^{\kappa} \frac{k^i (b(\mathbf{k}) + b^{\dagger}(\mathbf{k}))}{\sqrt{2} |\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j}{2}} \right)} d^3 k + \frac{g^2}{2} \int_{\epsilon}^{\kappa} \frac{k^i}{|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j}{2}} \right)^2} d^3 k - \frac{g^2}{2} \int_{\epsilon}^{\kappa} \frac{k^i}{|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)^2} d^3 k; \\ \widehat{\Gamma}_{\mathbf{P},\epsilon}^i \frac{j+1}{2} &\equiv \widehat{\Pi}_{\mathbf{P},\epsilon}^i \frac{j+1}{2} - \frac{\left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \widehat{\Pi}_{\mathbf{P},\epsilon}^i \frac{j+1}{2} \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)}{\left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|^2} \end{aligned}$$

Theorem 2.3bis

Convergence of the sequence of the ground states $H_{\mathbf{P},\epsilon}^w \frac{j+1}{2}$.

In order to arrive to a convergent sequence of ground eigenvectors, one starts from the cut-off ϵ and from the (normalized) vector $W_{\epsilon}(\nabla E^{\epsilon}(\mathbf{P})) \psi_{\mathbf{P}}^{\epsilon}$ and proceeds with the same iteration, on the basis of the results of lemma A1. Comparing the hamiltonians $H_{\mathbf{P},\epsilon}^w \frac{j+1}{2}$ and $\widehat{H}_{\mathbf{P},\epsilon}^w \frac{j+2}{2}$ we can build $\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}$ in terms of $\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}$ and finally we define $\phi_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \equiv W_{\epsilon} \frac{j+2}{2} \left(\nabla E^{\epsilon \frac{j+2}{2}}(\mathbf{P}) \right) W_{\epsilon}^{\dagger} \frac{j+2}{2} \left(\nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}$.

Like in the case $\mathbf{P} = 0$, in order to prove the convergence of the vectors $\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}$, one needs a more refined estimate of the contribution, to the difference between the generic vectors $\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}$ and $\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}$, given by the mixed terms containing the $\Gamma_{\mathbf{P},\epsilon}^i \frac{j+1}{2}$. For this purpose the procedure to follow is slightly more elaborated than in the case

where $\mathbf{P} = 0$, because it is necessary to compare the following vectors one after the other:

$$\phi_{\mathbf{P}}^{\epsilon \frac{j}{2}} \rightarrow \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \rightarrow \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \rightarrow \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \rightarrow \phi_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}.$$

I apply a proper adaptation of the lemmas 2.1 e 2.2. to estimate the contribution to the norm of $\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}$ given by the examined mixed terms. Apart from numeric differences, even in this case the improvement of the estimate consists in the bound of an expression like (14) of theorem 2.3 with a quantity of the order $\epsilon^{-\frac{j+1}{4}}$ (that means multiplicative constants that are uniform in j):

$$\frac{g^2}{4m^2} \left| \left(\Gamma_{\mathbf{P}, \epsilon \frac{j+1}{2}}^i \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \left(\frac{1}{H_{\mathbf{P}, \epsilon \frac{j+1}{2}}^w - E(j+1)} \right) \Gamma_{\mathbf{P}, \epsilon \frac{j+1}{2}}^i \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) \right|$$

where $E(j+1)$ is s.t. $\left| E(j+1) - E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right| = \frac{11}{20} \epsilon^{\frac{j+2}{2}}.$

Note

In the adaptation to the examined case of the derivation of the (13) from the (14), take care of lemma A1 and of the following difference:

$$\widehat{c}_{\mathbf{P}}(j+2) - c_{\mathbf{P}}(j+1) = -g^2 \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{2|\mathbf{k}|^2 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right)} d^3 k$$

The bound for the module of the above scalar product is the result of an inductive procedure like the one where $\mathbf{P} = 0$; it is for this reason that I will not repeat the entire proof but I will point out the key ingredients without further details.

I apply to both the factors of the scalar product the unitary operator

$$W_{\epsilon \frac{j+1}{2}} \left(\nabla E^{\epsilon \frac{j}{2}}(\mathbf{P}) \right) W_{\epsilon \frac{j+1}{2}}^{\dagger} \left(\nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right)$$

(such an operation is not required if $\mathbf{P} = 0$) and I obtain

$$\left(\widehat{\Gamma}_{\mathbf{P}, \epsilon}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \left(\frac{1}{\widehat{H}_{\mathbf{P}, \epsilon}^w \frac{j+1}{2} - E(j+1)} \right) \widehat{\Gamma}_{\mathbf{P}, \epsilon}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)$$

as in theorem 2.3 I go on with the following bound (from above)

$$\begin{aligned} & \left| \left(\widehat{\Gamma}_{\mathbf{P}, \epsilon}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \left(\frac{1}{\widehat{H}_{\mathbf{P}, \epsilon}^w \frac{j+1}{2} - E(j+1)} \right) \widehat{\Gamma}_{\mathbf{P}, \epsilon}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) \right| \leq \\ & \leq 2 \left| \left(\widehat{\Gamma}_{\mathbf{P}, \epsilon}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \left| \left(\frac{1}{H_{\mathbf{P}, \epsilon}^w \frac{j}{2} - E(j)} \right) \widehat{\Gamma}_{\mathbf{P}, \epsilon}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right| \right) \right| \leq \\ & \leq 2 \left| \left(\widehat{\Gamma}_{\mathbf{P}, \epsilon}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - \Gamma_{\mathbf{P}, \epsilon}^i \phi_{\mathbf{P}}^{\epsilon \frac{j}{2}}, \left(\frac{1}{H_{\mathbf{P}, \epsilon}^w \frac{j}{2} - E(j)} \right) \left(\widehat{\Gamma}_{\mathbf{P}, \epsilon}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - \Gamma_{\mathbf{P}, \epsilon}^i \phi_{\mathbf{P}}^{\epsilon \frac{j}{2}} \right) \right) \right| + \\ & + 4 \left| \left(\widehat{\Gamma}_{\mathbf{P}, \epsilon}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - \Gamma_{\mathbf{P}, \epsilon}^i \phi_{\mathbf{P}}^{\epsilon \frac{j}{2}}, \left| \left(\frac{1}{H_{\mathbf{P}, \epsilon}^w \frac{j}{2} - E(j)} \right) \left(\Gamma_{\mathbf{P}, \epsilon}^i \phi_{\mathbf{P}}^{\epsilon \frac{j}{2}} \right) \right| \right) \right| + \\ & + 2 \left| \left(\Gamma_{\mathbf{P}, \epsilon}^i \phi_{\mathbf{P}}^{\epsilon \frac{j}{2}}, \left| \left(\frac{1}{H_{\mathbf{P}, \epsilon}^w \frac{j}{2} - E(j)} \right) \Gamma_{\mathbf{P}, \epsilon}^i \phi_{\mathbf{P}}^{\epsilon \frac{j}{2}} \right| \right) \right| \end{aligned}$$

The proof goes on as in theorem 2.3 taking into account the following information:

1)

$$\begin{aligned} \widehat{\Pi}_{\mathbf{P}, \epsilon}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - \Pi_{\mathbf{P}, \epsilon}^i \phi_{\mathbf{P}}^{\epsilon \frac{j}{2}} &= -g \int_{\epsilon \frac{j+1}{2}}^{\frac{j}{2}} \frac{k^i (b(\mathbf{k}) + b^t(\mathbf{k}))}{\sqrt{2} |\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j}{2}} \right)} d^3 k + \frac{g^2}{2} \int_{\epsilon \frac{j+1}{2}}^{\kappa} \frac{k^i}{|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j}{2}} \right)^2} d^3 k + \\ & - \frac{g^2}{2} \int_{\epsilon \frac{j+1}{2}}^{\kappa} \frac{k^i}{|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)^2} d^3 k \end{aligned}$$

$$\begin{aligned} & \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \widehat{\Pi}_{\mathbf{P}, \epsilon}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) - \left(\phi_{\mathbf{P}}^{\epsilon \frac{j}{2}}, \Pi_{\mathbf{P}, \epsilon}^i \phi_{\mathbf{P}}^{\epsilon \frac{j}{2}} \right) = \\ & = 2Re \left(\Pi_{\mathbf{P}, \epsilon}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j}{2}}, \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - \phi_{\mathbf{P}}^{\epsilon \frac{j}{2}} \right) + \left(\Pi_{\mathbf{P}, \epsilon}^i \left(\phi_{\mathbf{P}}^{\epsilon \frac{j}{2}} - \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right), \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - \phi_{\mathbf{P}}^{\epsilon \frac{j}{2}} \right) + \\ & + \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \left\{ \widehat{\Pi}_{\mathbf{P}, \epsilon}^i - \Pi_{\mathbf{P}, \epsilon}^i \right\} \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) \end{aligned}$$

$$\begin{aligned}
2) \quad & \left\| \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} q^i b(\mathbf{q}) \frac{d^3 q}{|\mathbf{q}| \sqrt{2|\mathbf{q}|} \left(1 - \widehat{\mathbf{q}} \cdot \nabla E_{\mathbf{P}, \epsilon^{\frac{j}{2}}}^{\frac{j}{2}}\right)} \left(\frac{1}{H_{\mathbf{P}, \epsilon^{\frac{j}{2}}}^w - E(j)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+1}{2}}}^+} \text{ is of order } \epsilon^{\frac{j}{4}} \\
2bis) \quad & \left\| \int_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j}{2}}} q^i b^\dagger(\mathbf{q}) \frac{d^3 q}{|\mathbf{q}| \sqrt{2|\mathbf{q}|} \left(1 - \widehat{\mathbf{q}} \cdot \nabla E_{\mathbf{P}, \epsilon^{\frac{j}{2}}}^{\frac{j}{2}}\right)} \left(\frac{1}{H_{\mathbf{P}, \epsilon^{\frac{j}{2}}}^w - E(j)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+1}{2}}}^+} \text{ is of order } \epsilon^{\frac{j}{4}}
\end{aligned}$$

3) inductive hypothesis for evaluating:

$$\begin{aligned}
i) \quad & \left\| \frac{g^2}{4m^2} \left\| \frac{1}{H_{\mathbf{P}, \epsilon^{\frac{j}{2}}}^w - E(j)} \right\|^{\frac{1}{2}} d^3 k \Gamma_{\epsilon^{\frac{j}{2}}}^i \phi_{\epsilon^{\frac{j}{2}}}^{\frac{j}{2}} \right\|^2 \text{ of order } \epsilon^{-\frac{j}{4}} \\
ii) \quad & \left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} - \phi_{\mathbf{P}}^{\epsilon^{\frac{j}{2}}} \right\| \text{ of order } \epsilon^{\frac{j}{8}} = \sigma_{j-1}^{\frac{1}{4}}, \text{ which implies} \\
& \bullet \left\| \Pi_{\mathbf{P}, \epsilon^{\frac{j}{2}}}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} - \Pi_{\mathbf{P}, \epsilon^{\frac{j}{2}}}^i \phi_{\mathbf{P}}^{\epsilon^{\frac{j}{2}}} \right\| < \sqrt{2m} \left(E_{\mathbf{P}}^{\frac{j}{2}} - c_{\mathbf{P}}(j) \right)^{\frac{1}{2}} \left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} - \phi_{\mathbf{P}}^{\epsilon^{\frac{j}{2}}} \right\| \text{ then of order } \\
& \epsilon^{\frac{j}{8}} \text{ (the uniformity in } j \text{ of the bound of } \left(E_{\mathbf{P}}^{\frac{j}{2}} - c_{\mathbf{P}}(j) \right) \text{ follows from lemma A2)} \\
& \bullet \left| \nabla E_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} - \nabla E_{\mathbf{P}}^{\epsilon^{\frac{j}{2}}} \right| \leq C(g) \epsilon^{\frac{j}{8}} \text{ (see lemma A2)}
\end{aligned}$$

As for the estimate of

$$\left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} - \phi_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} \right\| = \left\| W_{\epsilon^{\frac{j+2}{2}}} \left(\nabla E_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}(\mathbf{P}) \right) W_{\epsilon^{\frac{j+2}{2}}}^\dagger \left(\nabla E_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}}(\mathbf{P}) \right) \phi_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} - \phi_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} \right\|$$

note that the difference between $\nabla E_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}$ and $\nabla E_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}}$ is bounded by a quantity of order $\epsilon^{\frac{j+1}{8}}$. This allows to neutralize the logarithmic divergence in $\epsilon^{\frac{j+2}{2}}$ which arises from the Weyl operators $W_{\epsilon^{\frac{j+2}{2}}} \left(\nabla E_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}(\mathbf{P}) \right) W_{\epsilon^{\frac{j+2}{2}}}^\dagger \left(\nabla E_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}}(\mathbf{P}) \right)$.

In conclusion, **tuning $\mathbf{g}$ uniformly in $\mathbf{j}$** as we saw in theorem 2.3, it is possible to bound the norm difference between $\phi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}$ and $\widehat{\phi}_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}}$ with $\epsilon^{\frac{j+1}{8}}$ and the difference between $\phi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}$ and $\phi_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}}$ with $\epsilon^{\frac{j+1}{16}}$.

Observation

The logarithmic divergence in $\epsilon^{\frac{j+2}{2}}$ related to the Weyl operators includes the one that follows from the first approximation of:

$$\left(\int_{\epsilon^{\frac{j+2}{2}}}^{\kappa} \left\| b(\mathbf{k}) W_{\epsilon^{\frac{j+2}{2}}}^b \left(\nabla E^{\epsilon^{\frac{j+2}{2}}}(\mathbf{P}) \right) \psi_{\mathbf{P}, \epsilon^{\frac{j+2}{2}}} \right\|^2 d^3k \right)^{\frac{1}{2}}$$

being $b(\mathbf{k}) \psi_{\mathbf{P}, \epsilon^{\frac{j+2}{2}}} = \frac{g}{\sqrt{2|\mathbf{k}|}} \left(\frac{1}{E^{\epsilon^{\frac{j+2}{2}}}(\mathbf{P}) - |\mathbf{k}| - H_{\mathbf{P}-\mathbf{k}, \epsilon^{\frac{j+2}{2}}}} \right) \psi_{\mathbf{P}, \epsilon^{\frac{j+2}{2}}}$ (see [2]) and $\left\| \psi_{\mathbf{P}, \epsilon^{\frac{j+2}{2}}} \right\| < 1$.

(see also lemma B2 in Appendix B).

Corollary 2.4

The sequence $\left\{ \phi_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} \right\}$ ($\phi_{\mathbf{P}}^{\epsilon}$ normalized vector) converges to a non-vanishing vector when the value of the coupling constant is less or equal to the $\bar{g}$ determined by theorems 2.3 and 2.3bis, therefore such as that $\left\| \phi_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} - \phi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} \right\| \leq \epsilon^{\frac{j+1}{16}}$.

Proof

This is a Cauchy sequence because $\forall l, j \quad l \geq j$

$$\begin{aligned} \left\| \phi_{\mathbf{P}}^{\epsilon^{\frac{l+1}{2}}} - \phi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} \right\| &\leq \left\| \phi_{\mathbf{P}}^{\epsilon^{\frac{l+1}{2}}} - \phi_{\mathbf{P}}^{\epsilon^{\frac{l}{2}}} + \phi_{\mathbf{P}}^{\epsilon^{\frac{l}{2}}} - \dots - \phi_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} + \phi_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} - \phi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} \right\| \leq \\ &\leq \epsilon^{\frac{l}{16}} + \epsilon^{\frac{l-1}{16}} + \dots + \epsilon^{\frac{j+1}{16}} \leq \epsilon^{\frac{j+1}{16}} \cdot \left(\frac{1}{1 - \epsilon^{\frac{1}{16}}} \right) \end{aligned}$$

The limit does not vanish since the following inequality holds, uniformly in j :

$$\left\| \phi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} \right\| \geq \left\| \phi_{\mathbf{P}}^{\epsilon} \right\| - \left(\epsilon^{\frac{1}{16}} + \epsilon^{\frac{2}{16}} + \dots + \epsilon^{\frac{j}{16}} \right) \geq 1 - \left(\frac{\epsilon^{\frac{1}{16}}}{1 - \epsilon^{\frac{1}{16}}} \right) \geq \frac{1 - 2\epsilon^{\frac{1}{16}}}{1 - \epsilon^{\frac{1}{16}}} > \frac{2}{3}.$$

3 Spectral regularity.

In this paragraph I will define a normalized vector $\phi_{\mathbf{P}}^\sigma$, that is the ground state of $H_{\mathbf{P},\sigma}^w |_{F_\sigma^+}$ ($\sigma \leq \epsilon$). It has a regularity property in $\mathbf{P}$ required for the construction of the scattering states in the next chapter (in particular I will use the vectors $W_\sigma^\dagger(\nabla E^\sigma(\mathbf{P})) \phi_{\mathbf{P}}^\sigma$).

We arrive at the vector $\phi_{\mathbf{P}}^\sigma$ through an intermediate (not normalized) vector $\tilde{\phi}_{\mathbf{P}}^\sigma$ from which it differs only for a phase term, apart from the normalization. The choice of the right phase is aimed at two results:

- the norm convergence of the vector $\phi_{\mathbf{P}}^\sigma$, for $\sigma \rightarrow 0$, to a vector $\phi_{\mathbf{P}}$;
- the acquisition of the following Hoelder property, with respect to $\mathbf{P}$:

$$\|\phi_{\mathbf{P}+\Delta\mathbf{P}}^\sigma - \phi_{\mathbf{P}}^\sigma\| \leq C |\Delta\mathbf{P}|^{\frac{1}{32}}$$

where C is a uniform constant in $\mathbf{P}, \mathbf{P} + \Delta\mathbf{P} \in \Sigma$ ($|\Delta\mathbf{P}| < \left(\frac{1}{6}\right)^4$) and in $0 < \sigma \leq \epsilon$

Definition of $\tilde{\phi}_{\mathbf{P}}^\sigma$

Initial conditions

We start from the initial infrared cut-off $\{\epsilon' : \epsilon \geq \epsilon' \geq \epsilon\sqrt{\epsilon}\}$ and from a coupling constant $\bar{g}$ such that, uniformly in ϵ' and in $\mathbf{P} \in \Sigma$, it is possible to perform the iterative procedure with the properties already shown when the starting cut-off is ϵ , in particular the validity of the theorem 2.3bis. It is also required that for the chosen value $\bar{g}$ we have $\forall \epsilon'$ and $\forall \mathbf{P} \in \Sigma$ $\left| \left(\phi_{\mathbf{P}}^{\epsilon'}, \psi_0 \right) \right| > 1 - r(\bar{g})$ where $r(\bar{g}) < \frac{2}{3}$.

Procedure

I perform the iteration shown in the previous chapter for each ϵ' . Given a σ ranging between $\epsilon^{\frac{j+2}{2}}$ and $\epsilon^{\frac{j+3}{2}}$ we can always write it as $\epsilon'^{\frac{j+2}{2}}$ where $\epsilon' = \epsilon'(\sigma) = \sigma^{\frac{2}{j+2}}$. I define

$$\tilde{\phi}_{\mathbf{P}}^\sigma \equiv \phi_{\mathbf{P}}^{\epsilon'(\sigma)^{\frac{j+2}{2}}}$$

Definition of $\phi_{\mathbf{P}}^\sigma$.

To go on with the definition I use the thesis of lemma 3.1 that will be later proved:

$$\left(\tilde{\phi}_{\mathbf{P}}^\sigma, \psi_0\right) \neq 0 \quad \forall \sigma < \epsilon, \forall \mathbf{P} \in \Sigma.$$

Since $\tilde{\phi}_{\mathbf{P}}^\sigma$ is ground state of $H_{\mathbf{P},\sigma}^w|_{F_\sigma^+}$ with a gap bigger than $\frac{\sigma}{2}$ and because of previous result, I realize that the normalized vector

$$\phi_{\mathbf{P}}^\sigma \equiv \frac{-\frac{1}{2\pi i} \oint \frac{1}{H_{\mathbf{P},\sigma}^w - E} dE \psi_0}{\left\| -\frac{1}{2\pi i} \oint \frac{1}{H_{\mathbf{P},\sigma}^w - E} dE \psi_0 \right\|}$$

(where $E \in \mathcal{C}$ and s.t. $|E - E_{\mathbf{P}}^\sigma| = \frac{\sigma}{4}$)
is the ground state of $H_{\mathbf{P},\sigma}^w|_{F_\sigma^+}$.

Lemma 3.1

$$\left(\tilde{\phi}_{\mathbf{P}}^\sigma, \psi_0\right) \neq 0 \quad \forall \sigma \leq \epsilon, \forall \mathbf{P} \in \Sigma$$

Proof

Knowing that:

- $\left\| \phi_{\mathbf{P}}^{\epsilon'} - \phi_{\mathbf{P}}^{\epsilon' \frac{i+2}{2}} \right\| \leq \frac{\epsilon' \frac{1}{16}}{1 - \epsilon' \frac{1}{16}} \leq \frac{1}{3} \quad (\text{corollary 2.4})$
- $\left(\tilde{\phi}_{\mathbf{P}}^\sigma, \psi_0\right) = \left(\tilde{\phi}_{\mathbf{P}}^\sigma - \phi_{\mathbf{P}}^{\epsilon'(\sigma)} + \phi_{\mathbf{P}}^{\epsilon'(\sigma)}, \psi_0\right) = \left(\tilde{\phi}_{\mathbf{P}}^\sigma - \phi_{\mathbf{P}}^{\epsilon'(\sigma)}, \psi_0\right) + \left(\phi_{\mathbf{P}}^{\epsilon'(\sigma)}, \psi_0\right)$
- for the hypothesis in the definition of $\tilde{\phi}_{\mathbf{P}}^\sigma$, $\left| \left(\phi_{\mathbf{P}}^{\epsilon'}, \psi_0\right) \right| > 1 - r(\overline{g})$ where $r(\overline{g}) < \frac{2}{3}$

we have that

$$\left| \left(\tilde{\phi}_{\mathbf{P}}^\sigma, \psi_0\right) \right| \geq \left| \left(\phi_{\mathbf{P}}^{\epsilon'(\sigma)}, \psi_0\right) \right| - \left| \left(\tilde{\phi}_{\mathbf{P}}^\sigma - \phi_{\mathbf{P}}^{\epsilon'(\sigma)}, \psi_0\right) \right| > 1 - r(\overline{g}) - \frac{1}{3} > 0$$

Theorem 3.2

For $\mathbf{P} \in \Sigma$, the limits $s - \lim_{\sigma \rightarrow 0} \phi_{\mathbf{P}}^{\sigma} \equiv \phi_{\mathbf{P}}$ and $\lim_{\sigma \rightarrow 0} E_{\mathbf{P}}^{\sigma} = E_{\mathbf{P}}$ exist.

Proof

I will write again $\tilde{\phi}_{\mathbf{P}}^{\sigma_2} - \tilde{\phi}_{\mathbf{P}}^{\sigma_1}$ in the following way

$$\tilde{\phi}_{\mathbf{P}}^{\sigma_2} - \tilde{\phi}_{\mathbf{P}}^{\sigma_1} = \tilde{\phi}_{\mathbf{P}}^{\sigma_2} - \phi_{\mathbf{P}}^{\epsilon_2(\sigma_2)^{\frac{l+2}{2}}} + \phi_{\mathbf{P}}^{\epsilon_2(\sigma_2)^{\frac{l+2}{2}}} - \phi_{\mathbf{P}}^{\epsilon_1(\sigma_1)^{\frac{m+2}{2}}} + \phi_{\mathbf{P}}^{\epsilon_1(\sigma_1)^{\frac{m+2}{2}}} - \tilde{\phi}_{\mathbf{P}}^{\sigma_1}.$$

Now, given an arbitrarily small δ , there exist $l(\delta), m(\delta)$ sufficiently large and a phase $e^{i\eta(\delta)}$ in which

$$\left\| \phi_{\mathbf{P}}^{\epsilon_1(\sigma_1)^{\frac{m+2}{2}}} - e^{i\eta(\delta)} \phi_{\mathbf{P}}^{\epsilon_2(\sigma_2)^{\frac{l+2}{2}}} \right\| \leq \delta.$$

This is possible essentially because of the convergence established in theorems 2.3 and 2.3bis and because, by construction, the ground state is unique until there is a cut-off.

Therefore $\left\| \tilde{\phi}_{\mathbf{P}}^{\sigma_2} - e^{-i\eta(\delta)} \tilde{\phi}_{\mathbf{P}}^{\sigma_1} \right\|$ can be bounded with a quantity of order $\sigma_2^{\frac{1}{8}} + \sigma_1^{\frac{1}{8}} + \delta$.

moreover

$$\begin{aligned} \left\| P_{\tilde{\phi}_{\mathbf{P}}^{\sigma_1}} \psi_0 - P_{\tilde{\phi}_{\mathbf{P}}^{\sigma_2}} \psi_0 \right\| &= \left\| \tilde{\phi}_{\mathbf{P}}^{\sigma_1} \left(\tilde{\phi}_{\mathbf{P}}^{\sigma_1}, \psi_0 \right) - \tilde{\phi}_{\mathbf{P}}^{\sigma_2} \left(\tilde{\phi}_{\mathbf{P}}^{\sigma_2}, \psi_0 \right) \right\| \leq \\ &\leq \left\| \tilde{\phi}_{\mathbf{P}}^{\sigma_1} \left(\tilde{\phi}_{\mathbf{P}}^{\sigma_1}, \psi_0 \right) - e^{i\eta} \tilde{\phi}_{\mathbf{P}}^{\sigma_2} \left(\tilde{\phi}_{\mathbf{P}}^{\sigma_1}, \psi_0 \right) \right\| + \left\| \tilde{\phi}_{\mathbf{P}}^{\sigma_2} \left(e^{-i\eta} \tilde{\phi}_{\mathbf{P}}^{\sigma_1}, \psi_0 \right) - \tilde{\phi}_{\mathbf{P}}^{\sigma_2} \left(\tilde{\phi}_{\mathbf{P}}^{\sigma_2}, \psi_0 \right) \right\| = \\ &\leq \left\| \tilde{\phi}_{\mathbf{P}}^{\sigma_1} - e^{i\eta} \tilde{\phi}_{\mathbf{P}}^{\sigma_2} \right\| \cdot \left| \left(\tilde{\phi}_{\mathbf{P}}^{\sigma_1}, \psi_0 \right) \right| + \left\| \tilde{\phi}_{\mathbf{P}}^{\sigma_2} \right\| \cdot \left\| \psi_0 \right\| \cdot \left\| e^{-i\eta} \tilde{\phi}_{\mathbf{P}}^{\sigma_1} - \tilde{\phi}_{\mathbf{P}}^{\sigma_2} \right\| \end{aligned}$$

It follows that $\phi_{\mathbf{P}}^{\sigma} \equiv \frac{P_{\tilde{\phi}_{\mathbf{P}}^{\sigma}} \psi_0}{\left\| P_{\tilde{\phi}_{\mathbf{P}}^{\sigma}} \psi_0 \right\|}$ converges strongly to a vector $\phi_{\mathbf{P}}$, with an error of order $\sigma^{\frac{1}{8}}$.

The convergence of $E_{\mathbf{P}}^{\sigma}$ follows from the estimates of the generalized version of lemma 1.3 and from the fact that the hamiltonian $H_{\mathbf{P}}$ is bounded from below.

Lemma 3.3

For $\mathbf{P}$ and $\mathbf{P} + \Delta\mathbf{P}$ belonging to $\Sigma \equiv \{\mathbf{P} : |\mathbf{P}| \leq \sqrt{m}\}$ and m sufficiently large the following Hoelder estimate on the energy gradient holds:

$$|\nabla E^\sigma(\mathbf{P}) - \nabla E^\sigma(\mathbf{P} + \Delta\mathbf{P})| \leq C |\Delta\mathbf{P}|^{\frac{1}{16}}$$

where the constant C is uniform in $0 \leq \sigma < \epsilon$, in $\mathbf{P}, \mathbf{P} + \Delta\mathbf{P} \in \Sigma$ where $|\Delta\mathbf{P}| \leq \left(\frac{1}{2}\right)^{\frac{4}{3}}$.

Proof

The idea is to perturb, in $\mathbf{P}$, the $\nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) \equiv \left(\psi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}}, \frac{\mathbf{P} - \mathbf{P}^{mes}}{m} \psi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} \right)$ where $\psi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}}$ is the normalized ground state of $H_{\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}}$. For this purpose I expand the resolvent

$$\frac{1}{H_{\mathbf{P} + \Delta\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}} \big|_{F^+_{|\Delta\mathbf{P}|^{\frac{1}{4}}}} - E}$$

($E \in \mathcal{C}$ and s.t. $\left| E - E_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} \right| = \frac{|\Delta\mathbf{P}|^{\frac{1}{4}}}{4}$), on the basis of the following information:

- $H_{\mathbf{P} + \Delta\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}} - H_{\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}} = -\frac{\Delta\mathbf{P}}{m} \cdot \mathbf{P}^{mes} + \frac{\Delta\mathbf{P}}{m} \cdot \mathbf{P} + \frac{|\Delta\mathbf{P}|^2}{2m}$;
- $H_{\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}} \big|_{F^+_{|\Delta\mathbf{P}|^{\frac{1}{4}}}}$ has unique ground state $\psi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}}$ of energy $E_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}}$ and corresponding gap bounded from below by $\frac{|\Delta\mathbf{P}|^{\frac{1}{4}}}{2}$ (theorem 1.5 in the continuum case);
- the norm $|\Delta P^i|^{\frac{1}{8}} \left\| \frac{(P^i - P^{mes^i})}{\sqrt{2m}} \left(\frac{1}{H_{\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}} \big|_{F^+_{|\Delta\mathbf{P}|^{\frac{1}{4}}}} - E} \right)^{\frac{1}{2}} \right\|$ is uniformly bounded in $|\Delta\mathbf{P}|$.

Therefore, for a sufficiently large m we have that

$$\sum_i |\Delta P^i|^{\frac{1}{4}} \left\| \left(\frac{1}{H_{\mathbf{P}+\Delta\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}|_{F^+} - E}} \right)^{\frac{1}{2}} \sqrt{\frac{2}{m}} \cdot \frac{(P^i - P^{mes^i})}{\sqrt{2m}} \left(\frac{1}{H_{\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}|_{F^+} - E}} \right)^{\frac{1}{2}} \right\| < \frac{1}{4}$$

$$\bullet \left\| -\frac{1}{2\pi i} \oint \frac{1}{H_{\mathbf{P}+\Delta\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}} - E}} dE \psi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} \right\| \geq 1 - \sum_{n=1}^{\infty} \left(\frac{|\Delta\mathbf{P}|^{\frac{3}{4}}}{2} \right)^n \geq \frac{2}{3}$$

for the constraint $|\Delta\mathbf{P}| \leq \left(\frac{1}{2}\right)^{\frac{4}{3}}$

it follows that:

$$\bullet -\frac{1}{2\pi i} \oint \frac{1}{H_{\mathbf{P}+\Delta\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}} - E}} dE \psi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} \text{ is ground state of } H_{\mathbf{P}+\Delta\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}} ;$$

$$\bullet \left\| \frac{-\frac{1}{2\pi i} \oint \frac{1}{H_{\mathbf{P}+\Delta\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}} - E}} dE \psi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}}}{-\frac{1}{2\pi i} \oint \frac{1}{H_{\mathbf{P}+\Delta\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}} - E}} dE \psi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}}} - \psi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} \right\| \leq C' |\Delta\mathbf{P}|^{\frac{3}{4}}$$

where C' is a constant uniform in $\mathbf{P} \in \Sigma$, $\mathbf{P} + \Delta\mathbf{P} \in \Sigma$ where $\Delta\mathbf{P} \in \left\{ \Delta\mathbf{P} : |\Delta\mathbf{P}| \leq \left(\frac{1}{2}\right)^{\frac{4}{3}} \right\}$.

Since:

$$1) \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P} + \Delta\mathbf{P}) - \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) = \left(\psi_{\mathbf{P}+\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}}, \frac{\mathbf{P}+\Delta\mathbf{P}-\mathbf{P}^{mes}}{m} \psi_{\mathbf{P}+\Delta\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} \right) - \left(\psi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}}, \frac{\mathbf{P}-\mathbf{P}^{mes}}{m} \psi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} \right)$$

$$2) H_{\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}} + 2\pi g^2 \kappa - \frac{(\mathbf{P}^{mes} - \mathbf{P})^2}{2m} \geq 0$$

we can conclude that

$$\left| \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) - \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P} + \Delta\mathbf{P}) \right| \leq C'' |\Delta\mathbf{P}|^{\frac{3}{4}} \quad (20)$$

where C'' is constant uniform in $\mathbf{P} \in \Sigma$, $\mathbf{P} + \Delta\mathbf{P} \in \Sigma$ where $\Delta\mathbf{P} \in \left\{ \Delta\mathbf{P} : |\Delta\mathbf{P}| \leq \left(\frac{1}{2}\right)^{\frac{4}{3}} \right\}$, and of order $\frac{1}{\sqrt{m}}$.

If $\sigma < |\Delta\mathbf{P}|^{\frac{1}{4}}$, in order to obtain the thesis of the lemma, I take advantage of the result of theorem 2.3bis together with lemma A2 in Appendix A:

$$\left| \nabla E^{|\Delta \mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) - \nabla E^\sigma(\mathbf{P}) \right| \leq C |\Delta \mathbf{P}|^{\frac{1}{16}} \text{ for } \mathbf{P} \in \Sigma.$$

If $\sigma \geq |\Delta \mathbf{P}|^{\frac{1}{4}}$ an estimate analogous to (20) holds.

Theorem 3.4

Under the hypotheses of lemma 3.3, the norm difference between $\phi_{\mathbf{P}}^\sigma$ and $\phi_{\mathbf{P}+\Delta \mathbf{P}}^\sigma$ is Hoelder with coefficient $\frac{1}{32}$ and multiplicative constant that is uniform in $0 \leq \sigma < \epsilon$, in $\mathbf{P}, \mathbf{P} + \Delta \mathbf{P} \in \Sigma$ where $|\Delta \mathbf{P}| \leq \left(\frac{1}{6}\right)^4$.

Proof

Preliminary definitions:

$$\begin{aligned} H_{\mathbf{P}, |\Delta \mathbf{P}|^{\frac{1}{4}}}^w &= \frac{1}{2m} \left(\mathbf{P}^{mes} - g \int_{|\Delta \mathbf{P}|^{\frac{1}{4}}}^\kappa \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{|\Delta \mathbf{P}|^{\frac{1}{4}}}(\mathbf{P})\right)} \left(b(\mathbf{k}) + b^\dagger(\mathbf{k})\right) d^3k \right)^2 \\ &+ \int \left(|\mathbf{k}| - \mathbf{k} \cdot \nabla E^{|\Delta \mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k + c_{\mathbf{P}} \left(|\Delta \mathbf{P}|^{\frac{1}{4}} \right) \\ H_{\mathbf{P}+\Delta \mathbf{P}, |\Delta \mathbf{P}|^{\frac{1}{4}}}^w &= \frac{1}{2m} \left(\mathbf{P}^{mes} - g \int_{|\Delta \mathbf{P}|^{\frac{1}{4}}}^\kappa \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{|\Delta \mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}+\Delta \mathbf{P})\right)} \left(b(\mathbf{k}) + b^\dagger(\mathbf{k})\right) d^3k \right)^2 + \\ &+ \int \left(|\mathbf{k}| - \mathbf{k} \cdot \nabla E^{|\Delta \mathbf{P}|^{\frac{1}{4}}}(\mathbf{P} + \Delta \mathbf{P}) \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k + c_{\mathbf{P}+\Delta \mathbf{P}} \left(|\Delta \mathbf{P}|^{\frac{1}{4}} \right) \\ H_{\mathbf{P}+\Delta \mathbf{P}, |\Delta \mathbf{P}|^{\frac{1}{4}}}^w - H_{\mathbf{P}, |\Delta \mathbf{P}|^{\frac{1}{4}}}^w &= c_{\mathbf{P}+\Delta \mathbf{P}} \left(|\Delta \mathbf{P}|^{\frac{1}{4}} \right) - c_{\mathbf{P}} \left(|\Delta \mathbf{P}|^{\frac{1}{4}} \right) + \\ &+ \int \left(-\mathbf{k} \cdot \nabla E^{|\Delta \mathbf{P}|^{\frac{1}{4}}}(\mathbf{P} + \Delta \mathbf{P}) + \mathbf{k} \cdot \nabla E^{|\Delta \mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k + \\ &+ \frac{1}{2m} \left(\Pi_{\mathbf{P}, |\Delta \mathbf{P}|^{\frac{1}{4}}} + \int_{|\Delta \mathbf{P}|^{\frac{1}{4}}}^\kappa \left[g \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{|\Delta \mathbf{P}|^{\frac{1}{4}}}(\mathbf{P})\right)} - g \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{|\Delta \mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}+\Delta \mathbf{P})\right)} \right] (b + b^\dagger) d^3k \right)^2 + \\ &- \frac{1}{2m} \left(\Pi_{\mathbf{P}, |\Delta \mathbf{P}|^{\frac{1}{4}}} \right)^2 = \end{aligned}$$

$$\begin{aligned}
&= c_{\mathbf{P}+\Delta\mathbf{P}} \left(|\Delta\mathbf{P}|^{\frac{1}{4}} \right) - c_{\mathbf{P}} \left(|\Delta\mathbf{P}|^{\frac{1}{4}} \right) + \\
&+ \int \left(\mathbf{k} \cdot \left(\nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) - \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P} + \Delta\mathbf{P}) \right) \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3k + \\
&+ \frac{1}{2m} \left(g \int_{|\Delta\mathbf{P}|^{\frac{1}{4}}}^\kappa \frac{\widehat{\mathbf{k}} \cdot \left(\nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) - \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P} + \Delta\mathbf{P}) \right)}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P} + \Delta\mathbf{P}) \right) \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) \right)} \left(b(\mathbf{k}) + b(\mathbf{k})^\dagger \right) d^3k \right)^2 + \\
&\frac{1}{2m} \Pi_{\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}} \left(g \int_{|\Delta\mathbf{P}|^{\frac{1}{4}}}^\kappa \frac{\widehat{\mathbf{k}} \cdot \left(\nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) - \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P} + \Delta\mathbf{P}) \right)}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P} + \Delta\mathbf{P}) \right) \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) \right)} \left(b(\mathbf{k}) + b(\mathbf{k})^\dagger \right) d^3k \right) + \\
&+ \left(g \int_{|\Delta\mathbf{P}|^{\frac{1}{4}}}^\kappa \frac{\widehat{\mathbf{k}} \cdot \left(\nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) - \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P} + \Delta\mathbf{P}) \right)}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P} + \Delta\mathbf{P}) \right) \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) \right)} \left(b(\mathbf{k}) + b(\mathbf{k})^\dagger \right) d^3k \right) \frac{1}{2m} \Pi_{\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}}
\end{aligned}$$

Considering that:

- the estimate (20) in lemma 3.3 holds:
 $\left| \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P}) - \nabla E^{|\Delta\mathbf{P}|^{\frac{1}{4}}}(\mathbf{P} + \Delta\mathbf{P}) \right| \leq C'' |\Delta\mathbf{P}|^{\frac{3}{4}};$
- the operator $\frac{H^w_{\mathbf{P}+\Delta\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}} - H^w_{\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}}}{|\Delta\mathbf{P}|^{\frac{3}{4}}}$ is form-bounded with respect to $H^w_{\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}}$;
- the gap of $E_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}}$ (that is the ground eigenvalue of $H^w_{\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}} |_{F^+_{|\Delta\mathbf{P}|^{\frac{1}{4}}}}$) is bounded from below by $\frac{|\Delta\mathbf{P}|^{\frac{1}{4}}}{2}$ (generalized version of theorem 1.5);

for the Kato's theorem on the analytic perturbation, the vector

$$\begin{aligned}
\phi_{\mathbf{P}+\Delta\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} &\equiv \frac{-\frac{1}{2\pi i} \oint_{\mathbf{P}+\Delta\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}} \frac{1}{H^w - E} dE \psi_0}{\left\| -\frac{1}{2\pi i} \oint_{\mathbf{P}+\Delta\mathbf{P}, |\Delta\mathbf{P}|^{\frac{1}{4}}} \frac{1}{H^w - E} dE \psi_0 \right\|} \\
& (E \in \mathcal{C} \text{ and s.t. } \left| E - E_{\mathbf{P}+\Delta\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} \right| = \frac{|\Delta\mathbf{P}|^{\frac{1}{4}}}{4})
\end{aligned}$$

can be obtained, for m sufficiently large, perturbing $\phi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}}$.

From the perturbation we have the estimate:

$$\left\| \phi_{\mathbf{P}+\Delta\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} - \phi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} \right\| \leq C''' |\Delta\mathbf{P}|^{\frac{1}{4}} \quad (20.1)$$

where the constant C''' is uniform in $\mathbf{P}, \mathbf{P} + \Delta\mathbf{P} \in \Sigma$ where $|\Delta\mathbf{P}| \leq \left(\frac{1}{6}\right)^4$.

In conclusion, for $\sigma < |\Delta\mathbf{P}|^{\frac{1}{4}}$, using a generalized version of corollary 2.4 for σ in the continuum, we have that:

$$\begin{aligned} & \left\| \phi_{\mathbf{P}+\Delta\mathbf{P}}^{\sigma} - \phi_{\mathbf{P}}^{\sigma} \right\| = \\ & = \left\| \phi_{\mathbf{P}+\Delta\mathbf{P}}^{\sigma} - \phi_{\mathbf{P}+\Delta\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} + \phi_{\mathbf{P}+\Delta\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} - \phi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} + \phi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} - \phi_{\mathbf{P}}^{\sigma} \right\| \leq \\ & \leq \left\| \phi_{\mathbf{P}+\Delta\mathbf{P}}^{\sigma} - \phi_{\mathbf{P}+\Delta\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} \right\| + \left\| \phi_{\mathbf{P}+\Delta\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} - \phi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} \right\| + \left\| \phi_{\mathbf{P}}^{|\Delta\mathbf{P}|^{\frac{1}{4}}} - \phi_{\mathbf{P}}^{\sigma} \right\| \leq \\ & \leq |\Delta\mathbf{P}|^{\frac{1}{32}} + |\Delta\mathbf{P}|^{\frac{1}{32}} + C''' |\Delta\mathbf{P}|^{\frac{1}{4}} \end{aligned}$$

If $\sigma \geq |\Delta\mathbf{P}|^{\frac{1}{4}}$ an estimate analogous to (20.1) holds.

Part II

Scattering theory.

The infrared features of the model produce some difficulties in understanding the scattering:

- the arbitrarily large number of mesons involved in the scattering makes the explanation of the asymptotic decoupling difficult. In particular a problem of consistency arises between the total emission of an infinite number of massless particles (mesons) and a “free” asymptotic dynamics for the nucleon. In other words we can say that a definition of “free” dynamics for the infra-particle (the nucleon) is required;
- the L.S.Z. Weyl meson operators (see below) do not converge on generic vectors of the Hilbert space

$$e^{iHt}e^{-iH^{mes.}t}e^{i(a(\varphi)+a^\dagger(\varphi))}e^{iH^{mes}t}e^{-iHt} \quad \varphi(\mathbf{y}) = \int e^{-i\mathbf{k}\mathbf{y}} \tilde{\varphi}(\mathbf{k}) d^3k \in S(R^3)$$

at least if one uses the same technique of the proof in [2].

Due to the absence of one particle states, from a conceptual point of view it is not possible to use the Haag and Ruelle theory for the construction of scattering states. Nevertheless, the decoupling mechanism in the H. and R. theory can be reproduced in terms of fixed time locality properties of the meson field and of the “current density field” of the nucleon .

According to this interpretative scheme I review the construction of the asymptotic nucleon, which was considered in [2]. My purposes are two:

- the first one is to give a minimal (with respect to the meson cloud) description of the nucleon out of the scattering
- the second one consists in finding a subspace of states which will be used in or-

der to prove the asymptotic convergence of the massless field. This subspace is the analogous of the one particle subspace in the regularized case (infrared cut-off in the interaction).

The starting point is the construction of the vector $\psi_G(t)$, at time t , that will approximate the minimal asymptotic nucleon state $\psi_G^{out(in)}$:

$\psi_G(t)$ is constructed starting from a one particle state for the hamiltonian H_{σ_t} , of wave function G in $\mathbf{P}$ -variables, to which a Weyl operator, in properly evolved meson variables (L.S.Z. Weyl operators, see later), is applied. The smearing function in Weyl operators has the right infrared behavior established in the spectral analysis. Its spectral distribution, near $\mathbf{k} = 0$, is labelled by the asymptotic nucleon mean velocity (constructed “a posteriori”); its frequency support goes from σ_t ($\sigma_t \rightarrow 0$ for $|t| \rightarrow +\infty$) to an arbitrarily small $\kappa_1 \neq 0$.

Intuitively, in the asymptotic limit it describes a “free” (one particle state) nucleon plus a **not totally removable** cloud of asymptotic mesons. The main difference with an analogous construction by Froehlich [1] is that the infrared cut-off of the cloud of interacting mesons is removed only asymptotically at a rate faster than $\frac{1}{t}$ (in accordance with the indetermination principle). The advantage consists in the fact that the construction of the vector $\psi_G(t)$ is always inside the Hilbert space and it makes it simpler to use locality (by the nucleon position $\mathbf{x}$) in proving the convergence. Moreover, the removal of the infrared cut-off is an “a posteriori” result and a byproduct of the decoupling.

I want to stress the fact that the coherent function in the definition of the minimal asymptotic nucleon states is arbitrary except for the infrared limit. Nevertheless the conceptual and mathematical role of the minimal asymptotic nucleon states is very useful in developing the scattering theory.

The mathematical counterpart of what I said is the following.

We construct a set of states $\psi_G(t)$ (where G is the $\mathbf{P}$ -wave function of the one σ_t -particle state) on which there exist the limits

$$s - \lim_{t \rightarrow \pm\infty} \psi_G(t) \equiv \psi_G^{out(in)}$$

Moreover on these states the functions f , continuous and of compact support, of the nucleon mean velocity have limit :

$$s - \lim_{t \rightarrow \pm\infty} e^{iHt} f\left(\frac{\mathbf{x}}{t}\right) e^{-iHt} \psi_G^{out(in)} = s - \lim_{t \rightarrow \pm\infty} e^{iHt} f\left(\frac{\mathbf{x}}{t}\right) e^{-iHt} \psi_G(t) \cong \psi_{f.G}^{out(in)}$$

The norm closure of the finite linear combinations $\{\psi_G^{out(in)}\}$ substitutes the subspace of states at one nucleon which we have in the regularized case

$$\begin{aligned} s - \lim_{t \rightarrow \pm\infty} e^{iHt} e^{-iH^{mes}t} e^{i(a(\varphi)+a^\dagger(\varphi))} e^{iH^{mes}t} e^{-iHt} \psi_G^{out(in)} &= \\ = s - \lim_{t \rightarrow \pm\infty} e^{iHt} e^{-iH^{mes}t} e^{i(a(\varphi)+a^\dagger(\varphi))} e^{iH^{mes}t} e^{-iHt} \psi_G(t) &\equiv \psi_{G,\varphi}^{out(in)} \\ (\varphi(\mathbf{y}) = \int e^{-i\mathbf{k}\mathbf{y}} \tilde{\varphi}(\mathbf{k}) d^3k, \tilde{\varphi}(\mathbf{k}) \in C_0^\infty(R^3 \setminus 0)) & \end{aligned}$$

By analogy, I will define the invariant (under space-time translation) subspaces $H^{1out(in)} \equiv \{\overline{\psi_G^{out(in)}}, G(\mathbf{P}) \in C_0^1(R^3 \setminus 0), \mathbf{P} \in \Sigma\}$.

Then I define $H^{out(in)} \equiv \{\overline{\psi_{G,\varphi}^{out(in)}}, G \in C_0^1(R^3 \setminus 0), \tilde{\varphi} \in C_0^\infty(R^3 \setminus 0)\}$. The last one is obtained by the vectors $\psi_{G,\varphi}^{out(in)}$, starting from $H^{1out(in)}$.

As I already said, these definitions are arbitrary in some sense. Nevertheless, through the (artificial) separation between $H^{1out(in)}$ and $H^{out(in)}$ I want to point out that:

- from a technical point of view, my construction of the scattering states is based on some (arbitrary) $H^{1out(in)}$;

- from a physical point of view, even if the meson cloud described by smearing functions φ is totally removable, the meson cloud linked to the vectors in $H^{1out(in)}$ is not completely removable. Then all the scattering states always contain asymptotic mesons, those ones of the spaces $H^{1out(in)}$ involved in the construction of the spaces $H^{out(in)}$.

Now, we make an observation which regards the coherent “static” factor $g \frac{1}{|\mathbf{k}| \sqrt{2|\mathbf{k}|(1-\hat{\mathbf{k}} \cdot \nabla E)}}$ founded in the spectral analysis. If we consider a σ -cut-off dynamics, we can easily compute the asymptotic nucleon mean velocity. This operator coincides with ∇E^σ if it is applied to the one nucleon particle states. Through a non-rigorous removal of the cut-off, the coherent factor can be thought in terms of the asymptotic nucleon mean velocity to be constructed.

Content of the chapters.

In chapter 4, I define the approximating vector $\psi_G(t)$ for the generic state of minimal asymptotic nucleon. Then I study the norm in time, and finally I prove its strong

convergence (paragraphs 4.1 and 4.2).

In chapter 5, I construct the scattering subspaces $H^{out(in)}$. Then I prove the existence of the asymptotic limits of the functions, continuous and of compact support, of the nucleon mean velocity. I also prove the existence of the asymptotic limit of L.S.Z. meson field Weyl operators and I discuss their commutation properties.

The construction will be explicitly performed in the case “out”. The case “in” is completely analogous.

4 Approximating vector $\psi_G(t)$.

I will consider a cubic region of volume $V = L^3$ inside Σ in the $\mathbf{P}$ -space, for simplicity of construction. For $\mathbf{P}$ in this region, we have:

- the existence of the ground state of $H_{\mathbf{P}}^g$ with the properties which implicitly follow from the results in chapter 3;
- small nucleon velocities $|\nabla E^\sigma(\mathbf{P})| < 1 \quad \forall \sigma$ (see lemma A2);
- $|\nabla E^\sigma(\mathbf{P} + \mathbf{k})| < 1 \quad \forall \sigma$, for $\mathbf{k} : 0 \leq |\mathbf{k}| \leq \kappa_1 \ll 1$ (κ_1 properly small).

I will assume that the following hypothesis holds in Σ :

hypothesis B1 $\exists m_r > 0 \quad \text{s.t.} \quad \forall \sigma \quad \forall \mathbf{P} \in \Sigma \quad \frac{\partial E^\sigma(\mathbf{P})}{\partial |\mathbf{P}|} \geq \frac{|\mathbf{P}|}{m_r} \quad \text{and} \quad \frac{\partial^2 E^\sigma(\mathbf{P})}{\partial^2 |\mathbf{P}|} \geq \frac{1}{m_r}$

I consider a time-dependent ($t \gg 1$) cell partition of the volume $V = L^3$. The linear dimension of each cell is $\frac{L}{2^n}$, where $n \in \mathbb{N}$, $n \geq 1$ is such that $(2^n)^{\frac{1}{\epsilon}} \leq t < (2^{n+1})^{\frac{1}{\epsilon}}$, $\epsilon > 0$. The exponent ϵ will be fixed only “a posteriori”. It follows that the number of cells is $N(t) = (2^n)^3$ where $n = [\log_2 t^\epsilon]$. I will call Γ_i the i^{th} cell, centered in $\overline{\mathbf{P}}_i$

Constructive prescription of $\psi_G(t)$.

1) I will consider the vector $\psi_{i,\sigma_t}^{(t)} = \int_{\Gamma_i} G(\mathbf{P}) \psi_{\mathbf{P},\sigma_t} d^3P$, where:

- $G(\mathbf{P}) \in C_0^1(R^3 \setminus 0)$;
- $\psi_{\mathbf{P}}^{\sigma_t} \equiv W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_t})\phi_{\mathbf{P}}^{\sigma_t}$ is the normalized ground state of $H_{\mathbf{P},\sigma_t}$ (I will use the index b in $W_{\sigma_{t_2}}^{b^\dagger}$ in order to distinguish it from Weyl operators in $a, a^\dagger$ variables);
- (t) is referred to the partition at time t ;
- $\|\psi_{i,\sigma_t}^{(t)}\| = \left(\int_{\Gamma_i} |G(\mathbf{P})|^2 d^3P\right)^{\frac{1}{2}}$ is of order $(N(t))^{-\frac{1}{2}}$

2)

I will dress each $\psi_{i,\sigma_t}^{(t)}$ by the proper $e^{iHt}e^{-iH^{mes}t}W_{\sigma_t}(\mathbf{v}_i)e^{iH^{mes}t}e^{-iH_{\sigma_t}t}$. In such a way, the vector will remain inside the Hilbert space $\mathcal{H}$ under removal of the cut-off. Therefore I will define:

$$\begin{aligned}\psi_G(t) &\equiv e^{iHt}e^{-iH^{mes}t}\sum_{i=1}^{N(t)} W_{\sigma_t}(\mathbf{v}_i)e^{iH^{mes}t}e^{i\gamma_{\sigma_t}(\mathbf{v}_i,\nabla E^{\sigma_t}(\mathbf{P}),t)}e^{-iE^{\sigma_t}(\mathbf{P})t}\psi_{i,\sigma_t}^{(t)} = \\ &= e^{iHt}\sum_{i=1}^{N(t)} W_{\sigma_t}(\mathbf{v}_i,t)e^{i\gamma_{\sigma_t}(\mathbf{v}_i,\nabla E^{\sigma_t}(\mathbf{P}),t)}e^{-iE^{\sigma_t}t}\psi_{i,\sigma_t}^{(t)}\end{aligned}$$

being

- $W_{\sigma_t}(\mathbf{v}_i) = e^{-g\int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k}) - a^\dagger(\mathbf{k})}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3k}{\sqrt{2}|\mathbf{k}|}}$, where $\mathbf{v}_i$ is the velocity corresponding to the cell center $\overline{\mathbf{P}}_i$ of Γ_i , $\mathbf{v}_i \equiv \nabla E^{\sigma_t}(\overline{\mathbf{P}}_i)$, κ_1 ($0 < \kappa_1 \ll 1$) is the integration upper bound for the frequency;
- $W_{\sigma_t}(\mathbf{v}_i, t) = e^{-iH^{mes}t}W_{\sigma_t}(\mathbf{v}_i)e^{iH^{mes}t} = e^{-g\int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|t} - a^\dagger(\mathbf{k})e^{-i|\mathbf{k}|t}}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3k}{\sqrt{2}|\mathbf{k}|}}$;
- $\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), t) = -\int_1^t \left\{ g^2 \int_{\sigma_t}^{\sigma_\tau^S} \int \frac{\cos(\mathbf{k} \cdot \nabla E^{\sigma_t}(\mathbf{P})\tau - |\mathbf{k}|\tau)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| \right\} d\tau =$
 $= -\int_1^t \left\{ g^2 \int_{\sigma_t}^{\sigma_\tau^S} \int \frac{\cos(\mathbf{q} \cdot \nabla E^{\sigma_t}(\mathbf{P}) - |\mathbf{q}|)}{(1 - \widehat{\mathbf{q}} \cdot \mathbf{v}_i)} d\Omega \frac{d|\mathbf{q}|}{\tau} \right\} d\tau$

$e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), t)}$ is a phase factor whose origin and definition will be clear later. In particular, in order to implement the convergence of $\psi_G(t)$, we will see that the integration bound σ_τ^S is a “slow” cut-off that will be substantially treated as a fixed cut-off, while the “speed” cut-off σ_τ will require the phase term. For this purpose it is necessary that σ_τ^S goes like $\tau^{-\alpha}$, where α is a positive number

sufficiently less than 1, and that σ_τ is of order $\tau^{-\beta}$, where β is sufficiently bigger than 1. In the proofs, I will take $\alpha = \frac{39}{40}$ and I will not fix the exponent β . On the basis of partial estimates, eventually β will be chosen equal to 128 in order to gain the strong convergence of the vector $\psi_G(t)$.

4.1 Control of the norm of $\psi_G(t)$.

I will study the scalar product:

$$\begin{aligned} & (\psi_G(t), \psi_G(t)) = \\ & = \sum_{i,j=1}^{N(t)} \left(W_{\sigma_t}(\mathbf{v}_i, t) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), t)} e^{-iH_{\sigma_t}t} \psi_{i,\sigma_t}^{(t)}, W_{\sigma_t}(\mathbf{v}_j, t) e^{i\gamma_{\sigma_t}(\mathbf{v}_j, \nabla E^{\sigma_t}(\mathbf{P}), t)} e^{-iH_{\sigma_t}t} \psi_{j,\sigma_t}^{(t)} \right) \end{aligned}$$

The diagonal terms of the above sum are easily under control. In fact, if $i = j$ we obtain:

$$\sum_{i=1}^{N(t)} \left(\psi_{i,\sigma_t}^{(t)}, \psi_{i,\sigma_t}^{(t)} \right) = \sum_{i=1}^{N(t)} \int_{\Gamma_i} |G(\mathbf{P})|^2 d^3P = \int_V |G(\mathbf{P})|^2 d^3P$$

The following step consists in proving that each mixed term of the sum $\sum_{i,j=1}^{N(t)}$ vanishes asymptotically with an order in t independent of the dimension of the cell. Therefore, by properly choosing the exponent ϵ that determines the rate of growth of $N(t) = [t^\epsilon]^3$, we obtain that the sum of mixed terms for $t \rightarrow +\infty$ vanishes.

Remarks on the notations

In the estimates I will produce in these paragraphs, I will generically call C the multiplicative constants which are uniform in the infrared cut-off and in the cells partition of the volume V . The bounds are intended from above, up to a different explicit warning. The time t is intended much greater than 1.

mixed terms

If $i \neq j$, I define:

$$M_{i,j}(t) \equiv \left(e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), t)} \psi_{i,\sigma_t}^{(t)}, e^{iH_{\sigma_t}t} W_{\sigma_t,i,j}(t) e^{i\gamma_{\sigma_t}(\mathbf{v}_j, \nabla E^{\sigma_t}(\mathbf{P}), t)} e^{-iH_{\sigma_t}t} \psi_{j,\sigma_t}^{(t)} \right)$$

$$W_{\sigma_t, i, j}(t) \equiv e^{-\int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|t} - a^\dagger(\mathbf{k})e^{-i|\mathbf{k}|t}}{|\mathbf{k}|} \cdot h_{i, j}(\widehat{\mathbf{k}}) \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} \text{ where } h_{i, j}(\widehat{\mathbf{k}}) = \frac{g\widehat{\mathbf{k}} \cdot (\mathbf{v}_j - \mathbf{v}_i)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_j) \cdot (1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)}$$

Now, I consider $M_{i, j}(t)$ as a two-variable function, by distinguishing the time variable t which parametrizes the partition from the time variable s of the evolution. For this purpose, I define:

$$\widehat{M}_{i, j}(t, s) \equiv \left(e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), s)} \psi_{i, \sigma_t}^{(t)}, e^{iH_{\sigma_t} s} W_{\sigma_t, i, j}(s) e^{i\gamma_{\sigma_t}(\mathbf{v}_j, \nabla E^{\sigma_t}(\mathbf{P}), s)} e^{-iH_{\sigma_t} s} \psi_{j, \sigma_t}^{(t)} \right)$$

with the constraint $s > t$ and the obvious property $\widehat{M}_{i, j}(t, t) = M_{i, j}(t)$.

I verify that:

$$\text{I)} \quad \widehat{M}_{i, j}(t, +\infty) = \lim_{s \rightarrow +\infty} \widehat{M}_{i, j}(t, s) = 0$$

$$\begin{aligned} \text{II)} \quad |M_{i, j}(t)| &= \left| \widehat{M}_{i, j}(t, t) - \widehat{M}_{i, j}(t, +\infty) + \widehat{M}_{i, j}(t, +\infty) \right| = \left| \widehat{M}_{i, j}(t, t) - \widehat{M}_{i, j}(t, +\infty) \right| \leq \\ &\leq \left| \int_t^{+\infty} \frac{d}{ds} \left(e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)}, e^{iH_{\sigma_t} s} W_{\sigma_t, i, j}(s) e^{i\gamma_{\sigma_t}(\mathbf{v}_j, \nabla E_{\sigma_t}, s)} e^{-iH_{\sigma_t} s} \psi_{j, \sigma_t}^{(t)} \right) ds \right| \leq \\ &\leq C \cdot t^{2\delta-3\epsilon} \cdot t^{-\frac{1}{112}} \cdot (\ln \sigma_t)^2. \end{aligned}$$

Proof of I)

For $s > t$

$$\widehat{M}_{i, j}(\lambda, t, s) \equiv \left(e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), s)} \psi_{i, \sigma_t}^{(t)}, e^{iH_{\sigma_t} s} W_{\sigma_t, i, j}^\lambda(s) e^{i\gamma_{\sigma_t}(\mathbf{v}_j, \nabla E^{\sigma_t}(\mathbf{P}), s)} e^{-iH_{\sigma_t} s} \psi_{j, \sigma_t} \right)$$

where:

$$\text{- } W_{\sigma_t, i, j}^\lambda(s) \equiv e^{-\lambda \int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|s} - a^\dagger(\mathbf{k})e^{-i|\mathbf{k}|s}}{|\mathbf{k}|} \cdot h_{i, j}(\widehat{\mathbf{k}}) \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}}, \lambda \text{ a real parameter;}$$

$$\begin{aligned} \text{- } \gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), s) &= \left\{ \begin{aligned} & - \int_1^s \left\{ g^2 \int_{\tau \cdot \sigma_t}^{\tau \cdot \tau} \tau^{-\frac{39}{40}} \int \frac{\cos(\mathbf{q} \cdot \nabla E^{\sigma_t}(\mathbf{P}) - |\mathbf{q}|)}{(1 - \widehat{\mathbf{q}} \cdot \mathbf{v}_i)} d\Omega \frac{d|\mathbf{q}|}{\tau} \right\} d\tau & \text{for } s : \quad s^{-\frac{39}{40}} \geq \sigma_t \\ & \gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), \sigma_t^{-\frac{40}{39}}) & \text{for } s : \quad s^{-\frac{39}{40}} < \sigma_t \end{aligned} \right. \end{aligned}$$

From the derivation with respect to the real parameter λ , we arrive at the differential equation:

$$\frac{d\widehat{M}_{i,j}(\lambda,t,s)}{d\lambda} = -\lambda C_{i,j} \widehat{M}_{i,j}(\lambda,t,s) + r_{\sigma}(\lambda,t,s) \quad (21)$$

($\psi_{i,\sigma_t}^{(t)} \in D(H_{\sigma_t})$, then it belongs to $D(a(f))$ and $D(a^\dagger(f))$, $f \in L^2(R^3)$, and the derivative with respect to λ is well defined)

where

$$C_{i,j} = \int_{\sigma_t}^{\kappa_1} \left| h_{i,j}(\widehat{\mathbf{k}}) \right|^2 \frac{d^3 k}{2|\mathbf{k}|^3}$$

$$r_{\sigma_t}(\lambda,t,s) =$$

$$\begin{aligned} &= - \left(W_{\sigma_t,i,j}^{\lambda\dagger}(s) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}),s)} e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)}, \int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|s}}{|\mathbf{k}|} \cdot h_{i,j}(\widehat{\mathbf{k}}) \frac{d^3 k}{\sqrt{2}|\mathbf{k}|} e^{i\gamma_{\sigma_t}(\mathbf{v}_j, \nabla E^{\sigma_t}(\mathbf{P}),s)} e^{-iH_{\sigma_t}s} \psi_{j,\sigma_t}^{(t)} \right) + \\ &+ \left(\int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|s}}{|\mathbf{k}|} \cdot h_{i,j}(\widehat{\mathbf{k}}) \frac{d^3 k}{\sqrt{2}|\mathbf{k}|} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}),s)} e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)}, W_{\sigma_t,i,j}^{\lambda}(s) e^{i\gamma_{\sigma_t}(\mathbf{v}_j, \nabla E^{\sigma_t}(\mathbf{P}),s)} e^{-iH_{\sigma_t}s} \psi_{j,\sigma_t}^{(t)} \right) \end{aligned}$$

$$\text{from which } \widehat{M}_{i,j}(1,t,s) = e^{-\frac{C_{i,j}}{2}} \widehat{M}_{i,j}(0,t,s) + \int_0^1 r_{\sigma_t}(\lambda,t,s) d\lambda$$

Note that:

- $\widehat{M}_{i,j}(0,t,s) = 0 \forall t,s$ since the $\mathbf{P}$ -supports of $\psi_{j,\sigma_t}^{(t)}, \psi_{i,\sigma_t}^{(t)}$ ($i \neq j$) are disjoint;
- thanks to lemma B2 (Appendix B) and by the control of the derivative with respect to s (theorem B4) one can verify the existence of

$$\begin{aligned} &s - \lim_{s \rightarrow +\infty} e^{iH_{\sigma_t}s} \int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|s}}{|\mathbf{k}|} \cdot h_{i,j}(\widehat{\mathbf{k}}) \frac{d^3 k}{\sqrt{2}|\mathbf{k}|} e^{-iH_{\sigma_t}s} e^{i\gamma_{\sigma_t}(\mathbf{v}_j, \nabla E^{\sigma_t}(\mathbf{P}),s)} \psi_{j,\sigma_t}^{(t)} \equiv \\ &\equiv a_{\sigma_t}^{\text{out}}(h) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}),\sigma_t^{-\frac{40}{39}})} \psi_{j,\sigma_t}^{(t)} \end{aligned}$$

- since the vector $\psi_{i,\sigma_t}^{(t)}$ is a Fock state for $\{a_{\sigma_t}^{\text{out}}(\mathbf{k}), \mathbf{k} : 0 \leq |\mathbf{k}| \leq \kappa_1\}$ (see theorem B5, Appendix B) $\lim_{s \rightarrow +\infty} r_{\sigma_t}(\lambda,t,s) = 0$; in fact

$$\begin{aligned} &\lim_{s \rightarrow +\infty} e^{iH_{\sigma_t}s} \int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|s}}{|\mathbf{k}|} \cdot h_{i,j}(\widehat{\mathbf{k}}) \frac{d^3 k}{\sqrt{2}|\mathbf{k}|} e^{-iH_{\sigma_t}s} e^{i\gamma_{\sigma_t}(\mathbf{v}_j, \nabla E^{\sigma_t}(\mathbf{P}),s)} \psi_{j,\sigma_t}^{(t)} = \\ &= \lim_{s \rightarrow +\infty} e^{iH_{\sigma_t}s} \int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|s}}{|\mathbf{k}|} \cdot h_{i,j}(\widehat{\mathbf{k}}) \frac{d^3 k}{\sqrt{2}|\mathbf{k}|} e^{-iH_{\sigma_t}s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}),\sigma_t^{-\frac{40}{39}})} \psi_{j,\sigma_t}^{(t)} = \end{aligned}$$

$$= a_{\sigma_t}^{out}(h) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), \sigma_t^{-\frac{40}{39}})} \psi_{j, \sigma_t}^{(t)} = 0$$

Therefore, starting from the equation(21), we have

$$\widehat{M}_{i,j}(t, +\infty) = \widehat{M}_{i,j}(\lambda = 1, t, +\infty) = \int_0^1 r_{\sigma_t}(\lambda, t, +\infty) d\lambda = 0$$

Proof of II)

Let us consider:

$$\begin{aligned} & \frac{d}{ds} \left(e^{iH_{\sigma_t}s} W_{\sigma_t}(\mathbf{v}_i, s) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), s)} e^{-iH_{\sigma_t}s} \right) \psi_{i, \sigma_t}^{(t)} = \\ & = ie^{iH_{\sigma_t}s} W_{\sigma_t}(\mathbf{v}_i, s) \left(\varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, s) + \frac{d\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), s)}{ds} \right) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), s)} e^{-iH_{\sigma_t}s} \psi_{i, \sigma_t}^{(t)} \end{aligned} \quad (22)$$

$$\text{where } \varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, s) \equiv g^2 \int_{\sigma_t}^{\kappa_1} \int \frac{\cos(\mathbf{k} \cdot \mathbf{x} - |\mathbf{k}|s)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| \quad (23)$$

Derivation of (22).

The term $ie^{iH_{\sigma_t}s} W_{\sigma_t}(\mathbf{v}_i, s) \varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, s) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} e^{-iH_{\sigma_t}s}$ is formally obtained from:

$$\begin{aligned} & ie^{iH_{\sigma_t}s} \left[H_{\sigma_t} - H^{mes}, e^{-g \int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|s} - a^\dagger(\mathbf{k})e^{-i|\mathbf{k}|s}}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|}} \right] e^{-iH_{\sigma_t}s} = \\ & = ie^{iH_{\sigma_t}s} \left[H_{I, \sigma_t}, e^{-g \int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|s} - a^\dagger(\mathbf{k})e^{-i|\mathbf{k}|s}}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|}} \right] e^{-iH_{\sigma_t}s} = \\ & = ie^{iH_{\sigma_t}s} \left[g \int_{\sigma_t}^{\kappa_1} \left(a(\mathbf{k}) e^{i\mathbf{k} \cdot \mathbf{x}} + a^\dagger(\mathbf{k}) e^{-i\mathbf{k} \cdot \mathbf{x}} \right) \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|}, e^{-g \int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|s} - a^\dagger(\mathbf{k})e^{-i|\mathbf{k}|s}}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|}} \right] e^{-iH_{\sigma_t}s} = \\ & = ie^{iH_{\sigma_t}s} e^{-g \int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|s} - a^\dagger(\mathbf{k})e^{-i|\mathbf{k}|s}}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 \mathbf{k}}{\sqrt{2}|\mathbf{k}|}} g^2 \int_{\sigma_t}^{\kappa_1} \int \frac{\cos(\mathbf{k} \cdot \mathbf{x} - |\mathbf{k}|s)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| e^{-iH_{\sigma_t}s} \end{aligned} \quad (24)$$

the last step follows from:

$$\begin{aligned} & \left[g \int_{\sigma_t}^{\kappa_1} \left(a(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{x}} + a^\dagger(\mathbf{k}) e^{-i\mathbf{k}\cdot\mathbf{x}} \right) \frac{d^3k}{\sqrt{2|\mathbf{k}|}}, -g \int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k}) e^{i|\mathbf{k}|s} - a^\dagger(\mathbf{k}) e^{-i|\mathbf{k}|s}}{|\mathbf{k}|(1-\widehat{\mathbf{k}}\cdot\mathbf{v}_i)} \frac{d^3k}{\sqrt{2|\mathbf{k}|}} \right] = \\ & = g^2 \int_{\sigma_t}^{\kappa_1} \int \frac{\cos(\mathbf{k}\cdot\mathbf{x}-|\mathbf{k}|s)}{(1-\widehat{\mathbf{k}}\cdot\mathbf{v}_i)} d\Omega d|\mathbf{k}| \end{aligned}$$

The formal steps are well defined in $D(H_{\sigma_t})$ from the operatorial point of view, because:

- $H_{\sigma_t}, H^{mes}, H_0$ and $g \int_{\sigma_t}^{\kappa_1} \left(a(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{x}} + a^\dagger(\mathbf{k}) e^{-i\mathbf{k}\cdot\mathbf{x}} \right) \frac{d^3k}{\sqrt{2|\mathbf{k}|}}$ have a common domain D of essential selfadjointness;
- $\frac{d(e^{iH^{mes}s} e^{-iH_{\sigma_t}s})}{ds}$ is closable;
- the sequences, obtained from the formal calculus by approximating (in the norm $\|H_0\psi\| + \|\psi\|$) the vectors in $D(H_{\sigma_t})$ with vectors in D , are convergent.

On the other side $\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s) = -\int_1^s \left\{ g^2 \int_{\sigma_t \cdot \tau}^{\frac{1}{40}} \int \frac{\cos(\mathbf{q}\cdot\nabla E^{\sigma_t}-|\mathbf{q}|)}{(1-\widehat{\mathbf{q}}\cdot\mathbf{v}_i)} d\Omega \frac{d|\mathbf{q}|}{\tau} \right\} d\tau$ from which

$$\begin{aligned} \frac{d\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)}{ds} &= -g^2 \int_{\sigma_t \cdot s}^{\frac{1}{40}} \int \frac{\cos(\mathbf{q}\cdot\nabla E^{\sigma_t}-|\mathbf{q}|)}{(1-\widehat{\mathbf{q}}\cdot\mathbf{v}_i)} d\Omega \frac{d|\mathbf{q}|}{s} \\ \frac{d\gamma_{\sigma_t}}{ds}(\mathbf{v}_i, \frac{\mathbf{x}}{s}, s) &\equiv -g^2 \int_{\sigma_t \cdot s}^{\frac{1}{40}} \int \frac{\cos(\mathbf{q}\cdot\frac{\mathbf{x}}{s}-|\mathbf{q}|)}{(1-\widehat{\mathbf{q}}\cdot\mathbf{v}_i)} d\Omega \frac{d|\mathbf{q}|}{s} \end{aligned}$$

I decompose $\varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, s)$ (formula (23)) in $\varphi_{\sigma_t, \mathbf{v}_i}^1(\mathbf{x}, s) + \varphi_{\sigma_t, \mathbf{v}_i}^2(\mathbf{x}, s)$, which are so defined

$$\begin{aligned} \varphi_{\sigma_t, \mathbf{v}_i}^1(\mathbf{x}, s) &\equiv \\ & g^2 \int_{s \cdot \sigma_t}^{\frac{1}{40}} \int \frac{\cos(\mathbf{k}\cdot\frac{\mathbf{x}}{s}-|\mathbf{k}|)}{(1-\widehat{\mathbf{k}}\cdot\mathbf{v}_i)} d\Omega \frac{d|\mathbf{k}|}{s} \quad for \quad s \leq \sigma_t^{-\frac{40}{39}} \\ &= \begin{cases} 0 & for \quad s > \sigma_t^{-\frac{40}{39}} \end{cases} \\ \varphi_{\sigma_t, \mathbf{v}_i}^2(\mathbf{x}, s) &\equiv \varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, s) - \varphi_{\sigma_t, \mathbf{v}_i}^1(\mathbf{x}, s) = \begin{cases} g^2 \int_{s^{-\frac{39}{40}}}^{\kappa_1} \int \frac{\cos(\mathbf{k}\cdot\mathbf{x}-|\mathbf{k}|s)}{(1-\widehat{\mathbf{k}}\cdot\mathbf{v}_i)} d\Omega d|\mathbf{k}| & for \quad s \leq \sigma_t^{-\frac{40}{39}} \\ \varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, s) & for \quad s > \sigma_t^{-\frac{40}{39}} \end{cases} \end{aligned}$$

The estimate of the norm of the expression (22) requires the introduction of some functions $\chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, s)$ defined in Appendix B. These functions tend, for $s \rightarrow +\infty$, to the characteristic functions of the transformed Γ_i cells, under the application $\mathbf{P} \rightarrow \nabla E^{\sigma_t}(\mathbf{P})$. Such a regularization is necessary to the estimates discussed in Appendix B. As we can see in the definition (b1) in Appendix B, an exponent δ is introduced to which a scale length $t^{-\frac{\delta}{6}}$ corresponds. This scale length has to be less than the (ϵ -dependent) linear dimension of the cell Γ_i in order to have consistency.

Analysis of the expression (22) :

$$\begin{aligned} & \varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, s) e^{-iE^{\sigma_t}(\mathbf{P})s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} + \frac{d\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), s)}{ds} e^{-iE^{\sigma_t}(\mathbf{P})s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} = \\ & = \varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, s) e^{-iE^{\sigma_t}(\mathbf{P})s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \left(1_{\Gamma_i}(\mathbf{P}) - \chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}(\mathbf{P}), s) \right) \psi_{i, \sigma_t}^{(t)} + \\ & + \frac{d\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), s)}{ds} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} e^{-iE^{\sigma_t}(\mathbf{P})s} \left(1_{\Gamma_i}(\mathbf{P}) - \chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}(\mathbf{P}), s) \right) \psi_{i, \sigma_t}^{(t)} + \end{aligned} \quad (25.0)$$

$$+ \varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, s) \left(\chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}(\mathbf{P}), s) - \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right) \right) e^{-iE^{\sigma_t}(\mathbf{P})s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} + \quad (25.1)$$

$$+ \varphi_{\sigma_t, \mathbf{v}_i}^2(\mathbf{x}, s) \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right) e^{-iE^{\sigma_t}(\mathbf{P})s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} + \quad (25.2)$$

$$+ \left(\varphi_{\sigma_t, \mathbf{v}_i}^1(\mathbf{x}, s) \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right) + \frac{d\gamma_{\sigma_t}}{ds}\left(\mathbf{v}_i, \frac{\mathbf{x}}{s}, s\right) \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right) \right) e^{-iE^{\sigma_t}(\mathbf{P})s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} + \quad (25.3)$$

$$- \frac{d\gamma_{\sigma_t}}{ds}\left(\mathbf{v}_i, \frac{\mathbf{x}}{s}, s\right) \left(\chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right) - \chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}(\mathbf{P}), s) \right) e^{-iE^{\sigma_t}(\mathbf{P})s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} + \quad (25.4)$$

$$+ \left(-\frac{d\gamma_{\sigma_t}}{ds}\left(\mathbf{v}_i, \frac{\mathbf{x}}{s}, s\right) + \frac{d\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), s)}{ds} \right) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} e^{-iE^{\sigma_t}(\mathbf{P})s} \chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}(\mathbf{P}), s) \psi_{i, \sigma_t}^{(t)} \quad (25.5)$$

Note

As regards the function $1_{\Gamma_i}(\mathbf{P})$, it is assumed that $\Gamma_i \cap \text{supp}G \neq \emptyset$.

The norm of each term on the right hand side is less than

$$C \cdot s^{-1} \cdot s^{2\delta - \frac{3\epsilon}{2}} \cdot s^{-\frac{1}{112}} \cdot (\ln \sigma_t)^2 \quad (26)$$

(25.0)

- because of lemma B1 and of lemma B3 (*Observation 1*) this term is bounded by $C \cdot \left| \frac{\ln(\sigma_t)}{s} \right| \cdot \frac{1}{s^{\frac{1}{12}}} \cdot t^{-\epsilon}$ ($\delta > 24\epsilon$);

(25.1)

- $|\varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, s)| \leq C \cdot \left| \frac{\ln(\sigma_t)}{s} \right|$ (*Observation 1*, lemma B3);
 $\left\| \left(\chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}(\mathbf{P}), s) \psi_{i, \sigma_t}^{(t)} - \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right) \right) e^{-iE^{\sigma_t}(\mathbf{P})s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} \right\| \leq$
 $\leq C \cdot s^{-\frac{1}{112}} \cdot s^{2\delta} \cdot |\ln \sigma_t| \cdot t^{-\frac{3\epsilon}{2}}$ (see lemma B2);

(25.2)

- $|\varphi_{\sigma_t, \mathbf{v}_i}^2(\mathbf{x}, s) \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right)| \leq C \cdot s^{-2} \cdot s^{\frac{39}{40}}$ as the support of $\chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right)$ leads to the estimate contained in *Observation 2*, lemma B3;

(25.3)

- since by definition of $\gamma_{\sigma_t}(\mathbf{v}_i, \cdot, s)$ we have $\frac{d\gamma_{\sigma_t}}{ds}(\mathbf{v}_i, \frac{\mathbf{x}}{s}, s) \equiv -\varphi_{\sigma_t, \mathbf{v}_i}^1(\mathbf{x}, s)$, this term is zero ;

(25.4)

- by lemma B3 $\left| \frac{d\gamma_{\sigma_t}}{ds}(\mathbf{v}_i, \frac{\mathbf{x}}{s}, s) \right| = |\varphi_{\sigma_t, \mathbf{v}_i}^1(\mathbf{x}, s)| \leq C \cdot \left| \frac{\ln(\sigma_t)}{s} \right|$. Besides, as already seen, $\left\| \left(\chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}(\mathbf{P}), s) \psi_{i, \sigma_t}^{(t)} - \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right) \right) e^{-iE^{\sigma_t}(\mathbf{P})s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} \right\| \leq$
 $\leq C \cdot s^{-\frac{1}{112}} \cdot s^{2\delta} \cdot |\ln \sigma_t| \cdot t^{-\frac{3\epsilon}{2}}$

(25.5)

- from Corollary B2 a bound of order $s^{-1} \cdot s^{-\frac{1}{112}} \cdot s^{2\delta} \cdot |\ln \sigma_t| \cdot t^{-\frac{3\epsilon}{2}}$ follows

Taking into account the estimate (26) and assuming the constraints

- $\delta > 24\epsilon$
- $2\delta + 3\epsilon < \frac{1}{112}$

I consider that:

$$|M_{i,j}(t)| = \left| \widehat{M}_{i,j}(t, t) - \widehat{M}_{i,j}(t, +\infty) + \widehat{M}_{i,j}(t, +\infty) \right| = \left| \widehat{M}_{i,j}(t, t) - \widehat{M}_{i,j}(t, +\infty) \right| \leq$$

$$\begin{aligned}
&\leq \left| \int_t^{+\infty} \frac{d}{ds} \left(e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), s)} \psi_{i, \sigma_t}^{(t)}, e^{iH_{\sigma_t}s} W_{\sigma, i, j}(s) e^{-iH_{\sigma_t}s} e^{i\gamma_{\sigma_t}(\mathbf{v}_j, \nabla E^{\sigma_t}(\mathbf{P}), s)} \psi_{j, \sigma_t}^{(t)} \right) ds \right| \leq \\
&\leq \int_t^{+\infty} 2 \left\| \frac{d}{ds} \left\{ e^{iH_{\sigma_t}s} W_{\sigma_t}(\mathbf{v}_i, s) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), s)} e^{-iH_{\sigma_t}s} \psi_{i, \sigma_t}^{(t)} \right\} \right\| \left\| \psi_{i, \sigma_t}^{(t)} \right\| ds \leq \\
&\leq C \cdot t^{-\frac{1}{112}} \cdot t^{2\delta-3\epsilon} \cdot (\ln \sigma_t)^2
\end{aligned}$$

Therefore, the sum of mixed terms is bounded by $C \cdot t^{-\frac{1}{112}} \cdot t^{2\delta+3\epsilon} \cdot (\ln \sigma_t)^2$ (where $\sigma_t = t^\beta$, $\beta > 1$, by hypothesis) which vanishes for $t \rightarrow +\infty$ if δ, ϵ are such that

- $\delta > 24\epsilon$
- $2\delta + 3\epsilon < \frac{1}{112}$

I will assume these constraints in the following paragraphs.

4.2 Strong convergence of $\psi_G(t)$ for $t \rightarrow \infty$.

I will display the Cauchy property of $\psi_G(t)$, by studying the norm of :

$$\begin{aligned}
&\psi_G(t_2) - \psi_G(t_1) = \\
&= e^{iHt_2} \sum_{i=1}^{N(t_1)} \sum_{l(i)} W_{\sigma_{t_2}}(\mathbf{v}_{l(i)}, t_2) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_{l(i)}, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_2)} e^{-iH_{\sigma_{t_2}}t_2} \psi_{l(i), \sigma_{t_2}}^{(t_2)} + \\
&- e^{iHt_1} \sum_{i=1}^{N(t_1)} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}(\mathbf{P}), t_1)} e^{-iH_{\sigma_{t_1}}t_1} \psi_{i, \sigma_{t_1}}^{(t_1)}
\end{aligned} \tag{27}$$

Notations

- If $t_2 > t_1 \gg 1$ and $N(t_2) \neq N(t_1)$, then $\sum_{j=1}^{N(t_2)} \equiv \sum_{i=1}^{N(t_1)} \sum_{l(i)}$ where $l(i)$ is the index which counts the sub-cells relative to t_2 -partition, contained in the i^{th} cell of the t_1 -partition. Therefore we have $1 \leq l(i) \leq \frac{N(t_2)}{N(t_1)}$;

- adding and subtracting the same quantities, I will rewrite the expression (27) as the sum of three contributions **A1, A2, B**. **A1** is a variation of the “dressing” $W_{\sigma_{t_2}}(\mathbf{v}_i, t_2)$ at fixed partition (that one relative to t_2), **A2** is the variation of the partition, the other variables remain fixed, while **B** is the variation from t_2 to t_1 , at fixed partition (the one relative to t_1).

$$\psi_G(t_2) - \psi_G(t_1) =$$

$$= e^{iHt_2} \sum_{i=1}^{N(t_1)} \sum_{l(i)} W_{\sigma_{t_2}} (\mathbf{v}_{l(i)}, t_2) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_{l(i)}, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_2)} e^{-iH_{\sigma_{t_2}} t_2} \psi_{l(i), \sigma_{t_2}}^{(t_2)} + \quad \mathbf{A1}$$

$$- e^{iHt_2} \sum_{i=1}^{N(t_1)} \sum_{l(i)} W_{\sigma_{t_2}} (\mathbf{v}_i, t_2) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_2)} e^{-iH_{\sigma_{t_2}} t_2} \psi_{l(i), \sigma_{t_2}}^{(t_2)} +$$

$$+ e^{iHt_2} \sum_{i=1}^{N(t_1)} \sum_{l(i)} W_{\sigma_{t_2}} (\mathbf{v}_i, t_2) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_2)} e^{-iH_{\sigma_{t_2}} t_2} \psi_{l(i), \sigma_{t_2}}^{(t_2)} + \quad \mathbf{A2}$$

$$- e^{iHt_2} \sum_{i=1}^{N(t_1)} W_{\sigma_{t_2}} (\mathbf{v}_i, t_2) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_2)} e^{-iH_{\sigma_{t_2}} t_2} \psi_{i, \sigma_{t_2}}^{(t_1)} +$$

$$+ e^{iHt_2} \sum_{i=1}^{N(t_1)} W_{\sigma_{t_2}} (\mathbf{v}_i, t_2) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_2)} e^{-iH_{\sigma_{t_2}} t_2} \psi_{i, \sigma_{t_2}}^{(t_1)} + \quad \mathbf{B}$$

$$- e^{iHt_1} \sum_{i=1}^{N(t_1)} W_{\sigma_{t_1}} (\mathbf{v}_i, t_1) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}(\mathbf{P}), t_1)} e^{-iH_{\sigma_{t_1}} t_1} \psi_{i, \sigma_{t_1}}^{(t_1)}$$

In the following paragraphs I will study the norm **A1,A2,B** in terms of t_1 and t_2 .

analysis of A1

Let us examine the square norm:

$$\left\| e^{iH_{\sigma_{t_2}} t_2} \sum_{i=1}^{N(t_1)} \sum_{l(i)} \left(W_{\sigma_{t_2}} (\mathbf{v}_{l(i)}, t_2) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_{l(i)}, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_2)} - W_{\sigma_{t_2}} (\mathbf{v}_i, t_2) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_2)} \right) e^{-iH_{\sigma_{t_2}} t_2} \psi_{l(i), \sigma_{t_2}}^{(t_2)} \right\|^2 =$$

$$= \sum_{i, l'=1}^{N(t_1)} \sum_{l(i), l'(i)} \left(e^{-iH_{\sigma_{t_2}} t_2} \psi_{l(i), \sigma_{t_2}}^{(t_2)}, (W_{\sigma_{t_2}}^\dagger (\mathbf{v}_{l(i)}, t_2) - W_{\sigma_{t_2}}^\dagger (\mathbf{v}_i, t_2)) (W_{\sigma_{t_2}} (\mathbf{v}_{l'(i)}, t_2) - W_{\sigma_{t_2}} (\mathbf{v}_i, t_2)) e^{-iH_{\sigma_{t_2}} t_2} \psi_{l'(i), \sigma_{t_2}}^{(t_2)} \right)$$

(the phases $e^{i\gamma}$ are omitted for reasons of space)

the sum of the terms where $i' \neq i$ and $l'(i) \neq l(i)$ vanish, for $t_2 \rightarrow +\infty$. Its rate is surely bounded (from above) by of a quantity of order $t_2^{-\frac{1}{112}} \cdot t_2^{2\delta+3\epsilon} \cdot (\ln \sigma_{t_2})^2$ as we can estimate by the same procedure used in the norm control. The remaining terms are of this type

$$\left(e^{-iH_{\sigma_{t_2}} t_2} \psi_{l(i), \sigma_{t_2}}^{(t_2)}, \left(W_{\sigma_{t_2}}^\dagger (\mathbf{v}_{l(i)}, t_2) - W_{\sigma_{t_2}}^\dagger (\mathbf{v}_i, t_2) \right) (W_{\sigma_{t_2}} (\mathbf{v}_{l(i)}, t_2) - W_{\sigma_{t_2}} (\mathbf{v}_i, t_2)) e^{-iH_{\sigma_{t_2}} t_2} \psi_{l(i), \sigma_{t_2}}^{(t_2)} \right) =$$

$$= \left(e^{-iH_{\sigma_{t_2}} t_2} \psi_{l(i), \sigma_{t_2}}^{(t_2)}, \left(2 - W_{\sigma_{t_2}}^\dagger (\mathbf{v}_i, t_2) W_{\sigma_{t_2}} (\mathbf{v}_{l(i)}, t_2) - W_{\sigma_{t_2}}^\dagger (\mathbf{v}_{l(i)}, t_2) W_{\sigma_{t_2}} (\mathbf{v}_i, t_2) \right) e^{-iH_{\sigma_{t_2}} t_2} \psi_{l(i), \sigma_{t_2}}^{(t_2)} \right)$$

For example, I examine:

$$\left(e^{-iH_{\sigma_{t_2}} t_2} \psi_{l(i), \sigma_{t_2}}^{(t_2)}, e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_2)} W_{\sigma_{t_2}}^\dagger (\mathbf{v}_i, t_2) W_{\sigma_{t_2}} (\mathbf{v}_{l(i)}, t_2) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_{l(i)}, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_2)} e^{-iH_{\sigma_{t_2}} t_2} \psi_{l(i), \sigma_{t_2}}^{(t_2)} \right)$$

As in the control of the norm, one considers

$$\left(e^{-iH_{\sigma_{t_2}} s} \psi_{l(i), \sigma_{t_2}}^{(t_2)}, e^{-i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}, s)} W_{\sigma_{t_2}}^\dagger(\mathbf{v}_i, s) W_{\sigma_{t_2}}(\mathbf{v}_{l(i)}, s) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_{l(i)}, \nabla E^{\sigma_{t_2}}, s)} e^{-iH_{\sigma_{t_2}} s} \psi_{l(i), \sigma_{t_2}}^{(t_2)} \right)$$

The limit for $s \rightarrow +\infty$ of the above expression is:

$$e^{-\frac{C_{l(i), i}}{2}} \left(e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_{l(i)}, \nabla E^{\sigma_{t_2}}(\mathbf{P}), \sigma_{t_2}^{-\frac{40}{39}})} \psi_{l(i), \sigma_{t_2}}^{(t_2)}, e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}(\mathbf{P}), \sigma_{t_2}^{-\frac{40}{39}})} \psi_{l(i), \sigma_{t_2}}^{(t_2)} \right)$$

where $C_{l(i), i} = \int_{\sigma_{t_2}}^{\kappa_1} |h_{l(i), i}(\hat{\mathbf{k}})|^2 \frac{d^3 k}{2|\mathbf{k}|^3}$ is bounded by $C \cdot t_1^{-\epsilon} \cdot |\ln(\sigma_{t_2})|$.

Summing on the cells, the total error is bounded by a quantity of order

$$t_2^{-\frac{1}{112}} \cdot t_2^{2\delta} \cdot (\ln \sigma_{t_2})^2$$

Then the discussion is restricted to

$$\sum_{i=1}^{N(t_1)} \sum_{l(i)} \left(2 - e^{-\frac{C_{l(i), i}}{2}} - e^{-\frac{C_{i, l(i)}}{2}} \right) \left(\psi_{l(i), \sigma_{t_2}}^{(t_2)}, \psi_{l(i), \sigma_{t_2}}^{(t_2)} \right)$$

that we can control by $\sum_{i=1}^{N(t_1)} \sum_{l(i)} \left(\left\| \psi_{l(i), \sigma_{t_2}}^{(t_2)} \right\|^2 \cdot C \cdot t_1^{-\epsilon} \cdot |\ln(\sigma_{t_2})| \right) \leq C \cdot t_1^{-\epsilon} \cdot |\ln(\sigma_{t_2})|$

analysis of A2

$$\left\| e^{iH_{t_2}} \sum_{i=1}^{N(t_1)} \sum_{l(i)} W_{\sigma_{t_2}}(\mathbf{v}_i, t_2) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_2)} e^{-iH_{\sigma_{t_2}} t_2} \left[\psi_{l(i), \sigma_{t_2}}^{(t_2)} - \psi_{i, \sigma_{t_2}}^{(t_1)} \right] \right\| = 0$$

because $\psi_{i, \sigma_{t_2}}^{(t_1)} = \int_{\Gamma_i} G(\mathbf{P}) \psi_{\mathbf{P}, \sigma_{t_2}} d^3 P = \sum_{l(i)} \int_{\Gamma_{l(i)}} G(\mathbf{P}) \psi_{\mathbf{P}, \sigma_{t_2}} d^3 P$ by definition.

analysis of B

In order to study the **B** term, I decompose it into five contributions and I estimate their norms.

$$\begin{aligned}
& \sum_i e^{iHt_2} W_{\sigma_{t_2}}(\mathbf{v}_i, t_2) W_{\sigma_{t_2}}^{b\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}, t_2)} W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}) e^{-iE^{\sigma_{t_2}} t_2} W_{\sigma_{t_2}}^b(\nabla E^{\sigma_{t_2}}) \psi_{i, \sigma_{t_2}}^{(t_1)} + \\
& - \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b\dagger}(\mathbf{v}_i) W_{\sigma_{t_1}}^b(\mathbf{v}_i) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} W_{\sigma_{t_1}}^{b\dagger}(\nabla E^{\sigma_{t_1}}) e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} = \\
& = \sum_i e^{iHt_2} W_{\sigma_{t_2}}(\mathbf{v}_i, t_2) W_{\sigma_{t_2}}^{b\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}, t_2)} e^{-iE^{\sigma_{t_2}} t_2} W_{\sigma_{t_2}}^b(\nabla E^{\sigma_{t_2}}) \psi_{i, \sigma_{t_2}}^{(t_1)} +
\end{aligned}$$

(BI)

$$\begin{aligned}
& - \sum_i e^{iHt_1} W_{\sigma_{t_2}}(\mathbf{v}_i, t_1) W_{\sigma_{t_2}}^{b^\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}, t_1)} e^{-iE^{\sigma_{t_2}} t_1} W_{\sigma_{t_2}}^b(\nabla E^{\sigma_{t_2}}) \psi_{i, \sigma_{t_2}}^{(t_1)} + \\
& + \sum_i e^{iHt_1} W_{\sigma_{t_2}}(\mathbf{v}_i, t_1) W_{\sigma_{t_2}}^{b^\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}, t_1)} e^{-iE^{\sigma_{t_2}} t_1} W_{\sigma_{t_2}}^b(\nabla E^{\sigma_{t_2}}) \psi_{i, \sigma_{t_2}}^{(t_1)} +
\end{aligned}
\tag{BII}$$

$$\begin{aligned}
& - \sum_i e^{iHt_1} W_{\sigma_{t_2}}(\mathbf{v}_i, t_1) W_{\sigma_{t_2}}^{b\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} + \\
& + \sum_i e^{iHt_1} W_{\sigma_{t_2}}(\mathbf{v}_i, t_1) W_{\sigma_{t_2}}^{b\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} +
\end{aligned}$$

BIII)

$$\begin{aligned}
& - \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} + \\
& + \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} +
\end{aligned}$$

BIV)

$$\begin{aligned}
& - \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b^\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_1}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} + \\
& + \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b^\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_1}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} +
\end{aligned}
\tag{BV}$$

$$-\sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b\dagger}(\mathbf{v}_i) W_{\sigma_{t_1}}^b(\mathbf{v}_i) W_{\sigma_{t_1}}^{b\dagger}(\nabla E^{\sigma_{t_1}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)}$$

control of BI)

The term **BI**) corresponds to

$$\sum_i \left\{ e^{iHt_2} W_{\sigma_{t_2}}(\mathbf{v}_i, t_2) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}, t_2)} e^{-iE^{\sigma_{t_2}} t_2} \psi_{i, \sigma_{t_2}}^{(t_1)} - e^{iHt_1} W_{\sigma_{t_2}}(\mathbf{v}_i, t_1) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}, t_1)} e^{-iE^{\sigma_{t_2}} t_1} \psi_{i, \sigma_{t_2}}^{(t_1)} \right\}$$

I estimate the contribution of the single cell by expressing the difference between times t_1 and t_2 as the integral of its derivative. Then I estimate the norm of the derivative:

$$\begin{aligned} & \frac{d}{ds} \left\{ e^{iHs} W_{\sigma_{t_2}}(\mathbf{v}_i, s) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}, s)} e^{-iE^{\sigma_{t_2}} s} \psi_{i, \sigma_{t_2}}^{(t_1)} \right\} ds = \\ & = \frac{d}{ds} \left\{ e^{iHs} W_{\sigma_{t_2}}(\mathbf{v}_i, s) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}, s)} e^{-iE^{\sigma_{t_2}} s} \psi_{i, \sigma_{t_2}}^{(t_1)} \right\} = \\ & = ie^{iHs} W_{\sigma_{t_2}}(\mathbf{v}_i, s) \left(\varphi_{\sigma_{t_2}, \mathbf{v}_i}(\mathbf{x}, s) + \frac{d\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}(\mathbf{P}), s)}{ds} \right) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}, s)} e^{-iE^{\sigma_{t_2}} s} \psi_{i, \sigma_{t_2}}^{(t_1)} + \end{aligned} \quad (28)$$

$$+ ie^{iHs} W_{\sigma_{t_2}}(\mathbf{v}_i, s) (H - H_{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}, s)} e^{-iE^{\sigma_{t_2}} s} \psi_{i, \sigma_{t_2}}^{(t_1)} \quad (29)$$

The formal steps are well defined as $\psi_{i, \sigma_{t_2}}^{(t_1)} \in D(H_{\sigma_{t_2}}) \equiv D(H)$.

estimate of (28)

Analogously to the proof of **II**) in paragraph 4.1, I decompose $\varphi_{\sigma_{t_2}, \mathbf{v}_i}(\mathbf{x}, s)$ in $\varphi_{\sigma_{t_2}, \mathbf{v}_i}^1(\mathbf{x}, s) + \varphi_{\sigma_{t_2}, \mathbf{v}_i}^2(\mathbf{x}, s)$, having defined:

$$\begin{aligned} \varphi_{\sigma_{t_2}, \mathbf{v}_i}^1(\mathbf{x}, s) &\equiv g^2 \int_{\sigma_{t_2}}^{s^{-\frac{39}{40}}} \int \frac{\cos(\mathbf{k} \cdot \mathbf{x} - |\mathbf{k}|s)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| \\ \varphi_{\sigma_{t_2}, \mathbf{v}_i}^2(\mathbf{x}, s) &\equiv g^2 \int_{s^{-\frac{39}{40}}}^{\kappa_1} \int \frac{\cos(\mathbf{k} \cdot \mathbf{x} - |\mathbf{k}|s)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| \end{aligned}$$

by steps analogous to the ones used in **II**) of paragraph 4.1, we obtain the following estimate of the norm of the expression (28):

$$C \cdot s^{-1} \cdot s^{-\frac{1}{112}} \cdot s^{2\delta} \cdot t_1^{-\frac{3\epsilon}{2}} \cdot (\ln \sigma_{t_2})^2$$

estimate of (29)

The norm of the vector

$$(H - H_{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}, t_2})} \psi_{i, \sigma_{t_2}}^{(t_1)} = e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}, t_2})} g \int_0^{\sigma_{t_2}} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) \frac{d^3 k}{\sqrt{2|\mathbf{k}|}} \psi_{i, \sigma_{t_2}}^{(t_1)}$$

can be estimated in the following way:

$$\begin{aligned} \left\| g \int_0^{\sigma_{t_2}} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) \frac{d^3 k}{\sqrt{2|\mathbf{k}|}} \psi_{i, \sigma_{t_2}}^{(t_1)} \right\| &\leq \left\| g \int_0^{\sigma_{t_2}} b(\mathbf{k}) \frac{d^3 k}{\sqrt{2|\mathbf{k}|}} \psi_{i, \sigma_{t_2}}^{(t_1)} \right\| + \left\| g \int_0^{\sigma_{t_2}} b^\dagger(\mathbf{k}) \frac{d^3 k}{\sqrt{2|\mathbf{k}|}} \psi_{i, \sigma_{t_2}}^{(t_1)} \right\| \leq \\ &\leq \left(g^2 \int_0^{\sigma_{t_2}} \frac{d^3 k}{2|\mathbf{k}|} \right)^{\frac{1}{2}} \cdot \left\| \psi_{i, \sigma_{t_2}}^{(t_1)} \right\| \leq C \cdot (\sigma_{t_2}) \cdot t_1^{-\frac{3\epsilon}{2}} \end{aligned}$$

(I used the fact that $b(\mathbf{k}) \psi_{i, \sigma_{t_2}}^{(t_1)} = 0$ for $\mathbf{k} : |\mathbf{k}| \leq \sigma_{t_2}$)

Therefore the norm of the term **BI**) is bounded by a quantity of order

$$t_1^{-\frac{1}{112}} \cdot t_1^{2\delta + \frac{3\epsilon}{2}} \cdot (\ln \sigma_{t_2})^2 + t_2 \cdot (\sigma_{t_2}) \cdot t_1^{\frac{3\epsilon}{2}}$$

(remember that the constraint $2\delta + 3\epsilon < \frac{1}{112}$ is assumed)

control of BII)

The contribution of a single cell vanishes with a rate related to the decrease of the cut-off. In fact

$$\begin{aligned} &\left\| e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}, t_1})} e^{-iE^{\sigma_{t_2}, t_1}} W_{\sigma_{t_2}}^b (\nabla E^{\sigma_{t_2}}) \psi_{i, \sigma_{t_2}}^{(t_1)} - e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}, t_1})} e^{-iE^{\sigma_{t_1}, t_1}} W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \right\| \leq \\ &\leq \left\| e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}, t_1})} e^{-iE^{\sigma_{t_2}, t_1}} W_{\sigma_{t_2}}^b (\nabla E^{\sigma_{t_2}}) \psi_{i, \sigma_{t_2}}^{(t_1)} - e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}, t_1})} e^{-iE^{\sigma_{t_2}, t_1}} W_{\sigma_{t_2}}^b (\nabla E^{\sigma_{t_2}}) \psi_{i, \sigma_{t_2}}^{(t_1)} \right\| + \\ &+ \left\| e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}, t_1})} e^{-iE^{\sigma_{t_2}, t_1}} W_{\sigma_{t_2}}^b (\nabla E^{\sigma_{t_2}}) \psi_{i, \sigma_{t_2}}^{(t_1)} - e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}, t_1})} e^{-iE^{\sigma_{t_1}, t_1}} W_{\sigma_{t_2}}^b (\nabla E^{\sigma_{t_2}}) \psi_{i, \sigma_{t_2}}^{(t_1)} \right\| + \\ &+ \left\| e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}, t_1})} e^{-iE^{\sigma_{t_1}, t_1}} W_{\sigma_{t_2}}^b (\nabla E^{\sigma_{t_2}}) \psi_{i, \sigma_{t_2}}^{(t_1)} - e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}, t_1})} e^{-iE^{\sigma_{t_1}, t_1}} W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \right\| \end{aligned}$$

first term of the sum

The module of $e^{i\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}, t_1})} - e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}, t_1})}$ can be estimated in terms of the module of the difference between the arguments of the exponentials:

$$\gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_1) - \gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}(\mathbf{P}), t_1) =$$

$$\begin{aligned}
&= - \int_1^{t_1} \left\{ g^2 \int_{\sigma_{t_1}}^{\sigma_{\tau}^L} \int \frac{\cos(\mathbf{k} \cdot \nabla E^{\sigma_{t_2}} \tau - |\mathbf{k}| \tau) - \cos(\mathbf{k} \cdot \nabla E^{\sigma_{t_1}} \tau - |\mathbf{k}| \tau)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| \right\} d\tau + \\
&- \int_1^{t_1} \left\{ g^2 \int_{\sigma_{t_2}}^{\sigma_{t_1}} \int \frac{\cos(\mathbf{k} \cdot \nabla E^{\sigma_{t_2}} \tau - |\mathbf{k}| \tau)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| \right\} d\tau = \\
&= 2 \int_1^{t_1} \left\{ g^2 \int_{\sigma_{t_1}}^{\sigma_{\tau}^L} \int \frac{\sin\left(\frac{\mathbf{k} \cdot (\nabla E^{\sigma_{t_2}} - \nabla E^{\sigma_{t_1}}) \tau}{2}\right) \sin\left(\frac{\mathbf{k} \cdot (\nabla E^{\sigma_{t_2}} + \nabla E^{\sigma_{t_1}}) \tau - 2|\mathbf{k}| \tau}{2}\right)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| \right\} d\tau + \\
&- \int_1^{t_1} \left\{ g^2 \int_{\sigma_{t_2}}^{\sigma_{t_1}} \int \frac{\cos(\mathbf{k} \cdot \nabla E^{\sigma_{t_2}} \tau - |\mathbf{k}| \tau)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| \right\} d\tau
\end{aligned}$$

then

$$\begin{aligned}
&\left| \gamma_{\sigma_{t_2}}(\mathbf{v}_i, \nabla E^{\sigma_{t_2}}(\mathbf{P}), t_1) - \gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}(\mathbf{P}), t_1) \right| \leq \\
&\leq C \cdot |\nabla E^{\sigma_{t_2}}(\mathbf{P}) - \nabla E^{\sigma_{t_1}}(\mathbf{P})| \cdot \int_1^{t_1} (\sigma_{\tau}^L)^2 \cdot \tau d\tau + C \cdot t_1 \cdot \sigma_{t_1} \leq \\
&\leq C \cdot (\sigma_{t_1})^{\frac{1}{4}} \int_1^{t_1} \tau^{-\frac{19}{20}} d\tau + C \cdot t_1 \cdot \sigma_{t_1}
\end{aligned}$$

(the bound of the module $\nabla E_{\mathbf{P}}^{\sigma_{t_2}} - \nabla E_{\mathbf{P}}^{\sigma_{t_1}}$ comes substantially from lemma A2, $\sigma_{t_1} = t_1^{-\beta}$ where $\beta > 1$, by hypothesis).

Then we have a bound by a quantity of order $t_1 \cdot (\sigma_{t_1})^{\frac{1}{4}}$.

second term of the sum

Taking in account the iterative procedure, the module $|e^{-iE^{\sigma_{t_2}} t_1} - e^{-iE^{\sigma_{t_1}} t_1}|$ is bounded by a quantity of order $(\sigma_{t_1}) \cdot t_1$

third term of the sum

$$\begin{aligned}
&\left\| e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_2}}^b(\nabla E^{\sigma_{t_2}}) \psi_{i, \sigma_{t_2}}^{(t_1)} - e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \right\| = \\
&= \left\| \int_{\Gamma_i} G(\mathbf{P}) W_{\sigma_{t_2}}^b(\nabla E^{\sigma_{t_2}}(\mathbf{P})) \psi_{\mathbf{P}, \sigma_{t_2}} d^3 P - \int_{\Gamma_i} G(\mathbf{P}) W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{\mathbf{P}, \sigma_{t_1}} d^3 P \right\| \leq \\
&\leq C \cdot (\sigma_{t_1})^{\frac{1}{8}} \cdot t_1^{-\frac{3\epsilon}{2}} \text{ (see theorem 3.2).}
\end{aligned}$$

Therefore the norm of **BII**) is surely bounded by:

$$C \cdot t_1^{\frac{3\epsilon}{2}+1} \cdot (\sigma_{t_1})^{\frac{1}{8}}$$

control of BIII)

$$\begin{aligned} & \sum_i e^{iHt_1} W_{\sigma_{t_2}}(\mathbf{v}_i, t_1) W_{\sigma_{t_2}}^{b^\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} + \\ & - \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b^\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \\ & \text{Having defined } \varphi_i = W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \\ & \text{the III) is written as:} \end{aligned}$$

$$\sum_i e^{iHt_1} \left(W_{\sigma_{t_2}}(\mathbf{v}_i, t_1) W_{\sigma_{t_2}}^{b^\dagger}(\mathbf{v}_i) - W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b^\dagger}(\mathbf{v}_i) \right) \varphi_i$$

I restrict the analysis to the single cell. For this purpose I examine

$$\begin{aligned} & W_{\sigma_{t_2}}(\mathbf{v}_i, t_1) W_{\sigma_{t_2}}^{b^\dagger}(\mathbf{v}_i) - W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b^\dagger}(\mathbf{v}_i) = \\ & = W_{\kappa_1}^{b^\dagger}(\mathbf{v}_i) e^{-g \int_{\sigma_{t_2}}^{\kappa_1} \frac{a(\mathbf{k})(e^{i|\mathbf{k}|t_1 - e^{i\mathbf{k} \cdot \mathbf{x}}}) - a^\dagger(\mathbf{k})(e^{-i|\mathbf{k}|t_1 - e^{-i\mathbf{k} \cdot \mathbf{x}}})}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} - W_{\kappa_1}^{b^\dagger}(\mathbf{v}_i) e^{-g \int_{\sigma_{t_1}}^{\kappa_1} \frac{a(\mathbf{k})(e^{i|\mathbf{k}|t_1 - e^{i\mathbf{k} \cdot \mathbf{x}}}) - c.c.}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} \\ & = W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b^\dagger}(\mathbf{v}_i) \left(e^{-g \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{a(\mathbf{k})(e^{i|\mathbf{k}|t_1 - e^{i\mathbf{k} \cdot \mathbf{x}}}) - a^\dagger(\mathbf{k})(e^{-i|\mathbf{k}|t_1 - e^{-i\mathbf{k} \cdot \mathbf{x}}})}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} - 1 \right) \end{aligned}$$

On the vectors belonging to the domain of the generator of the exponential we have

$$\begin{aligned} & e^{-g \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{a(\mathbf{k})(e^{i|\mathbf{k}|t_1 - e^{i\mathbf{k} \cdot \mathbf{x}}}) - a^\dagger(\mathbf{k})(e^{-i|\mathbf{k}|t_1 - e^{-i\mathbf{k} \cdot \mathbf{x}}})}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} - 1 = \\ & = - \int_0^1 e^{-g\lambda \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{a(\mathbf{k})(e^{i|\mathbf{k}|t_1 - e^{i\mathbf{k} \cdot \mathbf{x}}}) - a^\dagger(\mathbf{k})(e^{-i|\mathbf{k}|t_1 - e^{-i\mathbf{k} \cdot \mathbf{x}}})}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} d\lambda \cdot g \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{a(\mathbf{k})(e^{i|\mathbf{k}|t_1 - e^{i\mathbf{k} \cdot \mathbf{x}}}) - c.c.}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} \end{aligned}$$

Since the vector φ_i belongs to the domain of the generator, the following bound holds

$$\begin{aligned}
& \left\| g \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{a(\mathbf{k})(e^{i|\mathbf{k}|t_1} - e^{i\mathbf{k} \cdot \mathbf{x}}) - c.c.}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2|\mathbf{k}|}} \varphi_i \right\| \leq \left\| g \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{a(\mathbf{k})(e^{i|\mathbf{k}|t_1} - e^{i\mathbf{k} \cdot \mathbf{x}})}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2|\mathbf{k}|}} \varphi_i \right\| + \\
& + \left\| g \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{a^\dagger(\mathbf{k})(e^{-i|\mathbf{k}|t_1} - e^{-i\mathbf{k} \cdot \mathbf{x}})}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2|\mathbf{k}|}} \varphi_i \right\| \leq \\
& \leq 2 \left\| g \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{b(\mathbf{k})(e^{i|\mathbf{k}|t_1} - e^{i\mathbf{k} \cdot \mathbf{x}} - 1)}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2|\mathbf{k}|}} \varphi_i \right\| + \left(\varphi_i, g^2 \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{|e^{i|\mathbf{k}|t_1} - e^{i\mathbf{k} \cdot \mathbf{x}}|^2}{2|\mathbf{k}|^3(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)^2} d^3 k \varphi_i \right)^{\frac{1}{2}}
\end{aligned}$$

Analysis of $\left\| g \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{b(\mathbf{k})(e^{i|\mathbf{k}|t_1} - e^{i\mathbf{k} \cdot \mathbf{x}} - 1)}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2|\mathbf{k}|}} \varphi_i \right\|.$

1) the expression $b(\mathbf{k}) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)}$ is a well defined vector in $\mathbb{H}$ and it is strongly continuous in $\mathbf{k}$. Therefore:

$$\begin{aligned}
& \left\| \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{b(\mathbf{k})(e^{i|\mathbf{k}|t_1} - e^{i\mathbf{k} \cdot \mathbf{x}} - 1)}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2|\mathbf{k}|}} W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \right\| \leq \\
& \leq \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{|1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i|^{-1}}{|\mathbf{k}|^{\frac{3}{2}} \sqrt{2}} \left\| b(\mathbf{k}) (e^{i|\mathbf{k}|t_1} e^{-i\mathbf{k} \cdot \mathbf{x}} - 1) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \right\| d^3 k
\end{aligned}$$

2) estimate of

$$\begin{aligned}
& \left\| b(\mathbf{k}) (e^{i|\mathbf{k}|t_1} e^{-i\mathbf{k} \cdot \mathbf{x}} - 1) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \right\| \\
& \Leftrightarrow
\end{aligned}$$

1) In distributional sense, the following equality is valid:

$$b(\mathbf{k}) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_2}}(\mathbf{P})) = W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_2}}(\mathbf{P}))(b(\mathbf{k}) + f(\mathbf{k}, \mathbf{v}_i, \mathbf{P}))$$

$$\begin{aligned}
& \text{where } f(\mathbf{k}, \mathbf{v}_i, \mathbf{P}) = -\frac{g\chi_{\sigma_{t_2}}^\kappa(\mathbf{k})}{|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} + \frac{g\chi_{\sigma_{t_2}}^\kappa(\mathbf{k})}{|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \nabla E^{\sigma_{t_2}}(\mathbf{P}))} \\
& (\chi_{\sigma_{t_2}}^\kappa \text{ characteristic function of the set } \{\mathbf{k} : \sigma_{t_2} \leq |\mathbf{k}| \leq \kappa\});
\end{aligned}$$

$$\text{moreover, being } |\mathbf{k}| \leq \sigma_{t_1}, \quad b(\mathbf{k}) W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} = 0$$

Therefore for $\mathbf{k}$ such that $|\mathbf{k}| \leq \sigma_{t_1}$

$$b(\mathbf{k}) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b^\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} =$$

$$= f(\mathbf{k}, \mathbf{v}_i, \mathbf{P}) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)}.$$

The strong continuity in $\mathbf{k}$, $\sigma_{t_2} \leq |\mathbf{k}| \leq \sigma_{t_1}$, comes from the continuity of the function $f(\mathbf{k}, \mathbf{v}_i, \mathbf{P})$.

2) Estimate of

$$\left\| b(\mathbf{k}) \left(e^{i|\mathbf{k}|t_1} e^{-i\mathbf{k} \cdot \mathbf{x}} - 1 \right) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \right\|$$

Starting from the identity:

$$\begin{aligned} & e^{i|\mathbf{k}|t_1} W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E_{\mathbf{P}+\mathbf{k}}^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E_{\mathbf{P}+\mathbf{k}}^{\sigma_{t_1}}, t_1)} e^{-iE_{\mathbf{P}+\mathbf{k}}^{\sigma_{t_1}} t_1} f(\mathbf{k}, \mathbf{v}_i, \mathbf{P} + \mathbf{k}) e^{-i\mathbf{k} \cdot \mathbf{x}} W_{\sigma_{t_1}}^b(\nabla E_{\mathbf{P}}^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} + \\ & - W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}(\mathbf{P})) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}(\mathbf{P}), t_1)} e^{-iE_{\mathbf{P}}^{\sigma_{t_1}} t_1} f(\mathbf{k}, \mathbf{v}_i, \mathbf{P}) W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{i, \sigma_{t_1}}^{(t_1)} = \\ & = e^{i|\mathbf{k}|t_1} W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E_{\mathbf{P}+\mathbf{k}}^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E_{\mathbf{P}+\mathbf{k}}^{\sigma_{t_1}}, t_1)} e^{-iE_{\mathbf{P}+\mathbf{k}}^{\sigma_{t_1}} t_1} f(\mathbf{k}, \mathbf{v}_i, \mathbf{P} + \mathbf{k}) e^{-i\mathbf{k} \cdot \mathbf{x}} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{i, \sigma_{t_1}}^{(t_1)} + \\ & - W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}(\mathbf{P})) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}(\mathbf{P}), t_1)} e^{-iE^{\sigma_{t_1}}(\mathbf{P}) t_1} f(\mathbf{k}, \mathbf{v}_i, \mathbf{P}) e^{-i\mathbf{k} \cdot \mathbf{x}} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} + \\ & + W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}(\mathbf{P})) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}(\mathbf{P}), t_1)} e^{-iE^{\sigma_{t_1}}(\mathbf{P}) t_1} f(\mathbf{k}, \mathbf{v}_i, \mathbf{P}) e^{-i\mathbf{k} \cdot \mathbf{x}} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} + \\ & - W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}(\mathbf{P})) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}}(\mathbf{P}) t_1} f(\mathbf{k}, \mathbf{v}_i, \mathbf{P}) W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \end{aligned} \quad (31)$$

I estimate the expression (30) and the expression (31).

Estimate of (30)

Considering that:

$$\begin{aligned} 1. \quad & \left\| \left(W_{\sigma_{t_2}}^{b\dagger}(\nabla E_{\mathbf{P}+\mathbf{k}}^{\sigma_{t_2}}) - W_{\sigma_{t_2}}^{b\dagger}(\nabla E_{\mathbf{P}}^{\sigma_{t_2}}) \right) f(\mathbf{k}, \mathbf{v}_i, \mathbf{P} + \mathbf{k}) e^{-i\mathbf{k} \cdot \mathbf{x}} W_{\sigma_{t_1}}^b(\nabla E_{\mathbf{P}}^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \right\| \leq \\ & \leq C \cdot (\ln(\sigma_{t_2}) \cdot \ln(\sigma_{t_1}))^{\frac{1}{2}} \cdot t_1^{-\frac{3\epsilon}{2}} \cdot |\mathbf{k}|^{\frac{1}{16}} \cdot |\mathbf{k}|^{-\frac{3}{2}} \text{ (it is proved starting from an estimate analogous to the (b4) in lemma B2 and from the fact that } |\mathbf{k}| \leq \sigma_{t_1} \ll 1); \end{aligned}$$

$$2. \quad \left| e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}(\mathbf{P}+\mathbf{k}), t_1)} - e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}(\mathbf{P}), t_1)} \right| \leq C \cdot |\mathbf{k}|^{\frac{1}{16}} \cdot t_1^{\frac{1}{20}}$$

by comparing the arguments of the exponentials as in the discussion of **BII**);

$$3. \left| e^{i|\mathbf{k}|t_1} - 1 \right| < 2t_1 \cdot |\mathbf{k}|$$

$$4. |f(\mathbf{k}, \mathbf{v}_i, \mathbf{P}) - f(\mathbf{k}, \mathbf{v}_i, \mathbf{P} + \mathbf{k})| \leq C \cdot \frac{|\mathbf{k}|^{\frac{1}{16}}}{|\mathbf{k}|^{\frac{3}{2}}}$$

$$5. \left| e^{-iE^{\sigma_{t_1}}(\mathbf{P}+\mathbf{k})t_1} - e^{-iE^{\sigma_{t_1}}(\mathbf{P})t_1} \right| \leq C \cdot |\mathbf{k}|^{\frac{1}{16}} \cdot t_1$$

being $|\mathbf{k}| \leq \sigma_{t_1} \ll 1$, one can conclude that (30) is bounded by

$$C \cdot |\mathbf{k}|^{\frac{1}{16}} \cdot |\mathbf{k}|^{-\frac{3}{2}} \cdot t_1 \cdot t_1^{-\frac{3\epsilon}{2}} \cdot |\ln \sigma_{t_2}|$$

Estimate of (31)

The norm of the expression (31) is equal to

$$\begin{aligned} & \left\| e^{-i\mathbf{k} \cdot \mathbf{x}} W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} - W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \right\| = \\ & = \left\| e^{-i\mathbf{k} \cdot \mathbf{x}} \int_{\Gamma_i} G(\mathbf{P}) W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{\mathbf{P}, \sigma_{t_1}} d^3 P - \int_{\Gamma_i} G(\mathbf{P}) W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{\mathbf{P}, \sigma_{t_1}} d^3 P \right\| \end{aligned}$$

I observe that $e^{-i\mathbf{k} \cdot \mathbf{x}} W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{\mathbf{P}, \sigma_{t_1}} \in H_{\mathbf{P}-\mathbf{k}}$ and that the following equality holds in F^b :

$$I_{\mathbf{P}-\mathbf{k}} \left(e^{-i\mathbf{k} \cdot \mathbf{x}} W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{\mathbf{P}, \sigma_{t_1}} \right) = I_{\mathbf{P}} \left(W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{\mathbf{P}, \sigma_{t_1}} \right)$$

Then

$$\begin{aligned} & e^{-i\mathbf{k} \cdot \mathbf{x}} \int_{\Gamma_i} G(\mathbf{P}) W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{\mathbf{P}, \sigma_{t_1}} d^3 P - \int_{\Gamma_i} G(\mathbf{P}) W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{\mathbf{P}, \sigma_{t_1}} d^3 P = \\ & = \int_{\Gamma_i} G(\mathbf{P}) e^{-i\mathbf{k} \cdot \mathbf{x}} \left\{ W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{\mathbf{P}, \sigma_{t_1}} \right\} d^3 P - \int_{\Gamma_i} G(\mathbf{P}) W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}(\mathbf{P}-\mathbf{k})) \psi_{\mathbf{P}-\mathbf{k}, \sigma_{t_1}} d^3 P + \\ & \quad (32.1) \\ & + \int_{\Gamma_i} G(\mathbf{P}) W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}(\mathbf{P}-\mathbf{k})) \psi_{\mathbf{P}-\mathbf{k}, \sigma_{t_1}} d^3 P - \int_{\Gamma_i} G(\mathbf{P}-\mathbf{k}) W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}(\mathbf{P}-\mathbf{k})) \psi_{\mathbf{P}-\mathbf{k}, \sigma_{t_1}} d^3 P + \end{aligned}$$

(32.2)

$$+ \int_{\Gamma_i} G(\mathbf{P} - \mathbf{k}) W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}(\mathbf{P} - \mathbf{k})) \psi_{\mathbf{P}-\mathbf{k}, \sigma_{t_1}} d^3 P - \int_{\Gamma_i} G(\mathbf{P}) W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{\mathbf{P}, \sigma_{t_1}} d^3 P \quad (32.3)$$

The term (32.1) can be estimated starting from

$$\left\| I_{\mathbf{P}-\mathbf{k}} \left(e^{-i\mathbf{k} \cdot \mathbf{x}} \left\{ W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}(\mathbf{P})) \psi_{\mathbf{P}, \sigma_{t_1}} \right\} \right) - I_{\mathbf{P}-\mathbf{k}} \left(W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}(\mathbf{P} - \mathbf{k})) \psi_{\mathbf{P}-\mathbf{k}, \sigma_{t_1}} \right) \right\|_F$$

and then it is norm bounded by $C \cdot |\mathbf{k}|^{\frac{1}{32}} \cdot t_1^{-\frac{3\epsilon}{2}}$, for theorem 3.4.

Being $G \in C_0^1(R^3 \setminus 0)$, the norm of the term (32.2) is bounded by $C \cdot |\mathbf{k}| \cdot t_1^{-\frac{3\epsilon}{2}}$.

After having estimated a volume difference, the norm of the term (32.3) is bounded by a quantity of order $|\mathbf{k}|^{\frac{1}{2}} \cdot t_1^{-\epsilon}$.

In conclusion the norm of the (31) is bounded by $C \cdot |\mathbf{k}|^{\frac{1}{32}} \cdot t_1^{-\epsilon}$ ($|\mathbf{k}| \ll 1$).

$$\begin{aligned} \text{Then } \left\| g \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{b(\mathbf{k})(e^{i|\mathbf{k}|t_1 - i\mathbf{k} \cdot \mathbf{x}} - 1)}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|} \varphi_i \right\| &\leq C \cdot t_1 \cdot t_1^{-\epsilon} \cdot |\ln \sigma_{t_2}| \cdot \int_0^{\sigma_{t_1}} \frac{|\mathbf{k}|^{\frac{1}{32}}}{|\mathbf{k}|} d|\mathbf{k}| = \\ &= C \cdot t_1 \cdot t_1^{-\epsilon} \cdot |\ln \sigma_{t_2}| \cdot (\sigma_{t_1})^{\frac{1}{32}} \end{aligned}$$

Note

By the same steps, one obtains that

$$\left\| \left(e^{i|\mathbf{k}|t_1} e^{-i\mathbf{k} \cdot \mathbf{x}} - 1 \right) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \right\|$$

is bounded by a quantity of order $|\mathbf{k}|^{\frac{1}{32}} \cdot t_1 \cdot t_1^{-\epsilon} \cdot |\ln \sigma_{t_2}|$.

$$\textbf{Analysis of } \left(\varphi_i, g^2 \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{|e^{i|\mathbf{k}|t_1} - e^{i\mathbf{k} \cdot \mathbf{x}}|^2}{2|\mathbf{k}|^3 (1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)^2} d^3 k \varphi_i \right)^{\frac{1}{2}}$$

Note that $|e^{i|\mathbf{k}|t_1} - e^{i\mathbf{k} \cdot \mathbf{x}}|^2 = 2 - e^{-i|\mathbf{k}|t_1} e^{i\mathbf{k} \cdot \mathbf{x}} - e^{i|\mathbf{k}|t_1} e^{-i\mathbf{k} \cdot \mathbf{x}}$. Taking into account the result of the previous note, we have that:

$$\left(\varphi_i, g^2 \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{|e^{i|\mathbf{k}|t_1 - e^{i\mathbf{k} \cdot \mathbf{x}}}|^2}{2|\mathbf{k}|^3(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)^2} d^3k \varphi_i \right)^{\frac{1}{2}} \leq C \cdot t_1^{\frac{1}{2}} \cdot t_1^{-\frac{5\epsilon}{4}} \cdot (|\ln \sigma_{t_2}|)^{\frac{1}{2}} \cdot (\sigma_{t_1})^{\frac{1}{64}}$$

Therefore the norm of the term **BIII**) is surely bounded by a quantity of order:

$$t_1 \cdot t_1^{2\epsilon} \cdot |\ln \sigma_{t_2}| \cdot (\sigma_{t_1})^{\frac{1}{64}}$$

control of BIV)

$$\begin{aligned} & \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} + \\ & - \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_1}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \end{aligned}$$

For a single cell

$$\left\| \left(W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_2}}) - W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_1}}) \right) W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} \right\| \leq C \cdot (\sigma_{t_1})^{\frac{1}{4}} \cdot |\ln \sigma_{t_2}| \cdot t_1^{-\frac{3\epsilon}{2}};$$

therefore the norm of **BIV**) is bounded by $C \cdot t_1^{\frac{3\epsilon}{2}} \cdot (\sigma_{t_1})^{\frac{1}{4}} \cdot |\ln \sigma_{t_2}|$

control of BV)

$$\begin{aligned} & \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b\dagger}(\mathbf{v}_i) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_1}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} + \\ & - \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_1}}^{b\dagger}(\mathbf{v}_i) W_{\sigma_{t_1}}^b(\mathbf{v}_i) W_{\sigma_{t_1}}^{b\dagger}(\nabla E^{\sigma_{t_1}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} W_{\sigma_{t_1}}^b(\nabla E^{\sigma_{t_1}}) \psi_{i, \sigma_{t_1}}^{(t_1)} = \\ & = \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_1}}) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} \psi_{i, \sigma_{t_1}}^{(t_1)} + \\ & - \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} \psi_{i, \sigma_{t_1}}^{(t_1)} = \\ & = \sum_i e^{iHt_1} W_{\sigma_{t_1}}(\mathbf{v}_i, t_1) \left(W_{\sigma_{t_2}}^b(\mathbf{v}_i) W_{\sigma_{t_2}}^{b\dagger}(\nabla E^{\sigma_{t_1}}) - 1 \right) e^{i\gamma_{\sigma_{t_1}}(\mathbf{v}_i, \nabla E^{\sigma_{t_1}}, t_1)} e^{-iE^{\sigma_{t_1}} t_1} \psi_{i, \sigma_{t_1}}^{(t_1)} \end{aligned}$$

with the definitions

- $$W^b \big|_{\sigma_{t_2}}^{\sigma_{t_1}} (\mathbf{v}_i) \equiv W_{\sigma_{t_1}}^{b^\dagger} (\mathbf{v}_i) W_{\sigma_{t_2}}^b (\mathbf{v}_i) = e^{g \int_{\sigma_{t_1}}^{\kappa} \frac{b(\mathbf{k}) - b^\dagger(\mathbf{k})}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} e^{-g \int_{\sigma_{t_2}}^{\kappa} \frac{b(\mathbf{k}) - b^\dagger(\mathbf{k})}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}} =$$

$$= e^{-g \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{b(\mathbf{k}) - b^\dagger(\mathbf{k})}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}}$$
- $$W^{b^\dagger} \big|_{\sigma_{t_1}}^{\sigma_{t_2}} (\nabla E^{\sigma_{t_1}}) \equiv W_{\sigma_{t_2}}^{b^\dagger} (\nabla E^{\sigma_{t_1}}) W_{\sigma_{t_1}}^b (\nabla E^{\sigma_{t_1}}) = e^{g \int_{\sigma_{t_2}}^{\sigma_{t_1}} \frac{b(\mathbf{k}) - b^\dagger(\mathbf{k})}{|\mathbf{k}|(1 - \widehat{\mathbf{k}} \cdot \nabla E^{\sigma_{t_1}})} \frac{d^3 k}{\sqrt{2}|\mathbf{k}|}}$$

The discussion of this contribution requires the study of the squared norm and the control of mixed terms in the scalar product. In order to do it I will verify that the sum of the mixed terms vanishes and I will determine the rate. Then I will examine the diagonals terms.

mixed terms

As in par.4.1 “**Control of the norm..**”, I will consider the generic i-j term. I observe that it is possible to reply the same procedure since the operators, obtained from derivation with respect to the parameter λ , commute with $\left(W^b \big|_{\sigma_{t_2}}^{\sigma_{t_1}} (\mathbf{v}_i) W^{b^\dagger} \big|_{\sigma_{t_2}}^{\sigma_{t_1}} (\nabla E^{\sigma_{t_1}}) - 1 \right)$. Then the result is analogous: the sum of the modules of the mixed terms is bounded by $C \cdot t_1^{-\frac{1}{112}} \cdot t_1^{2\delta+3\epsilon} \cdot (\ln \sigma_{t_1})^2$.

diagonal terms

Considering that:

- the norm $\left\| \left(W^b \big|_{\sigma_{t_2}}^{\sigma_{t_1}} (\mathbf{v}_i) W^b \big|_{\sigma_{t_2}}^{\sigma_{t_1}} (\nabla E^{\sigma_{t_1}} (\mathbf{P})) - 1 \right) \psi_{i, \sigma_{t_1}}^{(t_1)} \right\|$ can be estimated by a quantity of order $\sup_{\mathbf{P} \in \Gamma_i} |\mathbf{v}_i - \nabla E^{\sigma_{t_1}} (\mathbf{P})| \cdot (|\ln \sigma_{t_2}|)^{\frac{1}{2}} \cdot t_1^{-\frac{3\epsilon}{2}}$
- $\sup_{\mathbf{P} \in \Gamma_i} |\mathbf{v}_i - \nabla E^{\sigma_{t_1}} (\mathbf{P})| \leq \sup_{\mathbf{P} \in \Gamma_i} \left| \nabla E^{\sigma_{t_1}} (\mathbf{P}) - \nabla E^{\sigma_{t_1}} (\overline{\mathbf{P}}_i) \right| \leq$
 $\leq C \cdot \sup_{\mathbf{P} \in \Gamma_i} \left| \mathbf{P} - \overline{\mathbf{P}}_i \right|^{\frac{1}{16}} \leq C \cdot t_1^{-\frac{\epsilon}{16}}$ (for the last step, see lemma 3.3)

the sum of the diagonal terms gives a contribution bounded by

$$C \cdot t_1^{-\frac{\epsilon}{32}} \cdot (\ln(\sigma_{t_2}))^{\frac{1}{4}}$$

Therefore it follows that the norm of the term **BV**) is bounded by

$$C \cdot \max \left(t_1^{-\frac{\epsilon}{32}}, \left(t_1^{-\frac{1}{112}} \cdot t_1^{2\delta+3\epsilon} \right)^{\frac{1}{2}} \right) \cdot |\ln \sigma_{t_2}|$$

Theorem 4.1

The vector $\psi_G(t)$ converges strongly, for $t \rightarrow +\infty$, with an error of order $\frac{1}{t^\rho}$ where $\rho > 0$ is a properly small coefficient.

Proof

We look at the bounds obtained for the norms of A1, A2 and B

$$\text{A1)} \ C \cdot \left\{ \left(t_1^{-\epsilon} \cdot |\ln \sigma_{t_2}| \right) + t_2^{-\frac{1}{112}} \cdot t_2^{2\delta+3\epsilon} \cdot (\ln \sigma_{t_2})^2 \right\}^{\frac{1}{2}}$$

$$\text{A2)} \ 0$$

$$\text{BI)} \ C \cdot t_1^{-\frac{1}{112}} \cdot t_1^{2\delta+\frac{3\epsilon}{2}} \cdot (\ln \sigma_{t_2})^2 + C \cdot t_2 \cdot (\sigma_{t_2}) \cdot t_1^{\frac{3\epsilon}{2}}$$

$$\text{BII)} \ C \cdot t_1^{\frac{3\epsilon}{2}} \cdot t_1 \cdot (\sigma_{t_1})^{\frac{1}{8}}$$

$$\text{BIII)} \ C \cdot t_1 \cdot t_1^{2\epsilon} \cdot |\ln \sigma_{t_2}| \cdot (\sigma_{t_1})^{\frac{1}{64}}$$

$$\text{BIV)} \ C \cdot t_1^{\frac{3\epsilon}{2}} \cdot (\sigma_{t_1})^{\frac{1}{4}} \cdot |\ln \sigma_{t_2}|$$

$$\text{BV)} \ C \cdot \max \left(t_1^{-\frac{\epsilon}{32}}, \left(t_1^{-\frac{1}{112}} \cdot t_1^{2\delta+3\epsilon} \right)^{\frac{1}{2}} \right) \cdot |\ln \sigma_{t_2}|$$

where we have tuned the time scales related to ϵ e δ in accordance to the constraints

- $24\epsilon < \delta$.
- $2\delta + 3\epsilon < \frac{1}{112}$

and we have chosen, for example, $\sigma_t = t^{-128}$. Thus we can observe that the time t_2 appears in the numerators only in $\ln(\sigma_{t_2})$ and that we can estimate

$$\|\psi_G(t_2) - \psi_G(t_1)\| < \frac{C}{M} \cdot \left(\frac{\ln(t_2)}{t_1^{3\rho}} \right)^2$$

where $\rho > 0$ e $\frac{C}{M} > 0$ are independent of t_1 e t_2 (for $t_2 \geq t_1 > \bar{t} \gg 1$).

Now, let us consider the sequence $t_1, t_1^2, t_1^3, \dots, t_1^n, \dots$ and put $t_1^n \leq t_2 < t_1^{n+1}$. Due to the norm properties, it follows that:

$$\begin{aligned} \|\psi_G(t_2) - \psi_G(t_1)\| &\leq \|\psi_G(t_1^2) - \psi_G(t_1)\| + \|\psi_G(t_1^3) - \psi_G(t_1^2)\| + \dots + \|\psi_G(t_2) - \psi_G(t_1^n)\| \leq \\ &\leq \frac{C}{M} \cdot \left\{ \left(\frac{2\ln(t_1)}{t_1^{3\rho}} \right)^2 + \left(\frac{3\ln(t_1)}{t_1^{2 \cdot 3\rho}} \right)^2 + \left(\frac{4\ln(t_1)}{t_1^{3 \cdot 3\rho}} \right)^2 + \dots + \left(\frac{(n+1)\ln(t_1)}{t_1^{n \cdot 3\rho}} \right)^2 \right\} = \\ &\leq \frac{C}{M \cdot t_1^\rho} \cdot \left\{ \left(\frac{2}{t_1^\rho} \cdot \frac{\ln(t_1)}{t_1^\rho} \right)^2 + \left(\frac{3}{t_1^{2\rho}} \cdot \frac{\ln(t_1)}{t_1^\rho} \right)^2 + \left(\frac{4}{t_1^{3\rho}} \cdot \frac{\ln(t_1)}{t_1^\rho} \right)^2 + \dots + \left(\frac{n+1}{t_1^{n\rho}} \cdot \frac{\ln(t_1)}{t_1^\rho} \right)^2 \right\} \end{aligned}$$

For t_1 sufficiently large $t_1 \geq \hat{t}_1 > \bar{t} \gg 1$ the sequence inside the brackets is convergent, and it is limited by a constant M .

I can conclude that $\forall t_1, t_2$ where $t_2 \geq t_1 \geq \hat{t}_1$

$$\|\psi_G(t_2) - \psi_G(t_1)\| \leq \frac{C}{t_1^\rho}$$

5 Scattering subspaces and asymptotic observables.

In this chapter I begin by constructing the subspace $H^{1out(in)}$ as the norm closure of the finite linear combinations of the vectors $\psi_G^{out(in)}$. The invariance requirement under space-time translations for the space $H^{1out(in)}$ implies a more general definition of the vector $\psi_G^{out(in)}$. Therefore it will be labelled as a $\psi_{G,\kappa_1,\tau,\mathbf{a}}^{out(in)}$. It corresponds to the evolution of $\psi_G^{out(in)}$ in the time τ and to the translation of a quantity $\mathbf{a}$.

I will verify that on $H^{1out(in)} \equiv \overline{\bigvee \psi_{G,\kappa_1,\tau,\mathbf{a}}^{out(in)}}$ the strong limits of the functions, continuous and of compact support, of nucleon mean velocity and the strong limits of the L.S.Z. Weyl operator associated to the meson field exist. Due to these results, we can

define the vectors $\psi_{G,\varphi}^{out(in)}$ (omitting $\kappa_1, \tau, \mathbf{a}$) obtained by applying the L.S.Z. Weyl operators, with smearing function φ , to the total set of $H^{1out(in)}$. The norm closure of finite linear combinations of the $\psi_{G,\varphi}^{out(in)}$ is a reasonable candidate for the scattering subspaces $H^{out(in)}$. The meaning of this definition is in the characterization of the $H^{out(in)}$ states, in terms of quantum numbers of the asymptotic variables which are defined on them: the asymptotic meson Weyl operators and the asymptotic nucleon mean velocity.

The spectral restriction on nucleon velocity (strictly less than 1) and the consequent restriction of $H^{out(in)}$ as subspaces of H are in accordance with the partial non-relativistic character of the model.

In theorem 5.1 the vectors $\psi_{G,\varphi}^{out(in)}$ are constructed. In theorem 5.2 we prove the convergence of continuous and of compact support functions of nucleon mean velocity on the vectors of $H^{out(in)}$. The corollary 5.3 is a check of the fact that the strong limits on $H^{out(in)}$ of the L.S.Z. Weyl operators generate a canonic Weyl algebra $\mathcal{A}^{out(in)}$, to which a free massless scalar field is associated.

Definition of the vector $\psi_{G,\kappa_1,\tau,\mathbf{a}}^{out(in)}$

I apply $e^{-i\mathbf{a}\cdot\mathbf{P}}e^{-iH\tau}$ to the generic vector ψ_G^{out} , constructed in the paragraphs 4.1,4.2,:

$$\begin{aligned} e^{-i\mathbf{a}\cdot\mathbf{P}}e^{-iH\tau}\psi_G^{out} &= \\ &= s\text{-}\lim_{t\rightarrow+\infty} e^{-i\mathbf{a}\cdot\mathbf{P}}e^{-iH\tau}e^{iHt} \sum_{i=1}^{N(t)} W_{\sigma_t}(\mathbf{v}_i, t) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), t)} e^{-iE^{\sigma_t}(t-\tau)} e^{-iE^{\sigma_t}\tau} \psi_{i,\sigma_t}^{(t)} = \\ &= s\text{-}\lim_{t\rightarrow+\infty} e^{iHt} \sum_{i=1}^{N(t)} e^{-g \int_{\sigma_t+\tau}^{\kappa_1} \frac{a(\mathbf{k})e^{i\mathbf{k}\cdot\mathbf{a}}e^{i|\mathbf{k}|(t+\tau)} - c.c.}{|\mathbf{k}|(1-\mathbf{k}\cdot\mathbf{v}_i)} \frac{d^3k}{\sqrt{2}|\mathbf{k}|}} e^{i\gamma_{\sigma_t+\tau}(\mathbf{v}_i, \nabla E^{\sigma_t+\tau}(\mathbf{P}), t+\tau)} e^{-i\mathbf{a}\cdot\mathbf{P}} e^{-iE^{\sigma_t+\tau}t} e^{-iE^{\sigma_t+\tau}\tau} \psi_{i,\sigma_t+\tau}^{(t+\tau)} \end{aligned}$$

The limit exists and the proof is similar to the one given for ψ_G^{out} , apart from some little marginal differences. Then I define

$$\psi_{G,\kappa_1,\tau,\mathbf{a}}^{out(in)} \equiv e^{-i\mathbf{a}\cdot\mathbf{P}}e^{-iH\tau}\psi_G^{out(in)}$$

The subspace of minimal asymptotic nucleon states is $H^{1out(in)} \equiv \overline{\bigvee \psi_{G,\kappa_1,\tau,\mathbf{a}}^{out(in)}}$. Later on I will simply call $\{\psi_G^{out(in)}\}$ the total set that generates $H^{1out(in)}$.

Theorem 5.1

The strong limits

$$s - \lim_{t \rightarrow +\infty} e^{iHt} e^{i(a(\varphi_t) + a^\dagger(\varphi_t))} e^{-iHt} \psi_G^{out} \equiv \psi_{G,\varphi}^{out}$$

exist, where $\tilde{\varphi}(\mathbf{k}) \in C_0^\infty(R^3 \setminus 0)$, $\varphi_t(\mathbf{y}) = \int e^{-i\mathbf{k}\mathbf{y} + i|\mathbf{k}|t} \tilde{\varphi}(\mathbf{k}) d^3k$ and $a^\dagger(\varphi) \equiv (a(\varphi))^\dagger = (\int a(\mathbf{k}) \tilde{\varphi}(\mathbf{k}) d^3k)^\dagger$

Proof

The convergence follows from the integrability of the norm

$$\begin{aligned} & \left\| \frac{d \left(e^{iHt} e^{i(a(\varphi_t) + a^\dagger(\varphi_t))} e^{-iHt} \right)}{dt} \psi_G^{out} \right\| = \\ & = \left\| e^{i(a(\varphi_t) + a^\dagger(\varphi_t))} \varphi(\mathbf{x}, t) e^{-iHt} \psi_G^{out} \right\| \leq \\ & \leq \left\| e^{i(a(\varphi_t) + a^\dagger(\varphi_t))} \varphi(\mathbf{x}, t) e^{-iHt} \psi_G^{out} - e^{i(a(\varphi_t) + a^\dagger(\varphi_t))} \varphi(\mathbf{x}, t) e^{-iHt} \psi_G(t) \right\| + \\ & + \left\| e^{i(a(\varphi_t) + a^\dagger(\varphi_t))} \varphi(\mathbf{x}, t) e^{-iHt} \psi_G(t) \right\| \leq \\ & \leq \left\| \varphi(\mathbf{x}, t) e^{-iHt} (\psi_G^{out} - \psi_G(t)) \right\| + \left\| \varphi(\mathbf{x}, t) e^{-iHt} \psi_G(t) \right\| \end{aligned}$$

where $\varphi(\mathbf{x}, t) \equiv -ig \int \left(\tilde{\varphi}(\mathbf{k}) e^{i|\mathbf{k}|t - i\mathbf{k} \cdot \mathbf{x}} + \overline{\tilde{\varphi}}(\mathbf{k}) e^{-i|\mathbf{k}|t + i\mathbf{k} \cdot \mathbf{x}} \right) \frac{d^3k}{\sqrt{2|\mathbf{k}|}}$

Both norms on the right hand side are bounded by quantities of order $\frac{1}{t^{1+\rho'}}$, $\rho' > 0$:

- the first one because $\sup_{\mathbf{x}} |\varphi(\mathbf{x}, t)| \leq \frac{M}{t}$ and $\|(\psi_G^{out} - \psi_G(t))\| \leq \frac{C}{t^\rho}$ (M and C are constants)
- as regards the second one, starting from the identity:

$$\begin{aligned} \varphi(\mathbf{x}, t) e^{-iHt} \psi_G(t) &= \varphi(\mathbf{x}, t) \sum_{i=1}^{N(t)} W_{\sigma_t}(\mathbf{v}_i, t) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), t)} e^{-iE^{\sigma_t} t} \psi_{i, \sigma_t}^{(t)} = \\ &= \sum_{i=1}^{N(t)} W_{\sigma_t}(\mathbf{v}_i, t) \varphi(\mathbf{x}, t) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), t)} e^{-iE^{\sigma_t} t} \int_{\Gamma_i} G(\mathbf{P}) \psi_{\mathbf{P}, \sigma_t} d^3P = \end{aligned}$$

$$= \sum_{i=1}^{N(t)} W_{\sigma_t}(\mathbf{v}_i, t) \varphi(\mathbf{x}, t) \left(1_{\Gamma_i}(\mathbf{P}) - \chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, t) \right) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), t)} e^{-iE^{\sigma_t}t} \int_{\Gamma_i} G(\mathbf{P}) \psi_{\mathbf{P}, \sigma_t} d^3P + \quad (33.1)$$

$$+ \sum_{i=1}^{N(t)} W_{\sigma_t}(\mathbf{v}_i, t) \varphi(\mathbf{x}, t) \left(\chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, t) - \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{t}, t\right) \right) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), t)} e^{-iE^{\sigma_t}t} \int_{\Gamma_i} G(\mathbf{P}) \psi_{\mathbf{P}, \sigma_t} d^3P + \quad (33.2)$$

$$+ \sum_{i=1}^{N(t)} W_{\sigma_t}(\mathbf{v}_i, t) \varphi(\mathbf{x}, t) \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{t}, t\right) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), t)} e^{-iE^{\sigma_t}t} \int_{\Gamma_i} G(\mathbf{P}) \psi_{\mathbf{P}, \sigma_t} d^3P \quad (33.3)$$

I exploit lemma B1 for 33.1, lemma B2 for 33.2, and Huygens principle in order to estimate $\sup_{\mathbf{x}} \left| \varphi(\mathbf{x}, t) \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{t}, t\right) \right|$ in the expression (33.3).

The scattering subspaces $H^{out(in)} \equiv \overline{\bigvee \psi_{G, \varphi}^{out(in)}}$ are invariant under space-time translations because the subspaces $H^{1out(in)}$ are invariant.

Theorem 5.2

The nucleon mean velocity functions f , which are continuous and of compact support, have asymptotically strong limits H^{out} ; in particular:

$$s - \lim_{t \rightarrow +\infty} e^{iHt} f\left(\frac{\mathbf{x}}{t}\right) e^{-iHt} \psi_{G, \varphi}^{out} = \psi_{G, \hat{f}}^{out} \quad \text{where} \quad \hat{f}(\mathbf{P}) \equiv \lim_{\sigma \rightarrow 0} f(\nabla E^\sigma(\mathbf{P}))$$

Proof

It is sufficient to prove it on the vectors $\psi_{G, \varphi}^{out}$ for functions $f \in C_0^\infty(R^3)$. Exploiting theorem 5.1 and the uniform boundedness in t of the operators $f\left(\frac{\mathbf{x}}{t}\right)$ and $e^{iHt} e^{i(a(\varphi_t) + a^\dagger(\varphi_t))} e^{-iHt}$ we obtain

$$\begin{aligned} s - \lim_{t \rightarrow +\infty} e^{iHt} f\left(\frac{\mathbf{x}}{t}\right) e^{-iHt} \psi_{G, \varphi}^{out} &= s - \lim_{t \rightarrow +\infty} e^{iHt} f\left(\frac{\mathbf{x}}{t}\right) e^{i(a(\varphi_t) + a^\dagger(\varphi_t))} e^{-iHt} \psi_G^{out} = \\ &= s - \lim_{t \rightarrow +\infty} e^{iHt} e^{i(a(\varphi_t) + a^\dagger(\varphi_t))} f\left(\frac{\mathbf{x}}{t}\right) e^{-iHt} \psi_G^{out} = \end{aligned}$$

$$= s - \lim_{t \rightarrow +\infty} e^{iHt} e^{i(a(\varphi_t) + a^\dagger(\varphi_t))} e^{-iHt} \psi_{G, \hat{f}}^{out}$$

(the last step is proved by the technique used in lemma B2)

The extension to all of H^{out} is automatic since $f\left(\frac{\mathbf{x}}{t}\right)$ is uniformly bounded in t

and moreover the set $\bigvee \psi_{G,\varphi}^{out}$ is a dense set in H^{out} , by construction of H^{out} .

Corollary 5.3

In the space H^{out} , the asymptotic meson algebra $\mathcal{A}^{out}$ is defined as the norm closure of the $*$ algebra generated by the set of unitary operators $\{W^{out}(\mu) : \tilde{\mu} \in C_0^\infty(R^3 \setminus 0)\}$ constructed in H^{out} as follows:

$$W^{out}(\mu) = s - \lim_{t \rightarrow +\infty} e^{iHt} e^{i(a(\mu_t) + a^\dagger(\mu_t))} e^{-iHt} \quad (34)$$

The following properties hold:

1) the generators $\{W^{out}(\mu) : \tilde{\mu} \in C_0^\infty(R^3 \setminus 0)\}$ satisfy the Weyl commutation rules

$$W^{out}(\mu) W^{out}(\eta) = W^{out}(\eta) W^{out}(\mu) e^{-h(\mu, \eta)} \quad \text{where} \quad h(\mu, \eta) = 2i \operatorname{Im} \int \tilde{\mu}(\mathbf{k}) \bar{\tilde{\eta}}(\mathbf{k}) d^3k;$$

2) for each fixed region $R^3 \setminus O_{\mathbf{a}}$ where $O_{\mathbf{a}}$ is a ball of radius $a \neq 0$, centered in the origin of R^3 , the group of the operators $W^{out}(\mu)$, where $\tilde{\mu} \in C_0^\infty(R^3 \setminus O_{\mathbf{a}})$, is strongly continuous with respect to $\tilde{\mu}$ in the $L^2(R^3 \setminus O_{\mathbf{a}}, d^3k)$ norm;

3) given the τ -evolved generators:

$$W_\tau^{out}(\mu) = s - \lim_{t \rightarrow +\infty} e^{iH(t+\tau)} e^{i(a(\mu_t) + a^\dagger(\mu_t))} e^{-iH(t+\tau)} = W^{out}(\mu_{-\tau}) \quad (35)$$

it is uniquely defined an automorphism α_τ of $\mathcal{A}^{out}$ starting from $\alpha_\tau(W^{out}(\mu)) = W^{out}(\mu_{-\tau})$. Therefore, being $\mu_{-\tau}$ the test function μ free evolved in the time τ , we can conclude that $\mathcal{A}^{out}$ is the Weyl algebra associated to the scalar massless field.

4) the algebra $\mathcal{A}^{out}$ commutes with the asymptotic nucleon mean velocity defined through theorem 5.2.

$\Leftrightarrow$

Proof

The existence of $s - \lim_{t \rightarrow +\infty} e^{iHt} e^{i(a(\mu_t) + a^\dagger(\mu_t))} e^{-iHt} \psi_{G,\varphi}^{out}$ is substantially the contents of theorem 5.1. The bounded operators $W^{out}(\mu)$, defined on the dense set

$\{\psi_{G,\varphi}^{out}\}$ of H^{out} , can be linearly extended to all of H^{out} by continuity. They leave the space H^{out} invariant and they are unitary in H^{out} .

1) On the vectors in H^{out} , $W^{out}(\mu)W^{out}(\eta)$ is the limit of the product of the approximating vectors (34), at the same time t . The last ones satisfy the property by construction. Therefore, the property is satisfied in the limit.

2) It is sufficient to prove that $W^{out}(\mu)$ is strongly continuous with respect to $\tilde{\mu}$ if it is applied to the total set $\{\psi_{G,\varphi}^{out}\}$:

- $W^{out}(\mu)\psi_{G,\varphi}^{out} = s - \lim_{t \rightarrow +\infty} e^{iHt} e^{i(a(\mu_t)+a^\dagger(\mu_t))} e^{-iHt} \psi_{G,\varphi}(t)$;
- at fixed t , the vector $e^{i(a(\mu_t)+a^\dagger(\mu_t))} e^{-iHt} \psi_{G,\varphi}(t)$ is strongly continuous with respect to $\tilde{\mu} \in C_0^\infty(R^3 \setminus O_a)$. This is due to the fact that

$$\begin{aligned} & e^{i(a(\mu_t)+a^\dagger(\mu_t))} \sum_{i=1}^{N(t)} W_{\sigma_t}(\mathbf{v}_i, t) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), t)} e^{-iE^{\sigma_t}t} \psi_{i,\sigma_t}^{(t)} = \\ & = \sum_{i=1}^{N(t)} W_{\sigma_t}(\mathbf{v}_i, t) e^{i(a(\mu_t)+a^\dagger(\mu_t))} e^{-h(\mu_t, \xi_{\mathbf{v}_i})} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}(\mathbf{P}), t)} e^{-iE^{\sigma_t}t} \psi_{i,\sigma_t}^{(t)} \end{aligned}$$

where $\xi_{\mathbf{v}_i} \equiv -ig \frac{\chi_{\sigma_i}^{\kappa_1}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1-\widehat{\mathbf{k}} \cdot \mathbf{v}_i)}$ and because $\psi_{i,\sigma_i}^{(t)}$ is in the domain of the generator of $e^{i(a(\mu_t)+a^\dagger(\mu_t))}$. Being $\tilde{\mu}_t \in C_0^\infty(R^3 \setminus O_a)$ and $\|\tilde{\mu}_t\|_{L^2(R^3 \setminus O_a, d^3k)} = \|\tilde{\mu}\|_{L^2(R^3 \setminus O_a, d^3k)}$, the vector is strongly continuous (with respect to $\tilde{\mu}$) uniformly in t . Since $\|W^{out}(\mu)\| = 1$, the property holds for each vector in H^{out} .

3) The τ -evolved generators $e^{iH\tau}W^{out}(\mu)e^{-iH\tau}$ are well defined because $e^{-iH\tau}:H^{out} \rightarrow H^{out}$. By inserting the expression (34) for $W^{out}(\mu)$, we arrive at the (35). The Weyl commutation rules are conserved by α_τ since

$$h(\mu_{-\tau}, \eta_{-\tau}) = 2iIm \int \tilde{\mu}(\mathbf{k}) \bar{\tilde{\eta}}(\mathbf{k}) d^3k = h(\mu, \eta)$$

4) Such a property is implicit in the construction of the asymptotic nucleon mean velocity (theorem 5.2).

$\Longleftrightarrow$

APPENDIX A

Preliminary remarks to the lemma A1

Like in lemma 1.4, it will be proved that the operator $(\Delta H_{\mathbf{P}}^w)_{\epsilon}^{\frac{j+1}{2}} \equiv \widehat{H}_{\mathbf{P}, \epsilon}^w + c_{\mathbf{P}}(j+1) - \widehat{c}_{\mathbf{P}}(j+2) - H_{\mathbf{P}, \epsilon}^w \psi_G^{out(in)}$ is small of order 1 with respect to $H_{\mathbf{P}, \epsilon}^w \big|_{F_{\epsilon}^+}$ in a generalized sense for g sufficiently small. We aim at expanding the resolvent

$$\frac{1}{\widehat{H}_{\mathbf{P}, \epsilon}^w - (E(j+1) + \widehat{c}_{\mathbf{P}}(j+2) - c_{\mathbf{P}}(j+1))} \big|_{F_{\epsilon}^+}$$

$$(E(j+1) \in \mathcal{C} \text{ s.t. } \left| E(j+1) - E_{\mathbf{P}}^{\epsilon} \right| = \frac{11\epsilon^{\frac{j+2}{2}}}{20}, \widehat{c}_{\mathbf{P}}(j+2) - c_{\mathbf{P}}(j+1) = -g^2 \int_{\epsilon}^{\frac{j+1}{2}} \frac{1}{2|\mathbf{k}|^2 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{\epsilon} \left(\frac{j+1}{2}(\mathbf{P}) \right) \right)} d^3 k)$$

in terms of $\frac{1}{H_{\mathbf{P}, \epsilon}^w - E(j+1)} \big|_{F_{\epsilon}^+}$ and $(\Delta H_{\mathbf{P}}^w)_{\epsilon}^{\frac{j+1}{2}}$.

I will only treat the case $\mathbf{P} = 0$, where the difference operator is written $(\Delta H^w)_{\epsilon}^{\frac{j+1}{2}} \equiv H_{\epsilon}^w + c(j+1) - c(j+2) - H_{\epsilon}^w$.

In the case $\mathbf{P} \neq 0$ the steps are analogous and well defined since the module of $\nabla E^{\sigma}(\mathbf{P})$ is uniformly bounded in σ by a positive constant less than 1, for $\mathbf{P} \in \Sigma$ (see lemma A2).

Lemma A1

Given the spectral properties in the beginning of paragraph 2.1, $(\Delta H^w)_{\epsilon}^{\frac{j+1}{2}} \big|_{F_{\epsilon}^+}$ is small with respect to H_{ϵ}^w for values of g less than a proper $\overline{g}$ and $\mathbf{P} \in \Sigma$, in the following sense:

given $E(j+1) \in C$ s.t. $\left| E(j+1) - E^{\epsilon \frac{j+1}{2}} \right| = \frac{11\epsilon \frac{j+2}{2}}{20}$,

$$\begin{aligned}
& \frac{1}{H^w_{\epsilon \frac{j+2}{2}} - (E(j+1) + c(j+2) - c(j+1))} \Big|_{F^+_{\epsilon \frac{j+2}{2}}} = \\
& = \frac{1}{H^w_{\epsilon \frac{j+1}{2}} + (\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon} - c(j+1) + c(j+2) - (E(j+1) - c(j+1) + c(j+2))} \Big|_{F^+_{\epsilon \frac{j+2}{2}}} = \\
& = \frac{1}{H^w_{\epsilon \frac{j+1}{2}} + (\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon} - E(j+1)} \Big|_{F^+_{\epsilon \frac{j+2}{2}}} = \\
& = \frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \Big|_{F^+_{\epsilon \frac{j+2}{2}}} + \frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \sum_{n=1}^{+\infty} \left(-(\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \right)^n \Big|_{F^+_{\epsilon \frac{j+2}{2}}}
\end{aligned}$$

where

$$\begin{aligned}
& \bullet \left\| \frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \left(-(\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \right)^n \right\|_{F^+_{\epsilon \frac{j+2}{2}}} \leq \frac{20(C(\bar{g}, \bar{m}))^n}{\epsilon \frac{j+2}{2}} \\
& \bullet 0 < C(\bar{g}, m) < \frac{1}{12} .
\end{aligned}$$

Proof

Like in lemma 1.4 we can state

$$\begin{aligned}
& \frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \left(-(\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \right)^n = \\
& = (-1)^n \frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} (\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \dots (\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} = \\
& = (-1)^n \left(\frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \dots \left(\frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} (\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \left(\frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \dots \left(\frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \\
& \textbf{Study of the norm of} \quad \left(\frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} (\Delta H^w)_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \left(\frac{1}{H^w_{\epsilon \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \Big|_{F^+_{\epsilon \frac{j+2}{2}}} .
\end{aligned}$$

$$\begin{aligned}
(\Delta H^w)_\epsilon^{\frac{j+1}{2}} &= -\frac{g}{2m} \int_\epsilon^{\frac{j+1}{2}} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) d^3k \cdot \Pi_{\epsilon^{\frac{j+1}{2}}} + \\
&- \Pi_{\epsilon^{\frac{j+1}{2}}} \cdot \frac{g}{2m} \int_\epsilon^{\frac{j+1}{2}} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) d^3k + \frac{g^2}{2m} \left(\int_\epsilon^{\frac{j+1}{2}} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) d^3k \right)^2
\end{aligned}$$

Making the calculations we have:

$$\begin{aligned}
\frac{g^2}{2m} \left(\int_\epsilon^{\frac{j+1}{2}} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}} \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) d^3k \right)^2 &= \frac{g^2}{2m} \sum_i \int_\epsilon^{\frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3k}{|\mathbf{k}|\sqrt{2}|\mathbf{k}|} \int_\epsilon^{\frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3k}{|\mathbf{k}|\sqrt{2}|\mathbf{k}|} + \\
&+ \frac{g^2}{2m} \sum_i \int_\epsilon^{\frac{j+1}{2}} k^i b^\dagger(\mathbf{k}) \frac{d^3k}{|\mathbf{k}|\sqrt{2}|\mathbf{k}|} \int_\epsilon^{\frac{j+1}{2}} k^i b^\dagger(\mathbf{k}) \frac{d^3k}{|\mathbf{k}|\sqrt{2}|\mathbf{k}|} + \\
&+ \frac{g^2}{m} \sum_i \int_\epsilon^{\frac{j+1}{2}} k^i b^\dagger(\mathbf{k}) \frac{d^3k}{|\mathbf{k}|\sqrt{2}|\mathbf{k}|} \int_\epsilon^{\frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3k}{|\mathbf{k}|\sqrt{2}|\mathbf{k}|} + \frac{g^2}{4m} \int_\epsilon^{\frac{j+1}{2}} \frac{d^3k}{|\mathbf{k}|}
\end{aligned}$$

We have to examine:

$$\begin{aligned}
1) & \frac{g^2}{2m} \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_\epsilon^{\frac{j+1}{2}} k^i b^\dagger(\mathbf{k}) \frac{d^3k}{|\mathbf{k}|\sqrt{2}|\mathbf{k}|} \int_\epsilon^{\frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3k}{|\mathbf{k}|\sqrt{2}|\mathbf{k}|} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \\
2) & \frac{g^2}{2m} \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_\epsilon^{\frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3k}{|\mathbf{k}|\sqrt{2}|\mathbf{k}|} \int_\epsilon^{\frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3k}{|\mathbf{k}|\sqrt{2}|\mathbf{k}|} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \\
3) & \frac{g^2}{m} \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} k^i b^\dagger(\mathbf{k}) \frac{d^3k}{|\mathbf{k}|\sqrt{2}|\mathbf{k}|} \int_\epsilon^{\frac{j+1}{2}} k^i b^\dagger(\mathbf{k}) \frac{d^3k}{|\mathbf{k}|\sqrt{2}|\mathbf{k}|} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \\
4) & \frac{g^2}{4m} \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_\epsilon^{\frac{j+1}{2}} \frac{d^3k}{|\mathbf{k}|} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+}
\end{aligned}$$

$$\begin{aligned}
5) & \frac{g}{2m} \left\| \left(\frac{1}{H_{\epsilon}^{w, \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \Pi_{\epsilon}^{i, \frac{j+1}{2}} \cdot \int_{\epsilon}^{\epsilon, \frac{j+1}{2}} k^i (b(\mathbf{k}) + b^\dagger(\mathbf{k})) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon}^{w, \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon}^{+, \frac{j+2}{2}}} \\
6) & \frac{g}{2m} \left\| \left(\frac{1}{H_{\epsilon}^{w, \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon}^{\epsilon, \frac{j+2}{2}} k^i (b(\mathbf{k}) + b^\dagger(\mathbf{k})) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \cdot \Pi_{\epsilon}^{i, \frac{j+1}{2}} \left(\frac{1}{H_{\epsilon}^{w, \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon}^{+, \frac{j+2}{2}}}
\end{aligned}$$

In order to control the above quantities, I will use the following estimate again and again

$$\left\| \int_{\epsilon}^{\epsilon, \frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon}^{w, \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon}^{+, \frac{j+2}{2}}} \leq \sqrt{40\pi} \cdot \epsilon^{\frac{j+1}{4}} \quad (\text{a1})$$

which is proved like the estimate (4) of lemma 1.4., by performing an unitary transformation.

I study the quadratic quantities (1,2,3,4) and then the mixed terms (5,6) which contain the $\Pi_{\epsilon}^{i, \frac{j+1}{2}}$. The following estimates are worked out:

1)

$$\begin{aligned}
& \left\| \left(\frac{1}{H_{\epsilon}^{w, \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon}^{\epsilon, \frac{j+1}{2}} k^i b^\dagger(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \int_{\epsilon}^{\epsilon, \frac{j+2}{2}} k^i b(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon}^{w, \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon}^{+, \frac{j+2}{2}}} \leq \\
& \leq \left\| \left(\frac{1}{H_{\epsilon}^{w, \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon}^{\epsilon, \frac{j+1}{2}} k^i b^\dagger(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \right\|_{F_{\epsilon}^{+, \frac{j+2}{2}}} \cdot \left\| \int_{\epsilon}^{\epsilon, \frac{j+2}{2}} k^i b(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon}^{w, \frac{j+1}{2}} - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon}^{+, \frac{j+2}{2}}}
\end{aligned}$$

if the norms on the right hand side exist.

Note that

$$\left\| \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \right\|_{F_{\epsilon \frac{j+2}{2}}^+} = \left\| \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left[\left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^\dagger \right\|_{F_{\epsilon \frac{j+2}{2}}^+}$$

the norm on the right hand side is controlled like

$$\left\| \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon \frac{j+2}{2}}^+}.$$

In conclusion

$$\begin{aligned} & \frac{g^2}{2m} \left\| \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon \frac{j+2}{2}}^+} \leq \\ & \leq C_1^i(g, m) \cdot \epsilon^{\frac{j+1}{2}} \end{aligned}$$

2)

$$\begin{aligned} & \left\| \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon \frac{j+2}{2}}^+} \leq \\ & \leq \left\| \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \right\|_{F_{\epsilon \frac{j+2}{2}}^+} \cdot \left\| \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} k^i b(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon \frac{j+2}{2}}^+} \end{aligned}$$

if the norms on the right hand side exist.

I evaluate the norm of $\int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left[\left(\frac{1}{H_{\epsilon \frac{j+1}{2}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^\dagger$ restricted to $F_{\epsilon \frac{j+2}{2}}^+$.

Given $\varphi \in D^b \cap F_{\epsilon \frac{j+2}{2}}^+$:

$$\begin{aligned}
& \left\| \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^\dagger \varphi \right\|^2 = \\
& = \left(\int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^\dagger \varphi, \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^\dagger \varphi \right) = \\
& \leq \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{(k^i)^2 d^3 k}{2|\mathbf{k}|^3} \left(\left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^\dagger \varphi, \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^\dagger \varphi \right) + \\
& + \left(\int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^\dagger \varphi, \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^\dagger \varphi \right) \leq \\
& \leq \pi \cdot \epsilon^{j+1} \cdot \frac{20}{\epsilon^{\frac{j+2}{2}}} \cdot \|\varphi\|^2 + 40\pi \epsilon^{\frac{j+1}{2}} \cdot \|\varphi\|^2
\end{aligned}$$

Then

$$\begin{aligned}
& \frac{g^2}{2m} \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \leq \\
& \leq C_2^i(g, m) \cdot \epsilon^{\frac{j}{2}}
\end{aligned}$$

3)

$$\begin{aligned}
& \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \leq \\
& \leq \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \cdot \left\| \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+}
\end{aligned}$$

• the norm $\left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+}$ was treated at point 1);

- the norm $\left\| \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+}$ can be controlled like the
- norm $\left\| \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^\dagger \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+}$ studied at point 2).

In conclusion:

$$\begin{aligned} & \frac{g}{m} \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \leq \\ & \leq C_3^i(g, m) \cdot \epsilon^{\frac{j}{2}} \end{aligned}$$

4)

$$\begin{aligned} & \frac{g^2}{4m} \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{d^3 k}{|\mathbf{k}|} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} = \\ & = \frac{g^2}{4m} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{d^3 k}{|\mathbf{k}|} \left\| \frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \leq C_4^i(g, m) \cdot \epsilon^{\frac{j}{2}} \end{aligned}$$

5)

$$\frac{g^2}{2m} \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \cdot \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i (b(\mathbf{k}) + b^\dagger(\mathbf{k})) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+}$$

5A)

$$\frac{1}{\sqrt{2m}} \cdot \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \cdot \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \leq$$

$$\leq \frac{1}{\sqrt{2m}} \cdot \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+}^{\frac{1}{2}} \cdot \left\| \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b(\mathbf{k}) \frac{d^3 \mathbf{k}}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+}^{\frac{1}{2}}$$

if the two norms on the right hand side exist.

In order to prove the existence and the bound of the second one see **1**).

As regards the bound of the first one, we can start from the equality (verified “a posteriori” thanks to the existence of the norm on the right hand side)

$$\left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+}^{\frac{1}{2}} = \left\| \Pi_{\epsilon^{\frac{j+1}{2}}}^i \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^{\dagger} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+}^{\frac{1}{2}}$$

and from the estimate ($\varphi \in D^b \cap F_{\epsilon^{\frac{j+2}{2}}}^+$)

$$\begin{aligned} & \frac{1}{2m} \left\| \Pi_{\epsilon^{\frac{j+1}{2}}}^i \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^{\dagger} \varphi \right\|^2 = \\ &= \frac{1}{2m} \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^{\dagger} \varphi, \Pi_{\epsilon^{\frac{j+1}{2}}}^i \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^{\dagger} \varphi \right) \leq \\ &\leq \frac{1}{2m} \sum_i \left(\Pi_{\epsilon^{\frac{j+1}{2}}}^i \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^{\dagger} \varphi, \Pi_{\epsilon^{\frac{j+1}{2}}}^i \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^{\dagger} \varphi \right) \leq \\ &\leq \left(\left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^{\dagger} \varphi, \left(\frac{1}{2m} \cdot \Pi_{\epsilon^{\frac{j+1}{2}}}^2 + H^{mes} \right) \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^{\dagger} \varphi \right) \leq \\ &= \left(\varphi, \left(H_{\epsilon^{\frac{j+1}{2}}}^w - c(j+1) \right) \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \left[\left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right]^{\dagger} \varphi \right) \leq \frac{40}{\epsilon^{\frac{j+2}{2}}} \|\varphi\|^2 \end{aligned}$$

(I used the fact that $|c(j+1)| \leq 1$ and $\left| E_{\epsilon^{\frac{j+1}{2}}}^{\epsilon^{\frac{j+1}{2}}} \right| \leq 1$)

5B)

Commuting and exploiting the properties of the norm we have:

$$\begin{aligned}
& \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \cdot \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \leq \\
& \leq \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} + \\
& + \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} (k^i)^2 b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \\
& \text{i) the term } \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \\
& \text{can be controlled by } \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \cdot \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i b(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} ; \\
& \text{ii) the term } \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} (k^i)^2 b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \\
& \text{is bounded by} \\
& \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} (k^i)^2 b^\dagger(\mathbf{k}) \frac{d^3 k}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \cdot \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \\
& \text{that is of order } \frac{\epsilon^{\frac{j+1}{2}}}{\epsilon^{\frac{j+2}{4}}}.
\end{aligned}$$

Summarizing:

$$\begin{aligned}
& \left\| \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \Pi_{\epsilon^{\frac{j+1}{2}}}^i \cdot \int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} k^i \left(b(\mathbf{k}) + b^\dagger(\mathbf{k}) \right) \frac{d^3 k}{|\mathbf{k}| \sqrt{2} |\mathbf{k}|} \left(\frac{1}{H_{\epsilon^{\frac{j+1}{2}}}^w - E(j+1)} \right)^{\frac{1}{2}} \right\|_{F_{\epsilon^{\frac{j+2}{2}}}^+} \leq \\
& \leq C_5^i(g, m) \cdot \frac{\epsilon^{\frac{j+1}{4}}}{\epsilon^{\frac{j+2}{4}}} = C_5^i(g, m) \cdot \epsilon^{-\frac{1}{4}}
\end{aligned}$$

6)

It is controlled like the expression 5)

Conclusion

If g is less than a limit value $\bar{g}$, the $\epsilon^{\frac{j+1}{2}}$ – independent constants $C_1^i(g, m), \dots, C_6^i(g, m)$ can be tuned in order to arrive at the thesis.

Lemma A2

Results about $\nabla E^\sigma(\mathbf{P})$ where $\mathbf{P} \in \Sigma$:

- 1) $|\nabla E^\sigma(\mathbf{P})| < v^{max} < 1 \quad \forall \sigma$;
- 2) $\left| \nabla E^{\epsilon^{\frac{j+1}{2}}}(\mathbf{P}) - \nabla E^{\epsilon^{\frac{j+2}{2}}}(\mathbf{P}) \right| < C(g) \cdot \epsilon^{\frac{j+1}{8}}$ for g sufficiently small and uniform in j ;
- 3) $\nabla E^\sigma(\mathbf{P})$ and $\frac{1}{\|\tilde{\phi}_\mathbf{P}^\sigma\|^2} \left(\tilde{\phi}_\mathbf{P}^\sigma, \Pi_{\mathbf{P}, \sigma} \tilde{\phi}_\mathbf{P}^\sigma \right)$ converge for $\sigma \rightarrow 0$ (for the definition of $\tilde{\phi}_\mathbf{P}^\sigma$ see chapter 3 “Spectral regularity”).

$\Leftrightarrow$

Proof

$$1) |\nabla E^\sigma(\mathbf{P})| = \frac{|(\psi_\mathbf{P}^\sigma, \mathbf{P} - \mathbf{P}^{mes} \psi_\mathbf{P}^\sigma)|}{m \|\psi_\mathbf{P}^\sigma\|^2} \leq \sqrt{\frac{2}{m}} \cdot \frac{1}{\|\psi_\mathbf{P}^\sigma\|} \cdot |(\psi_\mathbf{P}^\sigma, H_{\mathbf{P}, \sigma} + 2\pi g^2 \kappa \psi_\mathbf{P}^\sigma)|^{\frac{1}{2}} < 1$$

since, according to the initial hypotheses, we have

$$\frac{|(\psi_{\mathbf{P}}^\sigma, H_{\mathbf{P}, \sigma} \psi_{\mathbf{P}}^\sigma)|}{\|\psi_{\mathbf{P}}^\sigma\|^2} \leq |(\psi_0, H_{\mathbf{P}, \sigma} \psi_0)| = \frac{\mathbf{P}^2}{2m} \leq \frac{1}{2}$$

$$\frac{|(\psi_{\mathbf{P}}^\sigma, 2\pi g^2 \kappa \psi_{\mathbf{P}}^\sigma)|}{\|\psi_{\mathbf{P}}^\sigma\|^2} \leq \frac{1}{4}$$

2)

The property to be proved is involved in the proof of theorem 2.3bis. We assume an implication of the inductive hypothesis

$$\begin{aligned} & \left\| \frac{\widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}}}{\widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}}} - \frac{\phi_{\mathbf{P}}^{\frac{j+1}{2}}}{\phi_{\mathbf{P}}^{\frac{j+1}{2}}} \right\| = \left\| \frac{\widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}}}{\widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}}} - \frac{\phi_{\mathbf{P}}^{\frac{j+1}{2}}}{\phi_{\mathbf{P}}^{\frac{j+1}{2}}} \right\| = \left\| \frac{\widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}} - \phi_{\mathbf{P}}^{\frac{j+1}{2}}}{\widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}}} \right\| \leq \\ & \leq \frac{\left\| \widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}} - \phi_{\mathbf{P}}^{\frac{j+1}{2}} \right\|}{\left\| \widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}} \right\|} + \frac{\left\| \widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}} \right\| - \left\| \phi_{\mathbf{P}}^{\frac{j+1}{2}} \right\|}{\left\| \widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}} \right\|} \leq \\ & \leq \frac{2 \left\| \widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}} - \phi_{\mathbf{P}}^{\frac{j+1}{2}} \right\|}{\left\| \widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}} \right\|} \leq 3 \left\| -\widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}} + \phi_{\mathbf{P}}^{\frac{j+1}{2}} \right\| \end{aligned}$$

(by induction, from theorem 2.3bis (case $\mathbf{P} \neq 0$) we have:

$$\begin{aligned} \|\phi_{\mathbf{P}}^\epsilon\| & \leq \left\| \widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}} \right\| + \left\| -\widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}} + \phi_{\mathbf{P}}^{\frac{j+1}{2}} \right\| + \left\| -\widehat{\phi}_{\mathbf{P}}^{\frac{j+1}{2}} + \phi_{\mathbf{P}}^{\frac{j+1}{2}} \right\| + \dots + \left\| -\widehat{\phi}_{\mathbf{P}}^\epsilon + \phi_{\mathbf{P}}^\epsilon \right\| \\ \left\| \widehat{\phi}_{\mathbf{P}}^{\frac{j+2}{2}} \right\| & \geq \|\phi_{\mathbf{P}}^\epsilon\| - \left(\epsilon^{\frac{1}{16}} + \epsilon^{\frac{2}{16}} + \dots + \epsilon^{\frac{j+1}{16}} \right) \geq 1 - \left(\frac{\epsilon^{\frac{1}{16}}}{1 - \epsilon^{\frac{1}{16}}} \right) \geq \frac{1 - 2\epsilon^{\frac{1}{16}}}{1 - \epsilon^{\frac{1}{16}}} > \frac{2}{3} \end{aligned}$$

Let us analyze the difference between the gradients of energy.

$$\begin{aligned} m \nabla E_{\mathbf{P}}^{\frac{j+1}{2}} & = \mathbf{P} - \frac{\left(\phi_{\mathbf{P}}^{\frac{j+1}{2}}, \mathbf{P}^{mes} \phi_{\mathbf{P}}^{\frac{j+1}{2}} \right)}{\left\| \phi_{\mathbf{P}}^{\frac{j+1}{2}} \right\|^2} + \frac{1}{\left\| \phi_{\mathbf{P}}^{\frac{j+1}{2}} \right\|^2} \left(\phi_{\mathbf{P}}^{\frac{j+1}{2}}, g \int_{\epsilon^{\frac{j+1}{2}}}^{\kappa} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}}(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^\sigma)} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3 k \phi_{\mathbf{P}}^{\frac{j+1}{2}} \right) + \\ & - g^2 \int_{\epsilon^{\frac{j+1}{2}}}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\frac{j+1}{2}} \right)^2} d^3 k = \end{aligned}$$

$$= \mathbf{P} - \frac{\left(\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \Pi_{\mathbf{P}, \epsilon \frac{j+1}{2}} \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)}{\left\| \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|^2} - g^2 \int_{\epsilon \frac{j+1}{2}}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)^2} d^3 k$$

while

$$m \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} = \mathbf{P} - \frac{\left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, \widehat{\Pi}_{\mathbf{P}, \epsilon \frac{j+2}{2}} \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)}{\left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right\|^2} - g^2 \int_{\epsilon \frac{j+2}{2}}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)^2} d^3 k$$

then

$$\begin{aligned} m \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - m \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} - g^2 \int_{\epsilon \frac{j+2}{2}}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)^2} d^3 k + g^2 \int_{\epsilon \frac{j+1}{2}}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)^2} d^3 k = \\ = \frac{1}{\left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right\|^2} \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, \widehat{\Pi}_{\mathbf{P}, \epsilon \frac{j+2}{2}} \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right) - \frac{1}{\left\| \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|^2} \left(\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \Pi_{\mathbf{P}, \epsilon \frac{j+1}{2}} \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) \end{aligned} \quad (\text{a2})$$

$$\begin{aligned} m \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - m \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} + g^2 \int_{\epsilon \frac{j+1}{2}}^{\kappa} \frac{\mathbf{k} \left(-\widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} + \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) \left(2 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)^2 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)^2} d^3 k - g^2 \int_{\epsilon \frac{j+2}{2}}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)^2} d^3 k = \\ = \frac{1}{\left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right\|^2} \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, \widehat{\Pi}_{\mathbf{P}, \epsilon \frac{j+2}{2}} \left(\frac{\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}}{\left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right\|} - \frac{\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}}{\left\| \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|} \right) \right) + \\ + \frac{1}{\left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right\| \left\| \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|} \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, \widehat{\Pi}_{\mathbf{P}, \epsilon \frac{j+2}{2}} \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) - \frac{1}{\left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right\| \left\| \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|} \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, \Pi_{\mathbf{P}, \epsilon \frac{j+1}{2}} \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) + \\ + \frac{1}{\left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right\| \left\| \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|} \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, \Pi_{\mathbf{P}, \epsilon \frac{j+1}{2}} \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) - \frac{1}{\left\| \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|^2} \left(\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \Pi_{\mathbf{P}, \epsilon \frac{j+1}{2}} \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) \end{aligned}$$

Considering that

$$\begin{aligned} \widehat{\Pi}_{\mathbf{P}, \epsilon \frac{j+2}{2}}^i - \Pi_{\mathbf{P}, \epsilon \frac{j+1}{2}}^i = \\ = -g \int_{\epsilon \frac{j+2}{2}}^{\frac{j+1}{2}} \frac{k^i (b(\mathbf{k}) + b^\dagger(\mathbf{k}))}{\sqrt{2} |\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)} d^3 k + \frac{g^2}{2} \int_{\epsilon \frac{j+2}{2}}^{\kappa} \frac{k^i \left(-\widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} + \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) \left(2 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)^2 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)^2} d^3 k \end{aligned}$$

the equation (a2) can be written in the following way

$$\begin{aligned}
& m \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - m \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} + g^2 \int_{\epsilon \frac{j+1}{2}}^{\kappa} \frac{\mathbf{k} \left(\widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) \left(2 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)^2 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)^2} d^3 k + \\
& - \frac{1}{\left\| \frac{\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}}{\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}} \right\|} \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) \frac{g^2 \int_{\epsilon \frac{j+1}{2}}^{\kappa} \frac{\mathbf{k} \left(-\widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} + \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) \left(2 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)^2 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)^2} d^3 k = \\
& = \frac{1}{\left\| \frac{\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}}{\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}} \right\|} \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, \widehat{\Pi}_{\mathbf{P}, \epsilon \frac{j+2}{2}} \left(\frac{\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}}{\left\| \frac{\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}}{\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}} \right\|} - \frac{\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}}{\left\| \frac{\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}}{\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}} \right\|} \right) \right) + \frac{1}{\left\| \frac{\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}}{\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}} \right\|} \left(\left(\frac{\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}}{\left\| \frac{\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}}{\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}} \right\|} - \frac{\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}}{\left\| \frac{\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}}{\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}} \right\|} \right) \Pi_{\mathbf{P}, \epsilon \frac{j+1}{2}} \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) + \\
& + g^2 \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)^2} d^3 k - \frac{1}{\left\| \frac{\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}}{\phi_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}} \right\|} \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, g \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{\mathbf{k} (b(\mathbf{k}) + b^\dagger(\mathbf{k}))}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)} d^3 k \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)
\end{aligned}$$

On the left hand side of the equation, there is a quantity whose module is bigger than $C \cdot \left| \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} - \nabla E_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right|$ for $g \rightarrow 0$, where C is a positive constant that is uniform in j and converges to m for $g \rightarrow 0$. It is due to the result in point 1). On the right hand side, there is a quantity whose module is bounded by a g -dependent constant times $\epsilon^{\frac{j+1}{8}}$. Looking at the proof of theorem 2.3bis, the norm $\left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} - \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|$ is of order $\epsilon^{\frac{j+1}{8}}$ (the multiplicative constant, which is uniform in j , gets smaller by reducing g). Moreover the following bounds hold:

$$\begin{aligned}
& \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, g \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{k^i (b(\mathbf{k}) + b^\dagger(\mathbf{k}))}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)} d^3 k \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) = \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, g \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{k^i b^\dagger(\mathbf{k})}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)} d^3 k \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) = \\
& = g \int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{k^i}{\sqrt{2}|\mathbf{k}|^2 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)} \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, |\mathbf{k}|^{\frac{1}{2}} b^\dagger(\mathbf{k}) \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) d^3 k \leq \\
& \leq g \left(\int_{\epsilon \frac{j+2}{2}}^{\epsilon \frac{j+1}{2}} \frac{(k^i)^2}{2|\mathbf{k}|^4 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)^2} d^3 k \right)^{\frac{1}{2}} \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}}, H^{mes} \widehat{\phi}_{\mathbf{P}}^{\epsilon \frac{j+2}{2}} \right)^{\frac{1}{2}} \leq
\end{aligned}$$

$$\begin{aligned}
&\leq g \left(\int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{(k^i)^2}{2|\mathbf{k}|^4 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}\right)^2} d^3 k \right)^{\frac{1}{2}} \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}}, H^{mes} \widehat{\phi}_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} \right)^{\frac{1}{2}} \leq \\
&\leq g \left(\int_{\epsilon^{\frac{j+2}{2}}}^{\epsilon^{\frac{j+1}{2}}} \frac{(k^i)^2}{2|\mathbf{k}|^4 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}}\right)^2} d^3 k \right)^{\frac{1}{2}} \left(E_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} - \widehat{c}_{\mathbf{P}}(j+2) \right)^{\frac{1}{2}} \left(\widehat{\phi}_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}}, \widehat{\phi}_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} \right)^{\frac{1}{2}} \\
&\frac{\left\| \widehat{\Pi}_{\mathbf{P}, \epsilon^{\frac{j+2}{2}}}^i \widehat{\phi}_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} \right\|}{\left\| \widehat{\phi}_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} \right\|} \leq \sqrt{2m} \cdot \left(E_{\mathbf{P}}^{\epsilon^{\frac{j+2}{2}}} - c_{\mathbf{P}}(j+2) \right)^{\frac{1}{2}}, \quad \frac{\left\| \Pi_{\mathbf{P}, \epsilon^{\frac{j+1}{2}}}^i \phi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} \right\|}{\left\| \phi_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} \right\|} \leq \sqrt{2m} \cdot \left(E_{\mathbf{P}}^{\epsilon^{\frac{j+1}{2}}} - c_{\mathbf{P}}(j+1) \right)^{\frac{1}{2}}
\end{aligned}$$

Having established the bound $\left| \nabla E^{\epsilon^{\frac{j+1}{2}}}(\mathbf{P}) - \nabla E^{\epsilon^{\frac{j+2}{2}}}(\mathbf{P}) \right| < C(g) \cdot \epsilon^{\frac{j+1}{8}}$ for g sufficiently small, the j uniformity of $C(g)$ follows from the inductive procedure and the related tuning of the constant g . About the last point I do not give the details but a substantially analogous procedure is used in theorem 2.3. In conclusion the result is that for g less than a proper $\overline{g}$, the thesis of the point 2) of the lemma is proved jointly with theorem 2.3bis.

3)

The convergence of $\nabla E^{\sigma}(\mathbf{P})$ and of $\frac{1}{\|\tilde{\phi}_{\mathbf{P}}^{\sigma}\|^2} (\tilde{\phi}_{\mathbf{P}}^{\sigma}, \Pi_{\mathbf{P}, \sigma} \tilde{\phi}_{\mathbf{P}}^{\sigma})$ follows from the convergence proved in theorem 2.3bis. I will prove the second limit only.

I estimate the difference

$$\frac{1}{\|\tilde{\phi}_{\mathbf{P}}^{\sigma_2}\|^2} (\tilde{\phi}_{\mathbf{P}}^{\sigma_2}, \Pi_{\mathbf{P}, \sigma_2} \tilde{\phi}_{\mathbf{P}}^{\sigma_2}) - \frac{1}{\|\tilde{\phi}_{\mathbf{P}}^{\sigma_1}\|^2} (\tilde{\phi}_{\mathbf{P}}^{\sigma_1}, \Pi_{\mathbf{P}, \sigma_1} \tilde{\phi}_{\mathbf{P}}^{\sigma_1})$$

I recall the definition of $\tilde{\phi}_{\mathbf{P}}^{\sigma}$:

given σ between $\epsilon^{\frac{j+2}{2}}$ and $\epsilon^{\frac{j+3}{2}}$, it can be written as $\epsilon'^{\frac{j+2}{2}}$ where $\epsilon' = \sigma^{\frac{2}{j+2}} \{\epsilon' : \epsilon \geq \epsilon' \geq \epsilon\sqrt{\epsilon}\}$ and it can be defined $\tilde{\phi}_{\mathbf{P}}^{\sigma} \equiv \phi_{\mathbf{P}}^{\epsilon'(\sigma)^{\frac{j+2}{2}}}$, where we obtain $\phi_{\mathbf{P}}^{\epsilon'(\sigma)^{\frac{j+2}{2}}}$ by iteration starting from the cut-off ϵ' .

I rewrite $\tilde{\phi}_{\mathbf{P}}^{\sigma_2} - \tilde{\phi}_{\mathbf{P}}^{\sigma_1}$ in the following way

$$\tilde{\phi}_{\mathbf{P}}^{\sigma_2} - \tilde{\phi}_{\mathbf{P}}^{\sigma_1} = \tilde{\phi}_{\mathbf{P}}^{\sigma_2} - \phi_{\mathbf{P}}^{\epsilon_2(\sigma_2)\frac{l+2}{2}} + \phi_{\mathbf{P}}^{\epsilon_2(\sigma_2)\frac{l+2}{2}} - \phi_{\mathbf{P}}^{\epsilon_1(\sigma_1)\frac{m+2}{2}} + \phi_{\mathbf{P}}^{\epsilon_1(\sigma_1)\frac{m+2}{2}} - \tilde{\phi}_{\mathbf{P}}^{\sigma_1}$$

Now, fixed an arbitrarily small δ , there exist $l(\delta), m(\delta)$ sufficiently large and a phase $e^{i\eta(\delta)}$ such that

$$\left\| \frac{\phi_{\mathbf{P}}^{\epsilon_2(\sigma_2)\frac{l+2}{2}}}{\left\| \phi_{\mathbf{P}}^{\epsilon_2(\sigma_2)\frac{l+2}{2}} \right\|} - e^{i\eta(\delta)} \frac{\phi_{\mathbf{P}}^{\epsilon_1(\sigma_1)\frac{m+2}{2}}}{\left\| \phi_{\mathbf{P}}^{\epsilon_1(\sigma_1)\frac{m+2}{2}} \right\|} \right\| \leq \delta$$

This is substantially due to the convergence stated in the theorem 2.3bis and to the unicity of the ground state till there is a cut-off, by construction.

Then $\left\| \frac{\tilde{\phi}_{\mathbf{P}}^{\sigma_2}}{\left\| \tilde{\phi}_{\mathbf{P}}^{\sigma_2} \right\|} - e^{i\eta(\delta)} \frac{\tilde{\phi}_{\mathbf{P}}^{\sigma_1}}{\left\| \tilde{\phi}_{\mathbf{P}}^{\sigma_1} \right\|} \right\|$ is bounded by a quantity of order $\sigma_2^{\frac{1}{8}} + \sigma_1^{\frac{1}{8}} + \delta$.

Since

$$\frac{1}{\left\| \tilde{\phi}_{\mathbf{P}}^{\sigma_2} \right\|^2} \left(\tilde{\phi}_{\mathbf{P}}^{\sigma_2}, \Pi_{\mathbf{P}, \sigma_2} \tilde{\phi}_{\mathbf{P}}^{\sigma_2} \right) - \frac{1}{\left\| \tilde{\phi}_{\mathbf{P}}^{\sigma_1} \right\|^2} \left(\tilde{\phi}_{\mathbf{P}}^{\sigma_1}, \Pi_{\mathbf{P}, \sigma_1} \tilde{\phi}_{\mathbf{P}}^{\sigma_1} \right) = \frac{1}{\left\| \tilde{\phi}_{\mathbf{P}}^{\sigma_2} \right\|^2} \left(\tilde{\phi}_{\mathbf{P}}^{\sigma_2}, \Pi_{\mathbf{P}, \sigma_2} \tilde{\phi}_{\mathbf{P}}^{\sigma_2} \right) - \frac{1}{\left\| e^{i\eta} \tilde{\phi}_{\mathbf{P}}^{\sigma_1} \right\|^2} \left(e^{i\eta} \tilde{\phi}_{\mathbf{P}}^{\sigma_1}, \Pi_{\mathbf{P}, \sigma_1} e^{i\eta} \tilde{\phi}_{\mathbf{P}}^{\sigma_1} \right)$$

we can conclude that the limit exists by exploiting the bounds (from above) showed in point 2).

Calculation of $\widehat{H}_{\mathbf{P}, \epsilon}^{\frac{j+2}{2}}$.

$$\widehat{H}_{\mathbf{P}, \epsilon}^{\frac{j+2}{2}} = \frac{(\mathbf{P}_1 + \mathbf{P}_2)^2}{2m} + \frac{1}{2m} \left(\mathbf{P}^{mes} - g \int_{\epsilon}^{\kappa} \frac{\mathbf{k}(b(\mathbf{k}) + b^{\dagger}(\mathbf{k}))}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{\epsilon} \left(\frac{j+1}{2}(\mathbf{P}) \right) \right)} d^3 k + g^2 \int_{\epsilon}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1 - \widehat{\mathbf{k}} \cdot \nabla E^{\epsilon} \left(\frac{j+1}{2}(\mathbf{P}) \right) \right)^2} d^3 k \right)^2 +$$

$$\begin{aligned}
& -\frac{\mathbf{P}_2}{m} \cdot \left(\mathbf{P}^{mes} - g \int_{\epsilon}^{\kappa} \frac{\mathbf{k}(b(\mathbf{k})+b^\dagger(\mathbf{k}))}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)} d^3 k + g^2 \int_{\epsilon}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)^2} d^3 k \right) + \\
& + \int_0^\infty \left(|\mathbf{k}| - \mathbf{k} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k - g^2 \int_{\epsilon}^{\kappa} \frac{1}{2|\mathbf{k}|^2 \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)} d^3 k = \\
& = \frac{1}{2m} \left(\Pi_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} - g \int_{\epsilon}^{\kappa} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3 k + g^2 \int_{\epsilon}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)^2} d^3 k \right)^2 + \\
& - \frac{1}{m} \left(\frac{1}{\left\| \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|^2} \left(\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \Pi_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) + g^2 \int_{\epsilon}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)^2} d^3 k \right) \cdot \\
& \cdot \left(\Pi_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} - g \int_{\epsilon}^{\kappa} \frac{\mathbf{k}}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)} (b(\mathbf{k}) + b^\dagger(\mathbf{k})) d^3 k + g^2 \int_{\epsilon}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)^2} d^3 k \right) \\
& + \int_0^\infty \left(|\mathbf{k}| - \mathbf{k} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k - g^2 \int_{\epsilon}^{\kappa} \frac{1}{2|\mathbf{k}|^2 \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)} d^3 k = \\
& = \frac{1}{2m} \left(\Pi_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} - \frac{\left(\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \Pi_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right)}{\left\| \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|^2} - g \int_{\epsilon}^{\kappa} \frac{\mathbf{k}(b(\mathbf{k})+b^\dagger(\mathbf{k}))}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)} d^3 k + g^2 \int_{\epsilon}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)^2} d^3 k \right)^2 + \\
& + \int_0^\infty \left(|\mathbf{k}| - \mathbf{k} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P}) \right) b^\dagger(\mathbf{k}) b(\mathbf{k}) d^3 k - g^2 \int_{\epsilon}^{\kappa} \frac{1}{2|\mathbf{k}|^2 \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)} d^3 k + \\
& - \frac{1}{2m} \left(\frac{1}{\left\| \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right\|^2} \left(\phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}}, \Pi_{\mathbf{P}, \epsilon}^{\frac{j+1}{2}} \phi_{\mathbf{P}}^{\epsilon \frac{j+1}{2}} \right) + g^2 \int_{\epsilon}^{\kappa} \frac{\mathbf{k}}{2|\mathbf{k}|^3 \left(1-\widehat{\mathbf{k}} \cdot \nabla E^{\epsilon \frac{j+1}{2}}(\mathbf{P})\right)^2} d^3 k \right)^2 + \frac{\mathbf{P}^2}{2m}
\end{aligned}$$

APPENDIX B

Preliminary remarks

In the next lemmas I will consider an implicit hypothesis which is not proved in the spectral analysis but it is physically reasonable:

for $\mathbf{P} \in \Sigma$, there exists a constant $\frac{1}{m_r}$ (m_r means renormalized mass) such that the following inequalities hold, uniformly in $\sigma > 0$:

hypothesis B1

$$\frac{\partial E^\sigma(\mathbf{P})}{\partial |\mathbf{P}|} \geq \frac{|\mathbf{P}|}{m_r} \quad \text{and} \quad \frac{\partial^2 E^\sigma(\mathbf{P})}{\partial^2 |\mathbf{P}|} \geq \frac{1}{m_r}$$

Starting from this hypothesis, we obtain that the application $\mathbf{J}_\sigma : \mathbf{P} \rightarrow \nabla E^\sigma(\mathbf{P})$ is one to one and that the determinant of jacobian satisfies the inequality:

$$\det \mathbf{dJ}_\sigma = \frac{1}{|\mathbf{P}|^2} \left(\frac{\partial E^\sigma(\mathbf{P})}{\partial |\mathbf{P}|} \right)^2 \cdot \frac{\partial^2 E^\sigma(\mathbf{P})}{\partial^2 |\mathbf{P}|} \geq \frac{1}{m_r^3}$$

(I recall that the function $E^\sigma(\mathbf{P})$ is invariant under rotations and that it belongs to $C^\infty(R^3)$, see [2]) .

Then, given the region $O_{\nabla E^\sigma}$ and the corresponding $O_{\mathbf{P}} = \mathbf{J}_\sigma^{-1}(O_{\nabla E^\sigma})$ $O_{\mathbf{P}} \subset \Sigma$, we have the following relation between their volumes:

$$V_{O_{\mathbf{P}}} \leq m_r^3 V_{O_{\nabla E^\sigma}}$$

Remark on the notations

As in the previous chapters, I use the convention to generically call C the constants which are uniform in the variables we are treating. The bounds are intended from above, up to a different explicit warning.

Definition

As anticipated in the paragraph 4.1, the function $\chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, s)$ has to approximate the characteristic function of $\mathbf{J}_{\sigma_t}(\Gamma_i)$ for $s \rightarrow +\infty$ (where s is bigger or equal to the t ($\gg 1$) of the partition, the most general expression is $\chi_{\mathbf{v}_i}^{(t_1)}(\nabla E^{\sigma_{t_2}}, s)$ where the constraint is $s \geq t_1$). In particular, in order to approximate the region $\mathbf{J}_{\sigma_t}(\Gamma_i)$ from inside, I define $\chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, s) = \sum_l \chi_{\mathbf{v}_{l(i)}}^{l(t)}(\nabla E^{\sigma_t}, s)$ where $\text{supp}_{\nabla E^{\sigma_t}} \chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, s) \subset \text{supp} \mathbf{J}_{\sigma_t}(\Gamma_i)$ and where the $\chi_{\mathbf{v}_{l(i)}}^{l(t)}(\nabla E^{\sigma_t}, s)$ are constructed starting from the “model” function

$$\chi^{l(t)}(\mathbf{z}, s) = \chi_1^{l(t)}(z_1, s) \cdot \chi_2^{l(t)}(z_2, s) \cdot \chi_3^{l(t)}(z_3, s)$$

$$\chi_k^{l(t)}(z_k, s) \equiv \begin{cases} -s^{\frac{\delta}{2}} z_k + s^{\frac{\delta}{2}} \cdot \frac{1}{2s^{\frac{\delta}{6}}} & \text{for } -\frac{1}{s^{\frac{\delta}{2}}} + \frac{1}{2s^{\frac{\delta}{6}}} < z_k \leq \frac{1}{2s^{\frac{\delta}{6}}} \\ 1 & \text{for } -\frac{1}{2s^{\frac{\delta}{6}}} + \frac{1}{s^{\frac{\delta}{2}}} < z_k \leq \frac{1}{2s^{\frac{\delta}{6}}} - \frac{1}{s^{\frac{\delta}{2}}} \\ s^{\frac{\delta}{2}} z_k + s^{\frac{\delta}{2}} \cdot \frac{1}{2s^{\frac{\delta}{6}}} & \text{for } -\frac{1}{2s^{\frac{\delta}{6}}} < z_k \leq -\frac{1}{2s^{\frac{\delta}{6}}} + \frac{1}{s^{\frac{\delta}{2}}} \end{cases}$$

through a translation which sends the origin of the coordinates ∇E^{σ_t} space in $\mathbf{v}_{l(i)} \in \mathbf{J}_{\sigma_t}(\Gamma_i)$. Note that the support of $\mathbf{J}_{\sigma_t}(\Gamma_i)$ has a volume of order $\frac{1}{t^{3\epsilon}}$. In order to have a well defined $\chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, s)$ two requirements are necessary:

- the inequality $\delta > 6\epsilon$
 - a finite scale factor (related to m_r) for the variable z_k in the function $\chi_k^{l(t)}(z_k, s)$.
- To simplify the notations I will assume this factor equal to 1.

It comes out that the $\tilde{\chi}_{\mathbf{v}_{l(i)}}^{l(t)}(\mathbf{q}, s)$ have a behavior similar to

$$\tilde{\chi}^{l(t)}(\mathbf{q}, s) = \tilde{\chi}_1^{l(t)}(q_1, s) \cdot \tilde{\chi}_2^{l(t)}(q_2, s) \cdot \tilde{\chi}_3^{l(t)}(q_3, s)$$

$$\text{where } \tilde{\chi}_k^{l(t)}(q_k, s) \propto s^{\frac{\delta}{2}} \cdot \frac{\cos\left(q_k \left(\frac{1}{2s^{\frac{\delta}{6}}} - \frac{1}{s^{\frac{\delta}{2}}}\right)\right) - \cos\left(q_k \cdot \frac{1}{2s^{\frac{\delta}{6}}}\right)}{q_k^2}$$

and then

$$\int |\tilde{\chi}^{l(t)}(\mathbf{q}, s)| d^3q < C \cdot s^{3\frac{\delta}{2}} \quad \text{e} \quad \int_a^\infty |\tilde{\chi}^{l(t)}(\mathbf{q}, s)| d^3q < C \cdot \frac{1}{a} \cdot s^{\frac{\delta}{2}} \cdot s^\delta$$

from which

$$\int |\tilde{\chi}_{\mathbf{v}_i}^{l(t)}(\mathbf{q}, s)| d^3q \leq \sum_l \int |\tilde{\chi}_{\mathbf{v}_i}^{l(t)}(\mathbf{q}, s)| d^3q \leq C \cdot \frac{L^3}{t^{3\epsilon}} \cdot s^{3\frac{\delta}{6}} \cdot s^{3\frac{\delta}{2}} \leq C \cdot s^{2\delta}$$

$$\int_a^{+\infty} |\tilde{\chi}_{\mathbf{v}_i}^{l(t)}(\mathbf{q}, s)| d^3q \leq \sum_l \int_a^\infty |\tilde{\chi}_{\mathbf{v}_{l(i)}}^{l(t)}(q, s)| d^3q \leq C \cdot \frac{L^3}{t^{3\epsilon}} \cdot s^{3\frac{\delta}{6}} \frac{1}{a} \cdot s^{\frac{3\delta}{2}} < C \cdot \frac{1}{a} \cdot s^{2\delta}$$

Lemma B1

The norm $\left\| \left(1_{\Gamma_i}(\mathbf{P}) - \chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}(\mathbf{P}), s) \right) \psi_{i, \sigma_t}^{(t)} \right\|$ is bounded by a quantity of order $\frac{1}{s^{\frac{1}{12}}} \cdot t^{-\epsilon}$.

Proof

I define $\mathbf{J}_{\sigma_t}^{-1}(\hat{O}_{\nabla E^{\sigma_t}}^i) \equiv \left\{ \mathbf{P} \in \Sigma : \nabla E^{\sigma_t}(\mathbf{P}) \in \text{supp} \chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, s) \text{ and } \chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, s) \neq 1 \right\}$. Taking into account the definition of $\chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, s)$, the definition of the applica-

tion $\mathbf{J}_{\sigma_t} (\mathbf{J}_{\sigma_t} (\mathbf{P}) \propto \mathbf{P})$ and the hypothesis B1, the volume $\mathbf{J}_{\sigma_t}^{-1} (\widehat{O}_{\nabla E^{\sigma_t}}^i)$ is bounded by a quantity of order $\frac{1}{s^{\frac{2}{3}}} \cdot t^{-3\epsilon}$. On the other hand, the volume of the region $\text{supp} \mathbf{J}_{\sigma_t} (\Gamma_i) \setminus \text{supp}_{\nabla E^{\sigma_t}} \chi_{\mathbf{v}_i}^{(t)} (\nabla E^{\sigma_t}, s)$ is bounded by a quantity of order $\frac{1}{s^{\frac{2}{3}}} \cdot t^{-2\epsilon}$. Therefore:

$$\begin{aligned} & \left\| \left(1_{\Gamma_i} (\mathbf{P}) - \chi_{\mathbf{v}_i}^{(t)} (\nabla E^{\sigma_t} (\mathbf{P}), s) \right) \psi_{i, \sigma_t}^{(t)} \right\|^2 = \int_{\Gamma_i} \left| 1_{\Gamma_i} (\mathbf{P}) - \chi_{\mathbf{v}_i}^{(t)} (\nabla E^{\sigma_t}, s) \right|^2 |G(\mathbf{P})|^2 \|\psi_{\mathbf{P}, \sigma_t}\|^2 d^3 P \leq \\ & \leq C \cdot \frac{1}{s^{\frac{2}{3}}} \cdot t^{-2\epsilon} \end{aligned}$$

from which the thesis follows.

Lemma B2

In the constructive hypothesis fixed at the beginning of chapter 4, we have

$$\begin{aligned} & \left\| \int \widetilde{\chi}_{\mathbf{v}_i}^{(t)} (\mathbf{q}, s) \left(e^{-i\mathbf{q} \cdot \nabla E^{\sigma_t}} - e^{-i\mathbf{q} \cdot \frac{\mathbf{x}}{s}} \right) d^3 q e^{-iE^{\sigma_t}(\mathbf{P})s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} \right\| \leq \\ & \leq C \cdot s^{-\frac{1}{112}} \cdot s^{2\delta - \frac{3\epsilon}{2}} \cdot (\ln(\sigma_t))^2 \end{aligned}$$

(C is the same constant for all the cells and it is uniform in the partitions).

Proof

$$\begin{aligned} & \left\| \int \widetilde{\chi}_{\mathbf{v}_i}^{(t)} (\mathbf{q}, s) \left(e^{-i\mathbf{q} \cdot \nabla E^{\sigma_t}} - e^{-i\mathbf{q} \cdot \frac{\mathbf{x}}{s}} \right) d^3 q e^{-iE^{\sigma_t}(\mathbf{P})s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} \right\| = \\ & = \left\| e^{iE^{\sigma_t}(\mathbf{P})s} \int \widetilde{\chi}_{\mathbf{v}_i}^{(t)} (\mathbf{q}, s) \left(e^{-i\mathbf{q} \cdot \nabla E^{\sigma_t}} - e^{-i\mathbf{q} \cdot \frac{\mathbf{x}}{s}} \right) d^3 q e^{-iE^{\sigma_t}(\mathbf{P})s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} \right\| = \\ & = \left\| \int \widetilde{\chi}_{\mathbf{v}_i}^{(t)} (\mathbf{q}, s) \left[\left(e^{-i\mathbf{q} \cdot \nabla E^{\sigma_t}} - e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} \right) - e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} \left(e^{-i\mathbf{q} \cdot \frac{\mathbf{x}}{s}} - 1 \right) \right] d^3 q e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} \right\| \leq \\ & \leq \left\| \int \widetilde{\chi}_{\mathbf{v}_i}^{(t)} (\mathbf{q}, s) \left(e^{-i\mathbf{q} \cdot \nabla E^{\sigma_t}} - e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} \right) d^3 q e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} \right\| + \quad \textbf{i)} \\ & + \left\| \int \widetilde{\chi}_{\mathbf{v}_i}^{(t)} (\mathbf{q}, s) e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} \left(e^{-i\mathbf{q} \cdot \frac{\mathbf{x}}{s}} - 1 \right) d^3 q e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} \right\| \quad \textbf{ii)} \end{aligned}$$

$$\textbf{i)} \left\| \int \widetilde{\chi}_{\mathbf{v}_i}^{(t)} (\mathbf{q}, s) \left(e^{-i\mathbf{q} \cdot \nabla E^{\sigma_t}} - e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} \right) d^3 q e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t}, s)} \psi_{i, \sigma_t}^{(t)} \right\| \leq$$

$$\leq C \cdot \left\| \psi_{i,\sigma}^{(t)} \right\| \cdot \int_{\frac{1}{s^{34}}}^{+\infty} \left| \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \right| d^3 q + C \cdot \left\| \psi_{i,\sigma}^{(t)} \right\| \cdot \frac{\int_0^{s^{\frac{1}{34}}} \left| \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \right| d^3 q}{s^{\frac{1}{32}}}$$

It follows for these reasons:

- being the energy differentiable:
 $sE^{\sigma_t}(\mathbf{P}) - sE^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}) = -\mathbf{q} \cdot \nabla E^{\sigma_t}(\mathbf{P}')$ where $\mathbf{P}'$ is such that $|\mathbf{P} - \mathbf{P}'| \leq \left| \frac{\mathbf{a}}{s} \right|$
- for the lemma 3.3:
 $|\nabla E^{\sigma_t}(\mathbf{P}) - \nabla E^{\sigma_t}(\mathbf{P}')| \leq C \cdot |\mathbf{P} - \mathbf{P}'|^{\frac{1}{16}} \leq C \cdot \left| \frac{\mathbf{a}}{s} \right|^{\frac{1}{16}}$
- $\int \left| \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \right| d^3 q \leq C \cdot s^{2\delta}$
 $\int_a^{+\infty} \left| \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \right| d^3 q < C \cdot \frac{1}{a} \cdot s^{2\delta}$

Then the term **i**) is surely bounded by a quantity of order $\frac{s^{2\delta}}{s^{\frac{1}{34}}}$.

ii)

$$\begin{aligned} & \left\| \int \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} \cdot \left(e^{-i\mathbf{q} \cdot \frac{\mathbf{x}}{s}} - 1 \right) d^3 q e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t, s})} \psi_{i,\sigma_t}^{(t)} \right\| \leq \\ & \leq \left\| \int_0^{s^{\frac{1}{20}}} \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} \cdot \left(e^{-i\mathbf{q} \cdot \frac{\mathbf{x}}{s}} - 1 \right) d^3 q e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t, s})} \psi_{i,\sigma_t}^{(t)} \right\| + \\ & + \left\| \int_{s^{\frac{1}{20}}}^{+\infty} \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} \cdot \left(e^{-i\mathbf{q} \cdot \frac{\mathbf{x}}{s}} - 1 \right) d^3 q e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E^{\sigma_t, s})} \psi_{i,\sigma_t}^{(t)} \right\| \end{aligned} \quad (\text{b2})$$

first term of (b2)

$$\begin{aligned} & \left\| \int_0^{s^{\frac{1}{20}}} \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} \cdot \left(e^{-i\mathbf{q} \cdot \frac{\mathbf{x}}{s}} - 1 \right) d^3 q e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t, s})} \psi_{i,\sigma_t}^{(t)} \right\| = \\ & = \left\| \int_0^{s^{\frac{1}{20}}} \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \int_{\Gamma_i} e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} \cdot \left(e^{-i\mathbf{q} \cdot \frac{\mathbf{x}}{s}} - 1 \right) G(\mathbf{P}) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t, s})} \psi_{\mathbf{P}, \sigma_t} d^3 P d^3 q \right\| = \\ & = \left\| \int_0^{s^{\frac{1}{20}}} \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \int_{\Gamma_i} e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} e^{-i\mathbf{q} \cdot \frac{\mathbf{x}}{s}} G(\mathbf{P}) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t, s})} \psi_{\mathbf{P}, \sigma_t} d^3 P d^3 q + \right. \\ & \quad \left. - \int_0^{s^{\frac{1}{20}}} \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \int_{\Gamma_i} e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} G(\mathbf{P}) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t, s})} \psi_{\mathbf{P}, \sigma_t} d^3 P d^3 q \right\| = \\ & = \left\| \int_0^{s^{\frac{1}{20}}} \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \int_{\Gamma_i} e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{a}}{s}))s} e^{-i\mathbf{q} \cdot \frac{\mathbf{x}}{s}} G(\mathbf{P}) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t, s})} \psi_{\mathbf{P}, \sigma_t} d^3 P d^3 q + \right. \end{aligned}$$

$$\begin{aligned}
& - \int_0^{s^{\frac{1}{20}}} \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \int_{\Gamma_i} e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{q}}{s}))s} G(\mathbf{P}) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P} - \frac{\mathbf{q}}{s}}^{\sigma_t}, s)} \psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t} d^3 P d^3 q + \\
& + \int_0^{s^{\frac{1}{20}}} \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \int_{\Gamma_i} e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{q}}{s}))s} G(\mathbf{P}) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P} - \frac{\mathbf{q}}{s}}^{\sigma_t}, s)} \psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t} d^3 P d^3 q + \\
& - \int_0^{s^{\frac{1}{20}}} \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \int_{\Gamma_i} e^{i(E^{\sigma_t}(\mathbf{P} - \frac{\mathbf{q}}{s}) - E^{\sigma_t}(\mathbf{P}))s} G(\mathbf{P} - \frac{\mathbf{q}}{s}) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P} - \frac{\mathbf{q}}{s}}^{\sigma_t}, s)} \psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t} d^3 P d^3 q + \\
& + \int_0^{s^{\frac{1}{20}}} \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \int_{\Gamma_i} e^{i(E^{\sigma_t}(\mathbf{P} - \frac{\mathbf{q}}{s}) - E^{\sigma_t}(\mathbf{P}))s} G(\mathbf{P} - \frac{\mathbf{q}}{s}) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P} - \frac{\mathbf{q}}{s}}^{\sigma_t}, s)} \psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t} d^3 P d^3 q + \\
& - \int_0^{s^{\frac{1}{20}}} \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \int_{\Gamma_i} e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{q}}{s}))s} G(\mathbf{P}) e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t}, s)} \psi_{\mathbf{P}, \sigma_t} d^3 P d^3 q \Big\| \leq \\
& \leq \int_0^{s^{\frac{1}{20}}} \left| \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \right| \left\{ \int_{\Gamma_i} |G(\mathbf{P})|^2 \left\| e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t}, s)} I_{\mathbf{P}}(\psi_{\mathbf{P}, \sigma_t}) - e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P} - \frac{\mathbf{q}}{s}}^{\sigma_t}, s)} I_{\mathbf{P} - \frac{\mathbf{q}}{s}}(\psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t}) \right\|_F^2 d^3 P \right\}^{\frac{1}{2}} d^3 q + \\
& + \int_0^{s^{\frac{1}{20}}} \left| \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \right| \left\{ \int_{\Gamma_i} \left| G(\mathbf{P} - \frac{\mathbf{q}}{s}) e^{i(E^{\sigma_t}(\mathbf{P} - \frac{\mathbf{q}}{s}) - E^{\sigma_t}(\mathbf{P}))s} - G(\mathbf{P}) e^{i(E^{\sigma_t}(\mathbf{P}) - E^{\sigma_t}(\mathbf{P} + \frac{\mathbf{q}}{s}))s} \right|^2 \left\| I_{\mathbf{P} - \frac{\mathbf{q}}{s}}(\psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t}) \right\|_F^2 d^3 P \right\}^{\frac{1}{2}} d^3 q + \\
& + \int_0^{s^{\frac{1}{20}}} \left| \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \right| \left\{ \int_{O_{\frac{\mathbf{q}}{s}}} |G(\mathbf{P})|^2 \|I_{\mathbf{P}}(\psi_{\mathbf{P}, \sigma_t})\|_F^2 d^3 P \right\}^{\frac{1}{2}} d^3 q
\end{aligned}
\tag{b3.1}$$

$$\tag{b3.2}$$

$$\tag{b3.3}$$

bound of the term (b3.1)

$$\begin{aligned}
& \left\| e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t}, s)} I_{\mathbf{P}}(\psi_{\mathbf{P}, \sigma_t}) - e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P} - \frac{\mathbf{q}}{s}}^{\sigma_t}, s)} I_{\mathbf{P} - \frac{\mathbf{q}}{s}}(\psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t}) \right\|_F \leq \\
& \leq \left\| e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t}, s)} I_{\mathbf{P}}(\psi_{\mathbf{P}, \sigma_t}) - e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t}, s)} I_{\mathbf{P} - \frac{\mathbf{q}}{s}}(\psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t}) \right\|_F + \\
& + \left\| e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t}, s)} I_{\mathbf{P} - \frac{\mathbf{q}}{s}}(\psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t}) - e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P} - \frac{\mathbf{q}}{s}}^{\sigma_t}, s)} I_{\mathbf{P} - \frac{\mathbf{q}}{s}}(\psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t}) \right\|_F \leq \\
& \leq \left\| I_{\mathbf{P}}(W_{\sigma_t}^{b\dagger}(\nabla E_{\mathbf{P}}^{\sigma_t}) W_{\sigma_t}^b(\nabla E_{\mathbf{P}}^{\sigma_t}) \psi_{\mathbf{P}, \sigma_t}) - I_{\mathbf{P} - \frac{\mathbf{q}}{s}}(W_{\sigma_t}^{b\dagger}(\nabla E_{\mathbf{P} - \frac{\mathbf{q}}{s}}^{\sigma_t}) W_{\sigma_t}^b(\nabla E_{\mathbf{P} - \frac{\mathbf{q}}{s}}^{\sigma_t}) \psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t}) \right\|_F + \\
& + \left\| e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t}, s)} - e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P} - \frac{\mathbf{q}}{s}}^{\sigma_t}, s)} \right\| \left\| I_{\mathbf{P} - \frac{\mathbf{q}}{s}}(\psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t}) \right\|_F \leq \\
& \leq \left\| I_{\mathbf{P}}(W_{\sigma_t}^b(\nabla E_{\mathbf{P}}^{\sigma_t}) \psi_{\mathbf{P}, \sigma_t}) - I_{\mathbf{P} - \frac{\mathbf{q}}{s}}(W_{\sigma_t}^b(\nabla E_{\mathbf{P} - \frac{\mathbf{q}}{s}}^{\sigma_t}) \psi_{\mathbf{P} - \frac{\mathbf{q}}{s}, \sigma_t}) \right\|_F +
\end{aligned}$$

$$\begin{aligned}
& + \left\| I_{\mathbf{P}-\frac{\mathbf{a}}{s}} \left(\left(W_{\sigma_t}^{b^\dagger} (\nabla E_{\mathbf{P}}^{\sigma_t}) - W_{\sigma_t}^{b^\dagger} (\nabla E_{\mathbf{P}-\frac{\mathbf{a}}{s}}^{\sigma_t}) \right) W_{\sigma_t}^b (\nabla E_{\mathbf{P}-\frac{\mathbf{a}}{s}}^{\sigma_t}) \psi_{\mathbf{P}-\frac{\mathbf{a}}{s}, \sigma_t} \right) \right\|_F + \\
& + \left| e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t}, s)} - e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}-\frac{\mathbf{a}}{s}}^{\sigma_t}, s)} \right| \left\| I_{\mathbf{P}-\frac{\mathbf{a}}{s}} (\psi_{\mathbf{P}-\frac{\mathbf{a}}{s}, \sigma_t}) \right\|_F
\end{aligned}$$

The norm $\left\| I_{\mathbf{P}} (W_{\sigma_t}^b (\nabla E_{\mathbf{P}}^{\sigma_t}) \psi_{\mathbf{P}, \sigma_t}) - I_{\mathbf{P}-\frac{\mathbf{a}}{s}} (W_{\sigma_t}^b (\nabla E_{\mathbf{P}-\frac{\mathbf{a}}{s}}^{\sigma_t}) \psi_{\mathbf{P}-\frac{\mathbf{a}}{s}, \sigma_t}) \right\|_F$ is bounded by a quantity of order $\left(\frac{|\mathbf{q}|}{s}\right)^{\frac{1}{32}}$ (see theorem 3.4).

The norm of

$$\left\| I_{\mathbf{P}-\frac{\mathbf{a}}{s}} \left(\left(W_{\sigma_t}^{b^\dagger} (\nabla E_{\mathbf{P}}^{\sigma_t}) - W_{\sigma_t}^{b^\dagger} (\nabla E_{\mathbf{P}-\frac{\mathbf{a}}{s}}^{\sigma_t}) \right) W_{\sigma_t}^b (\nabla E_{\mathbf{P}-\frac{\mathbf{a}}{s}}^{\sigma_t}) \psi_{\mathbf{P}-\frac{\mathbf{a}}{s}, \sigma_t} \right) \right\|_F \quad (\text{b4})$$

can be estimated by the norm of

$$g \int_{\sigma_t}^{\kappa} \frac{\widehat{\mathbf{k}} \cdot (\nabla E^{\sigma_t}(\mathbf{P}) - \nabla E^{\sigma_t}(\mathbf{P} - \frac{\mathbf{a}}{s}))}{|\mathbf{k}| (1 - \widehat{\mathbf{k}} \cdot \nabla E^{\sigma_t}(\mathbf{P})) (1 - \widehat{\mathbf{k}} \cdot \nabla E^{\sigma_t}(\mathbf{P} - \frac{\mathbf{a}}{s}))} (b(\mathbf{k}) - b^\dagger(\mathbf{k})) \frac{d^3 k}{\sqrt{2}|\mathbf{k}|} I_{\mathbf{P}-\frac{\mathbf{a}}{s}} (W_{\sigma_t}^b (\nabla E^{\sigma_t}(\mathbf{P} - \frac{\mathbf{a}}{s})) \psi_{\mathbf{P}-\frac{\mathbf{a}}{s}, \sigma_t})$$

then it is substantially the product of the following quantities:

$$\begin{aligned}
& \bullet \left(\int_{\sigma_t}^{\kappa} \left(\frac{g \widehat{\mathbf{k}} \cdot (\nabla E^{\sigma_t}(\mathbf{P}) - \nabla E^{\sigma_t}(\mathbf{P} - \frac{\mathbf{a}}{s}))}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} (1 - \widehat{\mathbf{k}} \cdot \nabla E^{\sigma_t}(\mathbf{P})) (1 - \widehat{\mathbf{k}} \cdot \nabla E^{\sigma_t}(\mathbf{P} - \frac{\mathbf{a}}{s}))} \right)^2 d^3 k \right)^{\frac{1}{2}} \\
& \text{it is bounded by } C \cdot \left| \nabla E^{\sigma_t}(\mathbf{P} - \frac{\mathbf{a}}{s}) - \nabla E^{\sigma_t}(\mathbf{P}) \right| \cdot (\ln(\sigma_t))^{\frac{1}{2}} \leq \\
& \leq C \cdot \left| s^{-\frac{19}{20}} \right|^{\frac{1}{16}} \cdot (\ln(\sigma_t))^{\frac{1}{2}} \text{ (see lemma 3.3);} \\
& \bullet \left(\int_{\sigma_t}^{\kappa} \left\| b(\mathbf{k}) I_{\mathbf{P}-\frac{\mathbf{a}}{s}} (W_{\sigma_t}^b (\nabla E^{\sigma_t}(\mathbf{P} - \frac{\mathbf{a}}{s})) \psi_{\mathbf{P}-\frac{\mathbf{a}}{s}, \sigma_t}) \right\|_F^2 d^3 k \right)^{\frac{1}{2}} = \\
& = \left(\int_{\sigma_t}^{\kappa} \left\| I_{\mathbf{P}-\frac{\mathbf{a}}{s}} \left(W_{\sigma_t}^b (\nabla E^{\sigma_t}(\mathbf{P} - \frac{\mathbf{a}}{s})) \left(b(\mathbf{k}) + \frac{g \chi_{\sigma_t}^{\kappa}(\mathbf{k})}{\sqrt{2}|\mathbf{k}|^{\frac{3}{2}} (1 - \widehat{\mathbf{k}} \cdot \nabla E_{\mathbf{P}-\frac{\mathbf{a}}{s}}^{\sigma_t})} \right) \psi_{\mathbf{P}+\frac{\mathbf{a}}{s}, \sigma_t} \right) \right\|_F^2 d^3 k \right)^{\frac{1}{2}}
\end{aligned}$$

by using techniques like in [2], it is possible to give a bound with a uniform constant in t and in s . For our purposes, a bound of order $(\ln(\sigma_t))^{\frac{1}{2}}$ (uniform in s) is sufficient. For $\mathbf{P} \in \Sigma$, it comes from the following formula obtained in [2]

$$b(\mathbf{k}) \psi_{\mathbf{P}, \sigma_t} = \frac{g}{\sqrt{2|\mathbf{k}|}} \left(\frac{1}{E^{\sigma_t}(\mathbf{P}) - |\mathbf{k}| - H_{\mathbf{P}-\mathbf{k}, \sigma_t}} \right) \psi_{\mathbf{P}, \sigma_t}$$

For $s \leq \sigma_t^{-\frac{40}{39}}$, the module of $\left| e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t}, s)} - e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}+\frac{\mathbf{a}}{s}}^{\sigma_t}, s)} \right|$ can be estimated by the module of the difference of the exponents:

$$\begin{aligned} & \left| \int_1^s \left\{ g^2 \int_{\sigma_t}^{\sigma_\tau^L} \frac{\cos(\mathbf{k} \cdot \nabla E_{\mathbf{P}}^{\sigma_t} \tau - |\mathbf{k}| \tau) - \cos(\mathbf{k} \cdot \nabla E_{\mathbf{P}-\frac{\mathbf{a}}{s}}^{\sigma_t} \tau - |\mathbf{k}| \tau)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| \right\} d\tau \right| = \\ & = \left| 2 \int_1^s \left\{ g^2 \int_{\sigma_t}^{\sigma_\tau^L} \frac{\sin\left(\frac{\mathbf{k} \cdot (\nabla E_{\mathbf{P}}^{\sigma_t} - \nabla E_{\mathbf{P}-\frac{\mathbf{a}}{s}}^{\sigma_t}) \tau}{2}\right) \sin\left(\frac{\mathbf{k} \cdot (\nabla E_{\mathbf{P}}^{\sigma_t} + \nabla E_{\mathbf{P}-\frac{\mathbf{a}}{s}}^{\sigma_t}) \tau - 2|\mathbf{k}| \tau}{2}\right)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| \right\} d\tau \right| \leq \\ & \leq C \cdot \left| \nabla E_{\mathbf{P}}^{\sigma_t} - \nabla E_{\mathbf{P}-\frac{\mathbf{a}}{s}}^{\sigma_t} \right| \cdot \int_1^s (\sigma_\tau^L)^2 \cdot \tau d\tau \leq C \cdot \left| \frac{\mathbf{a}}{s} \right|^{\frac{1}{16}} \cdot \int_1^s \tau^{-\frac{19}{20}} d\tau \leq \\ & \leq C \cdot \left| \frac{s^{\frac{1}{20}}}{s} \right|^{\frac{1}{16}} \cdot s^{\frac{1}{20}} < C \cdot s^{-\frac{1}{112}} \end{aligned}$$

An analogous estimate holds for $s > \sigma_t^{-\frac{40}{39}}$.

Summarizing, the term (b3.1) is bounded by

$$\begin{aligned} & C \cdot \left(s^{-\frac{19}{20}} \right)^{\frac{1}{32}} \cdot s^{2\delta} \cdot t^{-\frac{3\epsilon}{2}} + C \cdot \left(s^{-\frac{19}{20}} \right)^{\frac{1}{16}} \cdot s^{2\delta} \cdot t^{-\frac{3\epsilon}{2}} \cdot |\ln \sigma_t| + C \cdot s^{-\frac{1}{112}} \cdot s^{2\delta} \cdot t^{-\frac{3\epsilon}{2}} \leq \\ & \leq C \cdot s^{-\frac{1}{112}} \cdot s^{2\delta} \cdot t^{-\frac{3\epsilon}{2}} \cdot |\ln \sigma_t| \end{aligned}$$

bound of the term (b3.2)

Being $G \in C_0^1(R^3 \setminus 0)$ and for the lemma 3.3 applied to $e^{i(E^{\sigma_t}(\mathbf{P}-\frac{\mathbf{a}}{s})-E^{\sigma_t}(\mathbf{P}))s} - e^{i(E^{\sigma_t}(\mathbf{P})-E^{\sigma_t}(\mathbf{P}+\frac{\mathbf{a}}{s}))s}$ there is a bounding with the quantity

$$C \cdot |\mathbf{q}| \cdot \left(\frac{|\mathbf{q}|}{s} \right)^{\frac{1}{16}} \cdot s^{2\delta} \cdot t^{-\frac{3\epsilon}{2}} \leq C \cdot s^{-\frac{1}{20}} \cdot \left(s^{-\frac{19}{20}} \right)^{\frac{1}{16}} \cdot s^{2\delta} \cdot t^{-\frac{3\epsilon}{2}} \leq C \cdot s^{2\delta} \cdot s^{-\frac{1}{112}} \cdot t^{-\frac{3\epsilon}{2}}$$

bound of the term (b3.3)

Starting from a difference between volumes, the expression (b3.3) is bounded by a quantity of order $\left(\frac{|\mathbf{q}|}{s} \right)^{\frac{1}{2}} \cdot s^{2\delta} \cdot t^{-\epsilon} \leq s^{-\frac{19}{40}} \cdot s^{2\delta} \cdot t^{-\epsilon}$

second term of the (b2)

It is bounded by $\left\| \psi_{i,\sigma_t}^{(t)} \right\| \cdot \int_{s^{\frac{1}{20}}}^{+\infty} \left| \tilde{\chi}_{\mathbf{v}_i}^{(t)}(\mathbf{q}, s) \right| d^3q$ then by

$$C \cdot s^{-\frac{1}{20}} \cdot s^{2\delta} \cdot t^{-\frac{3\epsilon}{2}}$$

In conclusion the sum of the terms **i)** and **ii)** is bounded by

$$C \cdot s^{-\frac{1}{112}} \cdot s^{2\delta} \cdot |\ln \sigma_t| \cdot t^{-\frac{3\epsilon}{2}}$$

having assumed $\frac{\epsilon}{2} \leq \frac{19}{40} - \frac{1}{112}$.

Corollary B2

From the previous proof it follows that for $s > t \gg 1$ the norm of

$$\left(g^2 \int_{\sigma_t \cdot s}^{s \cdot s^{-\frac{39}{40}}} \int \frac{\cos(\mathbf{q} \cdot \frac{\mathbf{x}}{s} - |\mathbf{q}|)}{(1 - \widehat{\mathbf{q}} \cdot \mathbf{v}_i)} d\Omega \frac{d|\mathbf{q}|}{s} - g^2 \int_{\sigma_t \cdot s}^{s \cdot s^{-\frac{39}{40}}} \int \frac{\cos(\mathbf{q} \cdot \nabla E^{\sigma_t} - |\mathbf{q}|)}{(1 - \widehat{\mathbf{q}} \cdot \mathbf{v}_i)} d\Omega \frac{d|\mathbf{q}|}{s} \right) e^{-iE^{\sigma_t}s} e^{i\gamma_{\sigma_t}(\mathbf{v}_i, \nabla E_{\mathbf{P}}^{\sigma_t}, s)} \psi_{i,\sigma_t}^{(t)}$$

is surely bounded by a quantity of order $s^{-1} \cdot s^{-\frac{1}{112}} \cdot s^{2\delta} \cdot |\ln \sigma_t| \cdot t^{-\frac{3\epsilon}{2}}$.

Lemma B3

I study the function $\varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, t)$ ($t \gg 1$) where $\sigma_t = \frac{1}{t^\alpha}$ and $\alpha > 0$

$$\varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, t) = g^2 \int_{\sigma_t}^{\kappa_1} \int \frac{\cos(\mathbf{k} \cdot \mathbf{x} - |\mathbf{k}|t)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| = g^2 \int \frac{\sin(\kappa_1 \widehat{\mathbf{k}} \cdot \mathbf{x} - \kappa_1 t) - \sin(\sigma_t \widehat{\mathbf{k}} \cdot \mathbf{x} - \sigma_t t)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i) \cdot (\widehat{\mathbf{k}} \cdot \mathbf{x} - t)} d\Omega$$

Observation 1

I analyze the behavior of $\varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, t)$ for $\mathbf{x} \in R^3$.

region $\mathbf{x}$: $|\mathbf{x}| < (1 - \eta)t$, $0 < \eta < 1$:

$$\begin{aligned} \left| g^2 \int \frac{\sin(\kappa_1 \widehat{\mathbf{k}} \cdot \mathbf{x} - \kappa_1 t) - \sin(\sigma_t \widehat{\mathbf{k}} \cdot \mathbf{x} - \sigma_t t)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i) \cdot (\widehat{\mathbf{k}} \cdot \mathbf{x} - t)} \cdot d\Omega \right| &\leq \int \left| g^2 \frac{2}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i) \cdot (\widehat{\mathbf{k}} \cdot \mathbf{x} - t)} \right| d\Omega \leq \\ &\leq \frac{1}{\eta t} \int \left| g^2 \frac{2}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} \right| d\Omega \end{aligned}$$

region $\mathbf{x}$: $(1 - \eta)t < |\mathbf{x}|$, $0 < \eta < 1$:

$$\varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, t) = g^2 \int_{\sigma_t}^{\kappa_1} \int \frac{\cos(\mathbf{k} \cdot \mathbf{x} - |\mathbf{k}|t)}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| = g^2 \int_{\sigma_t}^{\kappa_1} \int \cos(\mathbf{k} \cdot \mathbf{x} - |\mathbf{k}|t) \cdot \xi(\widehat{\mathbf{k}}, \mathbf{v}_i) d\Omega d|\mathbf{k}|$$

where $\xi(\widehat{\mathbf{k}}, \mathbf{v}_i) = \xi(\theta, \varphi, \mathbf{v}_i) \equiv \frac{1}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)}$. Note that being $|\mathbf{v}_i| \leq v^{max} < 1$

$$\exists \quad M \geq 0, M' \geq 0 \implies \left| \xi(\widehat{\mathbf{k}}, \mathbf{v}_i) \right| < M \text{ and } \left| \frac{d}{d \cos \theta} [\xi(\theta, \varphi, \mathbf{v}_i)] \right| < M';$$

I integrate by parts with respect to $d \cos \theta$

$$\begin{aligned} g^2 \int_{\sigma_t}^{\kappa_1} \int \frac{e^{i(\mathbf{k} \cdot \mathbf{x} - |\mathbf{k}|t)}}{(1 - \widehat{\mathbf{k}} \cdot \mathbf{v}_i)} d\Omega d|\mathbf{k}| &= \\ &= \int_{\sigma_t}^{\kappa_1} \int \frac{e^{-i(|\mathbf{k}| \cdot |\mathbf{x}| + |\mathbf{k}|t)}}{|\mathbf{x}|} \cdot \frac{\xi(\pi, \varphi, \mathbf{v}_i)}{i|\mathbf{k}|} d|\mathbf{k}| d\varphi - \int_{\sigma_t}^{\kappa_1} \int \frac{e^{i(|\mathbf{k}| \cdot |\mathbf{x}| - |\mathbf{k}|t)}}{|\mathbf{x}|} \cdot \frac{\xi(0, \varphi, \mathbf{v}_i)}{i|\mathbf{k}|} d|\mathbf{k}| d\varphi + \\ &- \int_{\sigma_t}^{\kappa_1} \int \frac{e^{i(\mathbf{k} \cdot \mathbf{x} - |\mathbf{k}|t)}}{|\mathbf{x}|} \cdot \frac{1}{i|\mathbf{k}|} \frac{d}{d \cos \theta} [\xi(\theta, \varphi, \mathbf{v}_i)] d|\mathbf{k}| d\Omega \end{aligned} \tag{b5}$$

this is bounded by a quantity of order $\frac{|\ln \sigma_t|}{t - \eta t}$.

In conclusion there exists a constant C , that is uniform in $\mathbf{v}_i$ belonging to the region $|\mathbf{v}_i| < v^{max} < 1$, such that $\forall \mathbf{x} \in R^3 \quad |\varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, t)| \leq C \cdot \frac{|\ln \sigma_t|}{t - \eta t}$. (b6)

Observation 2

I analyze the behavior of $\varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, t)$ for $(1 - \eta')t < |\mathbf{x}| < (1 - \eta)t$, where $0 < \eta < \eta' < 1$. I study the term (b5) (the other term has an analogous behavior)

$$\begin{aligned} & \int \frac{\kappa_1}{t^\alpha} \int \frac{e^{-i(|\mathbf{k}| \cdot |\mathbf{x}| + |\mathbf{k}|t)}}{|\mathbf{x}|} \cdot \frac{\xi(\pi, \varphi, \mathbf{v}_i)}{i|\mathbf{k}|} d|\mathbf{k}| d\varphi - \int \frac{\kappa_1}{t^\alpha} \int \frac{e^{i(|\mathbf{k}| \cdot |\mathbf{x}| - |\mathbf{k}|t)}}{|\mathbf{x}|} \cdot \frac{\xi(0, \varphi, \mathbf{v}_i)}{i|\mathbf{k}|} d|\mathbf{k}| d\varphi + \\ & - \int \frac{\kappa_1}{t^\alpha} \int \frac{e^{i(\mathbf{k} \cdot \mathbf{x} - |\mathbf{k}|t)}}{|\mathbf{x}|} \cdot \frac{1}{i|\mathbf{k}|} \frac{d}{d \cos \theta} [\xi(\theta, \varphi, \mathbf{v}_i)] d|\mathbf{k}| d\Omega \end{aligned} \quad (\text{b7})$$

By hypothesis, the following inequalities hold:

$$\begin{aligned} |\mathbf{x}| + t &> t \\ |\mathbf{x}| - t &< (1 - \eta)t - t \Rightarrow |\mathbf{x}| - t < -\eta t \Rightarrow ||\mathbf{x}| - t| > \eta t \\ |\mathbf{x}| &> (1 - \eta')t \end{aligned}$$

I consider, for example, the first term of the (b7) (the other ones have an analogous behavior) :

$$-i(|\mathbf{x}| + t) \cdot \int \frac{\kappa_1}{t^\alpha} \int \frac{e^{-i|\mathbf{k}| \cdot (|\mathbf{x}| + t)}}{|\mathbf{x}|} \cdot \frac{\xi(\pi, \varphi, \mathbf{v}_i)}{i|\mathbf{k}|} d|\mathbf{k}| d\varphi = \int \frac{\kappa_1}{t^\alpha} \int \frac{d}{d|\mathbf{k}|} \left(\frac{e^{-i|\mathbf{k}| \cdot (|\mathbf{x}| + t)}}{|\mathbf{x}|} \right) \cdot \frac{\xi(\pi, \varphi, \mathbf{v}_i)}{i|\mathbf{k}|} d|\mathbf{k}| d\varphi =$$

(now I integrate by part with respect to $d|\mathbf{k}|$)

$$\begin{aligned} &= \int \frac{e^{-i\kappa_1 \cdot (|\mathbf{x}| + t)}}{|\mathbf{x}|} \cdot \frac{\xi(\pi, \varphi, \mathbf{v}_i)}{i\kappa_1} d\varphi - \int \frac{e^{-it^{-\alpha} \cdot (|\mathbf{x}| + t)}}{|\mathbf{x}|} \cdot \frac{\xi(\pi, \varphi, \mathbf{v}_i)}{it^{-\alpha}} d\varphi + \\ &+ \int \frac{\kappa_1}{t^\alpha} \int \frac{e^{-i|\mathbf{k}| \cdot (|\mathbf{x}| + t)}}{|\mathbf{x}|} \cdot \xi(\pi, \varphi, \mathbf{v}_i) \cdot \frac{1}{i|\mathbf{k}|^2} \cdot d|\mathbf{k}| d\varphi \end{aligned}$$

from which

$$\begin{aligned} & \int \frac{\kappa_1}{t^\alpha} \int \frac{e^{-i|\mathbf{k}| \cdot (|\mathbf{x}| + t)}}{|\mathbf{x}|} \cdot \frac{\xi(\pi, \varphi, \mathbf{v}_i)}{i|\mathbf{k}|} d|\mathbf{k}| d\varphi = \int \frac{e^{-i\kappa_1 \cdot (|\mathbf{x}| + t)}}{|\mathbf{x}|(|\mathbf{x}| + t)} \cdot \frac{\xi(\pi, \varphi, \mathbf{v}_i)}{|\kappa_1|} d\varphi - \int \frac{e^{-it^{-\alpha} \cdot (|\mathbf{x}| + t)}}{|\mathbf{x}|(|\mathbf{x}| + t)} \cdot \frac{\xi(\pi, \varphi, \mathbf{v}_i)}{t^{-\alpha}} d\varphi + \\ & + \int \frac{\kappa_1}{t^\alpha} \int \frac{e^{-i|\mathbf{k}| \cdot (|\mathbf{x}| + t)}}{|\mathbf{x}|(|\mathbf{x}| + t)} \cdot \frac{\xi(\pi, \varphi, \mathbf{v}_i)}{|\mathbf{k}|^2} d|\mathbf{k}| d\varphi \end{aligned}$$

where each term on the right hand side is bounded by a quantity of order $\frac{t^\alpha}{t^2}$.

Conclusion:

in the region $(1 - \eta')t < |\mathbf{x}| < (1 - \eta)t$ we have that $|\varphi_{\sigma_t, \mathbf{v}_i}(\mathbf{x}, t)| \leq C_{\eta, \eta'} \cdot \frac{t^\alpha}{t^2}$ ($|\mathbf{v}_i| < v^{max} < 1$).

Theorem B4

Taking into account lemma B2, one can prove the existence of

$$s - \lim_{s \rightarrow +\infty} e^{iH_{\sigma_t}s} \int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|s}}{|\mathbf{k}|} \cdot h_{i,j}(\widehat{\mathbf{k}}) \frac{d^3k}{\sqrt{2|\mathbf{k}|}} e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)} \equiv a_{\sigma_t}^{out}(h) \psi_{i,\sigma_t}^{(t)}$$

The vectors $a_{\sigma_t}^{out(in)}(h) \psi_{i,\sigma_t}^{(t)}$ are in $D(H_{\sigma_t})$.

Proof

In order to prove the strong convergence, we check that the following quantity is integrable with respect to s :

$$\left\| \frac{d \left\{ e^{iH_{\sigma_t}s} \int_{\sigma_t}^{\kappa_1} \frac{a(\mathbf{k})e^{i|\mathbf{k}|s}}{|\mathbf{k}|} \cdot h_{i,j}(\widehat{\mathbf{k}}) \frac{d^3k}{\sqrt{2|\mathbf{k}|}} e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)} \right\}}{ds} \right\|$$

Formally :

$$\begin{aligned} \frac{d(e^{iH_{\sigma_t}s} a(h_s) e^{-iH_{\sigma_t}s})}{dt} &= \frac{d(e^{iH_{\sigma_t}s} e^{-iH^{mes}s} a(h) e^{iH^{mes}s} e^{-iH_{\sigma_t}s})}{dt} = \\ &= e^{iH_{\sigma_t}s} \left\{ i \int_{\sigma_t}^{\kappa_1} \tilde{h}_{i,j}(\mathbf{k}) e^{i(|\mathbf{k}|s - \mathbf{k} \cdot \mathbf{x})} \frac{1}{2|\mathbf{k}|^2} d^3k \right\} e^{-iH_{\sigma_t}s} \end{aligned}$$

The formal expressions are well defined from an operatorial point of view in $D(H_{\sigma_t})$.

Having defined $\hat{h}(\mathbf{x}, s) \equiv \left\{ i \int_{\sigma_t}^{\kappa_1} \tilde{h}_{i,j}(\mathbf{k}) e^{i(|\mathbf{k}|s - \mathbf{k} \cdot \mathbf{x})} \frac{1}{2|\mathbf{k}|^2} d^3k \right\}$, I consider the Hilbert inequality:

$$\begin{aligned} &\left\| \frac{d(e^{iH_{\sigma_t}s} a(h) e^{-iH_{\sigma_t}s})}{dt} \psi_{i,\sigma_t}^{(t)} \right\| \leq \\ &\leq \left\| \hat{h}(\mathbf{x}, s) \left(1_{\Gamma_i}(\mathbf{P}) - \chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, s) \right) e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)} \right\| + \left\| \hat{h}(\mathbf{x}, s) \left(\chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, s) - \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right) \right) e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)} \right\| + \\ &+ \left\| \hat{h}(\mathbf{x}, s) \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right) e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)} \right\| \leq \\ &\leq \sup_{\mathbf{x}} \left| \hat{h}(\mathbf{x}, s) \right| \cdot \left\| \left(1_{\Gamma_i}(\mathbf{P}) - \chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, s) \right) e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)} \right\| + \sup_{\mathbf{x}} \left| \hat{h}(\mathbf{x}, s) \right| \cdot \left\| \left(\chi_{\mathbf{v}_i}^{(t)}(\nabla E^{\sigma_t}, s) - \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right) \right) e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)} \right\| + \\ &+ \sup_{\frac{\mathbf{x}}{s} \in \mathbf{J}_{\sigma_t}(\Gamma_i)} \left| \hat{h}(\mathbf{x}, s) \cdot \chi_{\mathbf{v}_i}^{(t)}\left(\frac{\mathbf{x}}{s}, s\right) \right| \cdot \left\| \psi_{i,\sigma_t}^{(t)} \right\| \end{aligned}$$

Because of the results of lemmas B1, B2 and of the estimate in lemma B3, the first two terms on the right hand side are respectively bounded by $C \cdot s^{-\frac{\delta}{12}} \cdot \frac{|\ln \sigma_t|}{s} \cdot t^{-\epsilon}$ ($\delta > 24\epsilon$) and by $C \cdot s^{-1} \cdot s^{-\frac{1}{112}} \cdot s^{2\delta - \frac{3\epsilon}{2}} \cdot (\ln \sigma_t)^2$ (we assume the constraint $2\delta + 3\epsilon < \frac{1}{112}$). As regards the third term, the hypotheses on $G(\mathbf{P})$, $\mathbf{v}_i$ (see the note at pag.69) and the *observation 2* of lemma B3 ensure a vanishing of order $\frac{\sigma_t}{s^2} \cdot t^{-\frac{3\epsilon}{2}}$, for $s \rightarrow +\infty$.

The vectors $a_{\sigma_t}^{out(in)}(h) \psi_{i,\sigma_t}^{(t)}$ belong to $D(H)$.

For each s , $H_{\sigma_t} e^{iH_{\sigma_t}s} a(h_s) e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)}$ is well defined because $a(h_s) \psi_{i,\sigma_t}^{(t)} \subset D(H_{\sigma_t})$. H_{σ_t} is a closed operator. Therefore it is sufficient to prove the convergence, for $s \rightarrow +\infty$, of:

$$\begin{aligned} H_{\sigma_t} e^{iH_{\sigma_t}s} a(h_s) e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)} &= e^{iH_{\sigma_t}s} H_{\sigma_t} a(h_s) e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)} = \\ &= \left\{ e^{iH_{\sigma_t}s} a(h_s) e^{-iH_{\sigma_t}s} E^{\sigma_t}(\mathbf{P}) \psi_{i,\sigma_t}^{(t)} + e^{iH_{\sigma_t}s} [H_{\sigma_t} - H^{mes}, a(h_s)] e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)} + e^{iH_{\sigma_t}s} [H^{e.m.}, a(h_s)] e^{-iH_{\sigma_t}s} \psi_{i,\sigma_t}^{(t)} \right\} \end{aligned}$$

Looking at the first part of the theorem and being $[H^{e.m.}, a(h_s)] = - \int_{\sigma_t}^{\kappa_1} a(\mathbf{k}) e^{i|\mathbf{k}|s} \frac{\tilde{h}_{i,j}(\mathbf{k})}{\sqrt{2|\mathbf{k}|}} d^3k$, each term in the above expression has limit.

Theorem B5

If for each $\mathbf{P}$ in Γ_i and for each $\mathbf{k}$ in $supp \tilde{h}(\mathbf{k})$, $\tilde{h}(\mathbf{k}) = \frac{h_{i,j}(\widehat{\mathbf{k}})}{|\mathbf{k}| \sqrt{2|\mathbf{k}|}} \chi_{\sigma_t}^{\kappa_1}(\mathbf{k})$, it happens that $\mathbf{P} + \mathbf{k} \in \Sigma$, then

$$a_{\sigma_t}^{out(in)}(h) \psi_{i,\sigma_t}^{(t)} = 0$$

Proof

Starting from the spectral decomposition with respect to $\mathbf{P}$ operators, we obtain that $\int a_{\sigma_t}^{out}(\mathbf{k}) \tilde{h}(\mathbf{k}) \left(\psi_{i,\sigma_t}^{(t)} \right)_{\mathbf{P}+\mathbf{k}} d^3k$ is a vector in $H_{\mathbf{P}}$ and that it belongs to the domain of $H_{\mathbf{P},\sigma_t}$. Then the procedure consists in studying the mean value of the positive operator $H_{\mathbf{P},\sigma_t} - E^{\sigma_t}(\mathbf{P})$ on it and in taking into account the condition $|\nabla E^{\sigma_t}|_{\mathbf{P}+\mathbf{q}} < 1$ $\forall \mathbf{q} \in supp \tilde{h}$ (if $\mathbf{P} \in \Sigma$) to estimate $E^{\sigma_t}(\mathbf{P} + \mathbf{k}) - |\mathbf{k}| - E^{\sigma_t}(\mathbf{P}) < 0$ and to conclude that the vector is zero.

References

- [1] E. Nelson. “Interaction of non relativistic particles with a quantized scalar field” J. Math. Phys.,5:1190-1197, 1964.
- [2] J. Froehlich. “On the infrared problem in a model of scalar electrons and massless, scalar bosons.” *Ann. Inst. H. Poincaré*, Sect. A, XIX (1): 1-103, 1973.
- [3] J. Dereziński, C. Gerard “Asymptotic completeness in quantum field theory. Massive Pauli-Fierz hamiltonians” Rev. Math.Phys., 11(1999), 4, 383-450.
- [4] V.Bach, J. Froehlich, I.M. Segal. “Quantum electrodynamics of confined non relativistic particles” *Adv. Math.*, 137(2): 299-395, 1998.